



Organisation for Economic Co-operation and Development

ENV/JM/MONO(2017)15/REV1

Unclassified

English - Or. English

11 April 2023

ENVIRONMENT DIRECTORATE
JOINT MEETING OF THE CHEMICALS COMMITTEE AND THE WORKING PARTY ON
CHEMICALS, PESTICIDES AND BIOTECHNOLOGY

Cancels & replaces the same document of 25 July 2019

**GUIDANCE DOCUMENT NO 263 ON INTEGRATED APPROACHES TO TESTING AND
ASSESSMENT (IATA) FOR SERIOUS EYE DAMAGE AND EYE IRRITATION**

**SERIES ON TESTING AND ASSESSMENT
Number 263
(Second Edition)**

1IMPORTANT NOTE: After the publication of the second edition of the document in July 2019, new Test Guidelines have been published. In particular TG 467 and TG 492B can be used standalone to predict eye hazard potential across the full range of serious eye damage/eye irritation including no irritation responses. This Guidance Document will be updated shortly in a third edition to reflect recent changes.

JT03516409

DRAFT 6 Nov 2023

OECD Environment, Health and Safety Publications

Series on Testing and Assessment

No. 263

GUIDANCE DOCUMENT NO 263 ON INTEGRATED APPROACHES TO TESTING AND
ASSESSMENT (IATA) FOR SERIOUS EYE DAMAGE AND EYE IRRITATION

(Second Edition)

IOMC

INTER-ORGANIZATION PROGRAMME FOR THE SOUND MANAGEMENT OF CHEMICALS

A cooperative agreement among FAO, ILO, UNDP, UNEP, UNIDO, UNITAR, WHO, World Bank and OECD

Environment Directorate
ORGANISATION FOR ECONOMIC CO-OPERATION AND DEVELOPMENT
Paris 2018

About the OECD

The Organisation for Economic Co-operation and Development (OECD) is an intergovernmental organisation in which representatives of 35 industrialised countries in North and South America, Europe and the Asia and Pacific region, as well as the European Commission, meet to co-ordinate and harmonise policies, discuss issues of mutual concern, and work together to respond to international problems. Most of the OECD's work is carried out by more than 200 specialised committees and working groups composed of member country delegates. Observers from several countries with special status at the OECD, and from interested international organisations, attend many of the OECD's workshops and other meetings. Committees and working groups are served by the OECD Secretariat, located in Paris, France, which is organised into directorates and divisions.

The Environment, Health and Safety Division publishes free-of-charge documents in twelve different series: **Testing and Assessment; Good Laboratory Practice and Compliance Monitoring; Pesticides; Biocides; Risk Management; Harmonisation of Regulatory Oversight in Biotechnology; Safety of Novel Foods and Feeds; Chemical Accidents; Pollutant Release and Transfer Registers; Emission Scenario Documents; Safety of Manufactured Nanomaterials; and Adverse Outcome Pathways.** More information about the Environment, Health and Safety Programme and EHS publications is available on the OECD's World Wide Web site (www.oecd.org/chemicalsafety/).

This publication was developed in the IOMC context. The contents do not necessarily reflect the views or stated policies of individual IOMC Participating Organizations.

The Inter-Organisation Programme for the Sound Management of Chemicals (IOMC) was established in 1995 following recommendations made by the 1992 UN Conference on Environment and Development to strengthen co-operation and increase international co-ordination in the field of chemical safety. The Participating Organisations are FAO, ILO, UNDP, UNEP, UNIDO, UNITAR, WHO, World Bank and OECD. The purpose of the IOMC is to promote co-ordination of the policies and activities pursued by the Participating Organisations, jointly or separately, to achieve the sound management of chemicals in relation to human health and the environment.

This publication is available electronically, at no charge.

**For this and many other Environment,
Health and Safety publications, consult the OECD's
World Wide Web site (www.oecd.org/ehs)**

or contact:

**OECD Environment Directorate,
Environment, Health and Safety Division
2, rue André-Pascal
75775 Paris cedex 16
France**

Fax : (33-1) 44 30 61 80

E-mail : ehscont@oecd.org

© OECD 2018

Applications for permission to reproduce or translate all or part of this material should be made to: Head of Publications Service, RIGHTS@oecd.org, OECD, 2 rue André-Pascal, 75775 Paris Cedex 16, France

SUMMARY

This document has two aims:

First, it suggests an Integrated Approach on Testing and Assessment (IATA) for serious eye damage and eye irritation hazard identification, in view of replacing the "sequential testing strategy", which is currently provided in the supplement to OECD TG 405 and which requires adaptation to technical progress.

Second, the document provides key information characteristics of each of the individual information sources comprising the IATA. Furthermore it provides guidance on how and when to integrate existing and/or newly generated information for decision making, including decisions on the need for further testing or final decisions on classification and labelling regarding the potential eye hazard effects of test chemicals.

This Guidance Document was originally approved by the 29th Meeting of the WNT in April 2017, and further updated in 2018 to reflect revisions of Test Guidelines, in particular TG 438 and TG 492.

Table of Contents

SUMMARY	6
List of acronyms	9
1. INTRODUCTION AND SCOPE	1211
2. COMPOSITION OF THE IATA FOR SERIOUS EYE DAMAGE AND EYE IRRITATION	1413
3. INTEGRATION OF IN VITRO TESTS INTO TOP-DOWN AND BOTTOM-UP TESTING APPROACHES	2019
4. DESCRIPTION OF THE ELEMENTS OF THE IATA FOR SERIOUS EYE DAMAGE AND EYE IRRITATION	2523
4.1. Module 1: Existing human data on serious eye damage and eye irritation	2523
4.2. Module 2: In vivo animal data according to OECD TG 405 on serious eye damage and eye irritation	2826
4.2.1. Description and use of the in vivo rabbit eye test method (OECD TG 405) within the IATA	2826
4.2.2. Considerations on the intrinsic characteristics of the in vivo rabbit eye test method	3230
4.3. Module 3: In vitro data from OECD adopted test methods on serious eye damage and eye irritation	3334
4.3.1. Bovine Corneal Opacity and Permeability (BCOP) test method (OECD TG 437)	4136
4.3.2. Isolated Chicken Eye (ICE) test method (OECD TG 438)	4640
4.3.3. Short Time Exposure (STE) test method (OECD TG 491)	5044
4.3.4. Reconstructed human Cornea-like Epithelium Eye Irritation Test (RhCE EIT) (OECD TG 492)	5448
4.3.5. Fluorescein Leakage (FL) test method (OECD TG 460)	5851
4.4. Module 4: Other existing animal data from non-OECD adopted test methods on serious eye damage and eye irritation	7254
4.5. Module 5: Other data from non-OECD adopted alternative test methods on serious eye damage and eye irritation	7254
4.5.1. OECD Guidance Document 160 on the use of histopathology as an additional endpoint	7355
4.5.2. In vitro test methods that underwent validation studies	8062
4.5.3. Promising optimized in vitro test methods	8466
4.6. Module 6: Existing human, in vivo and in vitro data on skin corrosion	8567
4.7. Module 7: Physico-chemical properties (existing, measured or estimated)	8567
4.8. Module 8: Non-testing data on serious eye damage and eye irritation	8769
4.8.1. (Q)SAR, expert systems, grouping and read-across (substances)	8769

4.8.2. Bridging principles and theory of additivity (mixtures)	8974
4.8.3. Overview of non-testing data on serious eye damage and eye irritation	9173
4.9. Module 9: Phases and elements of Weight of Evidence (WoE) approaches	9476
4.9.1. Place/role of WoE in the IATA	9577
4.9.2. Coverage of relevant sources of information	9577
4.9.3. Assessment of data quality	9577
4.9.5. Adequacy of information	9577
4.9.6. Non-testing data	9678
4.9.7. Existing human data	9678
4.9.8. Evaluation of consistency of the data	9779
4.9.10. Assessment of the coverage of relevant parameters and observations	9779
4.9.11. Conclusions of WoE	9779

ANNEX I – EXAMPLE OF MATRIX FOR WEIGHT OF EVIDENCE ANALYSES 11695

ANNEX II – EXAMPLES OF WEIGHT OF EVIDENCE EVALUATIONS 11897

Tables

Table 2.1. Parts and Modules of the IATA	1544
Table 4.1. Regulatory use, applicability, limitations and performance of the OECD adopted in vitro test methods for eye hazard identification.	3432
Table 4.2. Overview of non-OECD adopted test methods useful in supporting eye hazard identification. Note that this is likely to be non-exhaustive. Furthermore it is recommended to check latest status of those methods under discussions at the OECD level.	7456
Table 4.3. Overview of available (Q)SARs for serious eye damage and eye irritation. Note that this list is likely to be non-exhaustive and does not imply endorsement by OECD of any of the listed models for a particular prediction	9072

List of acronyms

A.I.S.E.	International Association for Soaps, Detergents and Maintenance Products
AOP	Adverse Outcome Pathway
BCOP	Bovine Corneal Opacity and Permeability (test)
BfR	German Federal Institute for Risk Assessment
CAMVA	Chorio-Allantoic Membrane Vascular Assay
CAS	Chemical Abstracts Service (number)
Cat.	Category
C&L	Classification and Labelling
CV	Coefficient of Variation
DA	Defined Approach
<u>DAL</u>	<u>Defined Approach for Liquids</u>
DIP	Data interpretation procedure
ET50	Time of exposure to reduce tissue viability of 50%
GD	Guidance Document
GHS	Globally Harmonized System for classification and labelling (UN GHS, 2015)
ECHA	European Chemicals Agency
EIT	Eye Irritation Test
EPAA	European Partnership for Alternative Approaches to Animal Testing
EURL ECVAM	European Union Reference Laboratory for Alternatives to Animal Testing
ESAC	EURL ECVAM Scientific Advisory Committee
EVEIT	Ex Vivo Eye Irritation Test
EU	European Union
FL	Fluorescein Leakage (test)
GLP	Good Laboratory Practices
HCE	Human Corneal Epithelium
HET-CAM	Hen's Egg Test on the Chorio-Allantoic Membrane
HPLC	High Performance Liquid Chromatography
IATA	Integrated Approach on Testing and Assessment
ICCVAM	US Interagency Coordinating Committee on Validation of Alternative Methods
ICE	Isolated Chicken Eye (test)
IRE	Isolated Rabbit Eye (test)

IVIS In Vitro Irritancy Score

JaCVAM Japanese Centre for the Validation of Alternative Methods

JRC European Commission – Directorate General Joint Research Centre

LIS LLBO Irritancy Score

LLBO Laser Light-based Opacimeter

LVET Low Volume Eye Test

MAGAM Multinational Analysis of data from Poison control centres on corrosive Eye lesions of Automatic dishwashing detergent and other detergent and cleaning products

MDCK Madin-Darby Canine Kidney (cells)

MoA Mode of Action

MTT 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide / Thiazolyl blue tetrazolium bromide

NRR Neutral Red Release (test)

OECD Organisation for Economic Co-operation and Development

PorCORA Porcine Ocular Cornea Opacity/Reversibility Assay

QMRF (Q)SAR Model Reporting Format

QPRF (Q)SAR Prediction Reporting Format

(Q)SAR(Quantitative) Structure-Activity Relationship

RBC Red Blood Cell (test)

REACH EU Regulation 1907/2006 on the Registration, Evaluation, Authorisation and restriction of Chemicals

RhCE Reconstructed human Cornea-like Epithelium

SAR Structure-Activity Relationship

SIRC Statens Seruminstitut Rabbit Cornea (cells)

SMI Slug Mucosal Irritation (test)

STE Short Time Exposure (test)

TG Test Guideline

TTT Time-to-Toxicity

UN United Nations

UN GHS Cat. 1 Serious eye damage/irreversible effects on eye

UN GHS Cat. 2/2A Eye irritation/reversible effects on the eye

UN GHS Cat. 2B Mildly irritating to eyes

UN GHS No Cat. No category/no need for classification

UPLC Ultra Performance Liquid Chromatography

US United States

US EPA US Environmental Protection Agency

UVCB Substances of Unknown and Variable Composition and Biologicals

VRM [Validated Reference Method](#)

WNT OECD Working Group of the National Coordinators of the Test Guidelines Programme

WoE Weight of Evidence

DRAFT 6 Nov 2023

1. INTRODUCTION AND SCOPE

1. The objective of the present Guidance Document (GD) is to establish an Integrated Approach on Testing and Assessment (IATA) for hazard identification of serious eye damage and eye irritation potential of test chemicals –(or the absence thereof) that provides adequate information for classification and labelling according to the United Nations Globally Harmonised System (UN GHS, [2015/2023](#)). This GD describes various types of information sources (i.e., physical chemical properties, in vitro and in chemico test methods) and how to use them with the context of an IATA, in order to provide a guidance on using available sources to replace the *in vivo* Draize eye test method or use it only as a last resolution. This GD was originally developed when several in vitro and ex vivo methods became available, giving a choice of assays that are not all providing the same type of information. The GD provides insights into specific aspects of various Test Guidelines and other methods, and can guide users in interpreting data and choosing what other information source/assay type would be conducive to a conclusion on classification. Users should first refer to the data requirements in specific sectors to understand what regulations require. Additionally, in 2023 the UN GHS Chapter 3.3 on eye damage/irritation was updated to include the use of non-animal testing methods for classification of eye hazards in a tiered strategy, which represents a globally harmonized approach for classification (see 3.3.2.8 in UN GHS, 2023).

2. **Serious eye damage** refers to the production of tissue damage in the eye, or serious physical decay of vision, which is not fully reversible ~~(i.e., within 21 days of application in the rabbit test according to OECD TG 405)~~, occurring after exposure of the eye to a test chemical. Test chemicals that have the potential to induce serious eye damage/irreversible effects on the eye are classified as UN GHS Category 1 (UN, [2017/2023](#)). **Eye irritation** refers to the production of changes in the eye, which are fully reversible (i.e., within 21 days in the rabbit test according to OECD TG 405), occurring after exposure of the eye, to a test chemical (UN, [2017/2023](#)). Test chemicals that have the potential to induce eye irritation/reversible effects on the eye are classified as UN GHS Category 2 (UN, [2015/2023](#)). For regulatory authorities requiring more than one classification for reversible eye irritation, Categories 2A and 2B are used, where Category 2A uses the same classification criteria as Category 2 but in which a Category 2B is assigned when the irritant effects triggering Category 2A effects are fully reversible within 7 days of observation (UN, [2015/2023](#)). Finally, test **chemicals not classified** for eye irritation or serious eye damage are defined as those that do not meet the requirements for classification as UN GHS Category 1 or 2 (2A or 2B), and are referred to as UN GHS No Category (No Cat.) (UN, [2015/2023](#)). A test chemical can be an individual (mono- or multi-constituent) substance or a mixture, and represents what is tested without a priori defining the applicability domain for a specific test method.

3. Since 2002, the OECD Test Guideline (TG) 405 on in vivo acute eye irritation and corrosion contains a supplement describing a sequential testing and evaluation strategy for eye irritation/corrosion (OECD, [2012a/2023a](#)). While this supplement is not covered by the OECD Council decision on Mutual Acceptance of Data (MAD), it has provided valuable guidance on how to consider existing information and organise the generation of new testing data on acute eye hazard effects. In its revised version from 2012, the sequential testing and evaluation strategy calls for the use of validated and accepted in vitro and/or ex vivo test methods for identification of serious eye damage (UN GHS Cat. 1), eye irritation (UN GHS Cat. 2 or UN GHS Cat. 2A and 2B), and insufficient eye hazard effects to require classification (i.e., UN GHS No Cat.), before conducting an in vivo animal test. The use of an in vivo animal test is recommended only as a last resort with the purpose of minimising animal use.

4. Since the adoption in 2002 and revision in 2012 of this sequential testing strategy within OECD TG 405, a number of Test Guidelines on in vitro methods have been adopted and/or revised for the identification of test chemicals inducing serious eye damage (UN GHS Cat. 1) or for the identification of test chemicals not requiring classification for eye irritation and serious eye damage hazards (UN GHS No Cat.), notably OECD TG 437, TG 438, TG 460, TG 491, and TG 492, TG 494, and TG 496 (OECD, 2012a, 2013a, 2013b, 2018a, 2015a, 20175b, 2023b, 2023c, 2023d, 2023e, 2023f, 2021, 2023g). OECD TG 492B is a stand-alone in vitro method that has been adopted for the identification of test chemicals not requiring classification (UN GHS No Cat), requiring classification for eye irritation (UN GHS Cat 2) and requiring classification for serious eye damage (UN GHS Cat 1) (OECD 2022a). Furthermore, two rule-based Defined Approaches (DA) have been adopted for the eye hazard identification of non-surfactant liquids according to the three UN GHS hazard categories (TG 467, OECD 2022b). In addition, methods not adopted by the OECD (i.e., not yet validated, not yet accepted by the OECD or implemented within specific country regulatory requirements) may provide further information required by some authorities, e.g. on specific mechanistic insights such as reversibility of effects and effects on the vascular system. The suitability of such data for regulatory purposes needs to be judged on a case by case basis.

5. Updates to the sequential testing and evaluation strategy supplement within OECD TG 405 are therefore required in view of providing guidance on the use, combination and generation of new data, where required. Furthermore, based on the growing experience with the composition and use of IATAs for this specific human health endpoint (UN, 2015, 2023; ECHA, 2015), and the adoption in 2014 of the Guidance Document No. 203 on an Integrated Approach on Testing and Assessment for Skin Corrosion and Irritation (OECD, 2014a, 2017), such revision is timely in order to incorporate current scientific and regulatory considerations and practices for the identification of eye hazards.

6. For these reasons, the OECD Working Group of the National Coordinators for the Test Guidelines (WNT) approved in 2015, a project jointly proposed by the US and the European Commission to develop a Guidance Document on an Integrated Approach on Testing and Assessment (IATA) for serious eye damage and eye irritation. The IATA is composed of well described and characterised "Modules", each of which contain one to several individual information sources of similar type. The strengths and limitations as well as the potential role and contribution of each Module and their individual information sources in the IATA for the identification of serious eye damage, eye irritation and no need for classification are described with the purpose of minimizing the use of animals to the extent possible, while ensuring human safety.

2. COMPOSITION OF THE IATA FOR SERIOUS EYE DAMAGE AND EYE IRRITATION

7. The IATA groups the various individual information sources in "modules" according to the type of information provided. Nine modules were identified as relevant elements of the IATA for eye hazard identification, which can be grouped in three major parts as described in Table 2.1. The different individual information sources associated with each module are described in chapter 4 in a detailed and consistent manner in terms of their applicability, limitations and performance characteristics.

8. The three Parts that guide the assessment of serious eye damage and eye irritation hazards are Part 1 on existing and non-testing data, Part 2 on a weight of evidence analysis, and Part 3 on the generation of new testing data. Under Part 1 of the IATA (existing and non-testing data), existing and available information is retrieved from literature and databases and other reliable sources for Modules 1 to 6, while Module 7 covers physico-chemical properties (primarily pH, which can be existing, measured or estimated) and Module 8 covers non-testing methods, including (Q)SAR, expert systems, grouping and read-across (for substances), and bridging principles and theory of additivity (for mixtures). Part 2 is equivalent to Module 9 and consists of the phases and elements of a weight of evidence (WoE) approach. If the WoE analysis is inconclusive regarding the identification or non-identification of serious eye damage and eye irritation hazard potential, new testing, starting with in vitro methods, needs to be conducted in Part 3 (testing data), in which animal testing is foreseen only as a last resort and after considering the newly obtained in vitro data together with other available information in a second WoE evaluation.

9. A schematic outline of the IATA for eye hazard identification focusing on classification and labelling (C&L) is presented in Figure 2.1. Briefly, the collected existing and non-testing information from Part 1 is evaluated in a WoE approach. If the WoE is conclusive, decision for C&L can be taken accordingly. If it is inconclusive, all available information from the WoE should be considered to formulate a hypothesis of the most likely classification for eye hazard potential of the test chemical, i.e. classified (UN GHS Cat. 1, Cat. 2, Cat. 2A or Cat. 2B), no need for classification (UN GHS No Cat.), or high certainty of not inducing serious eye damage (Non-Cat. 1) (see also chapter 3). This hypothesis will then guide the sequence of prospective testing to e.g. a Top-Down or Bottom-Up approach (Scott et al., 2010) [or Cat. 2 testing methods](#).

Table 2.1. Parts and Modules of the IATA

Part (*)	Modules
Part 1: Existing information, physico-chemical properties and non-testing methods	1. Existing human data on serious eye damage and eye irritation
	2. Existing <i>in vivo</i> animal data according to OECD TG 405 on serious eye damage and eye irritation
	3. Existing <i>in vitro</i> data from OECD adopted test methods on serious eye damage and eye irritation <ul style="list-style-type: none"> a) OECD TG 437 on the BCOP test method b) OECD TG 438 on the ICE test method c) OECD TG 491 on the STE test method d) OECD TG 492 on the RhCE test methods e) OECD TG 460 on the FL test method f) OECD TG 467 on Defined Approaches g) OECD TG 492B on the RhCE test method h) OECD TG 494 on the Vitrigel test method i) OECD TG 496 on the Macromolecular test method
	4. Other existing animal data from non-OECD adopted test methods on serious eye damage and eye irritation
	5. Other data from non-OECD adopted alternative test methods on serious eye damage and eye irritation
	6. Existing data on skin corrosion (human, animal and <i>in vitro</i>)
	7. Physicochemical properties (existing, measured or estimated) such as pH and acid/alkaline reserve
	8. Non-testing data on serious eye damage and eye irritation <ul style="list-style-type: none"> a) Substances: (Q)SAR, expert systems, grouping and read-across b) Mixtures: bridging principles and theory of additivity
Part 2: WoE analysis	9. Phases and elements of WoE approaches
Part 3: New testing	4. Testing on OECD adopted <i>in vitro</i> test methods for serious eye damage and eye irritation
	6. Testing on other non-OECD adopted alternative test methods for serious eye damage and eye irritation
	3. As a last resort, testing on <i>in vivo</i> animal test method according to OECD TG 405 for serious eye damage and eye irritation

* While the three Parts are considered as a sequence, the order of Modules 1 to 8 of Part 1 (here shown in decreasing order of complexity) might be arranged as appropriate. Furthermore, if sufficient and adequate data exist, each module may lead on its own to a classification decision or the absence of classification where relevant, as described in the figure.




Figure 2.1. Detailed IATA for serious eye damage and eye irritation. C&L: Classification and labelling (i.e., UN GHS Cat. 1 or Cat. 2); NC: UN GHS No Category.

* While the three Parts are considered as a sequence, the order of Modules 1 to 8 of Part 1 (here shown in decreasing order of complexity) might be arranged as appropriate. Furthermore, if sufficient and adequate data exist, each module may lead on its own to a classification decision or the absence of classification where relevant, as described in the figure.

Commented [EA1]: TO DO: Remove and/or in the last box when the figure will be updated.

Commented [EA2R1]: PART 3 upper box: Add boxes add the right hand side with "Cat. 2 methods" "OECD adopted *in vitro* test method(s) or DA(s) for eye hazard identification (NC, Cat. 2, Cat. 1) (Module 4)"

- a For example results obtained with other existing in vivo test methods (e.g., the FHSA method 16CFR 1500.42 ([U.S. CPSC, 2003](#)[2015](#))) might be used to derive a final classification, which might include also identification of UN GHS No Category. Furthermore, results obtained with optimized non-OECD adopted test methods (e.g., Isolated Rabbit Eye Test) might be used to identify UN GHS Cat. 1 test chemicals. Finally, negative results obtained with optimized non-OECD adopted test methods might be used in a WoE approach.
- b For example, the application of bridging principles might be used to derive a classification of the tested mixture, which might include also identification of UN GHS No Category. In contrast, results obtained from (Q)SARs might be used in a WoE approach.
- c The use of additional in vitro test methods suitable for identifying UN GHS Cat. 1, based if possible on different mechanisms of action, may be considered in case a negative result is obtained with a first in vitro test method used for this purpose. This is due to the fact that a single in vitro test method aiming at the identification of UN GHS Cat. 1 may not cover all mechanisms of action resulting in serious eye damage (e.g. persistence of effects) and may therefore produce a certain amount of false negatives (see chapters 3 and 4.3).
- d The use of additional OECD adopted in vitro test methods for identifying UN GHS No Cat. may be considered in case a positive result is obtained with a first in vitro test method used for this purpose. This is due to the fact that the currently OECD adopted in vitro test method aiming at the identification of UN GHS No Cat. produce a significant amount of false positives (see chapters 3 and 4.3).
- e In cases where the WoE evaluation in Part 2 indicates that a classification is warranted with a high degree of certainty, testing with an in vitro test method for identification of UN GHS No Cat. may be waived, and the next steps in the strategy should be undertaken.
- f ~~In cases where the WoE evaluation indicates that the test chemical is not UN GHS Cat. 1 or UN GHS No Cat., then OECD UN GHS Cat. 2 testing methods should be utilized. UN GHS Cat. 2 classification is to be considered only in cases where the WoE evaluation indicates that the test chemical is not UN GHS Cat. 1 with a high degree of certainty.~~

10. The structure provided by the three Parts and the information on the nine Modules described in Table 2.1 allow for composing an IATA. Ideally, this IATA should be universally applicable and ensure human safety, while making maximum use of existing data, being resource efficient and eliminating or at least minimising the requirement for animal testing.

11. While the three Parts are considered as a sequence, Modules 1 to 8 of Part 1 might be arranged as appropriate. This will be especially helpful in cases in which information on one or a few Modules cannot be outweighed by any other information, so that a conclusion on the eye hazard potential can be drawn without considering further Modules. Existing information on Modules 1 to 6 can be retrieved by a comprehensive literature and database search. Indeed, in recent years, large databases have become available on the internet, e.g., the European C&L Inventory –and the dissemination site for chemicals registered under REACH-. The search should be performed systematically using search terms such as a CAS (Chemical Abstract Service) number or chemical name. Note that in case relevant information is identified, rights to use this information for regulatory purposes may need to be obtained. Whereas Modules 1 to 5 directly relate to eye hazard, Module 6 requires a different search for in vitro and in vivo skin corrosion data following e.g. its recommended IATA (OECD GD 203, [2014a](#)[2017](#)) that can also impact the final classification of the test chemical.

12. In case the existing information (Modules 1 to 6 within Part 1) does not allow for an unequivocal decision regarding the serious eye damage and eye irritation potential (or the absence thereof) of the substance/mixture, the relevant physico-chemical data and/or non-testing data (i.e., (Q)SAR, expert systems, grouping and read across for substances as well as bridging principles and additivity approach

for mixtures) should be considered. If not retrieved from database searches or available estimates are doubtful, pH and potentially acidity and alkalinity reserve, as well as other physico-chemical parameters may be measured. Regarding Module 8 (non-testing methods), the OECD QSAR Toolbox –may be considered as a starting point to retrieve information as it allows for (i) the retrieval of a first set of existing experimental (physico-chemical and toxicological) data on the target substance(s), (ii) the identification of analogues (for read-across) and retrieval of their existing experimental (physico-chemical and toxicological) data and (iii) the characterisation of these substances with mechanistic and other profilers, including structural alerts for serious eye damage and eye irritation. Further existing data on analogues identified with the Toolbox can then be retrieved by repeating the above literature and database search for these compounds. If data from several (Q)SAR models on a substance are already available and are known to disagree, it may not be helpful to generate other (Q)SAR predictions. If, however no (Q)SAR analysis has been performed, the generation of (Q)SAR information might be helpful to supplement the existing data and come to a conclusion on C&L. Importantly, it is always necessary to carefully consider how well the prediction from each (Q)SAR model falls within the applicability domain of that model.

13. In the analysis of the WoE (Module 9), each data element is characterised for its quality, relevance, coverage (e.g., serious eye damage, eye irritation and/or no need for classification) and associated uncertainty. The decision on inclusion or exclusion of each of the different pieces of existing information is to be based on these parameters (see chapter 4.9). When consistency is seen among "qualified" data elements, WoE may reach a conclusion that the relevant endpoint or information requirement has been sufficiently covered and further testing is not necessary. When on the other hand, insufficient information remains after the "non-qualified" data have been rejected/put aside and/or when the remaining information is inconsistent or contradictory, WoE may lead to a conclusion that further testing is necessary (Part 3 of the IATA), in which case it should also inform on which test(s) to conduct to fill the identified gap(s) (see chapter 3).

14. The WoE assessment needs to be transparently explained and documented to enable a logical flow especially if leading to a final decision/conclusion on classification and labelling. While a WoE approach implies the weighing of each available piece of information on a case by case basis, the modules included in the IATA differ a priori with respect to their intrinsic weight e.g. based on considerations of relevance relating to the species of interest or biological and mechanistic aspects. The following relative a priori weights are nevertheless indicative only and will depend on the quality of the individual data in each specific case. Typically, the relative a priori weights of the modules can be expected to be as follows, based on regulatory acceptance of data when it is of equal quality:

- Good quality and relevant existing human data (Module 1) would be expected to carry the highest weight when the adverse ocular effect and its magnitude can be reliably attributed to the test chemical of interest, however most often such information is not available so that human data on eye hazard effects are generally rather used in a WoE approach.
- This is followed by, with equal weights, in vivo rabbit data according to OECD TG 405 (Module 2) and in vitro data from OECD adopted test methods (Module 3). In particular, it is important to critically appraise the intrinsic characteristics (e.g., uncertainty, variability, drivers of classification) of both the in vivo and the in vitro test methods of Modules 2 and 3 (see chapters 4.2.2 and 4.3).
- Other in vivo animal and in vitro data from non-OECD adopted test methods on serious eye damage and eye irritation (Modules 4 and 5), data indicating skin corrosion (Module 6), physico-chemical information (Module 7) and non-testing methods (Module 8) would typically carry less intrinsic weight.

An example for a simple approach for documenting a WoE evaluation is presented in Annex 1, and examples of evaluations are given for detergents and agrochemical mixtures in annex 2.

15. Before conducting prospective testing for serious eye damage and eye irritation hazard identification, it is strongly recommended to i) consider all existing available test data and ii) generate information whenever possible by means of alternative methods to animal testing such as in vitro methods, (Q)SAR models, grouping or read-across. Evaluating existing data is key to avoid unnecessary animal testing. It can also represent a time and cost efficient way to derive a conclusion on serious eye damage and eye irritation hazard potential, if the available data allow for it.

16. Acknowledging that the applicability of the individual information sources of this IATA to mixtures may vary and that such applicability may depend on the information available in each specific case to be assessed, the IATA is considered applicable to both substances and mixtures. Indeed, data on mixtures can be used for all modules relating to the testing and/or non-testing of eye hazard effects, i.e., modules 1 to 5 and modules 7 to 8 (for details see chapter 4 and paragraph 22).

17. The individual sources of information described in Modules 1 to 8 (Table 2.1) have been characterised as described in chapter 4 and comprise the following information headlines:

- Regulatory use (UN GHS Classification), i.e., the UN GHS Classification that can be derived from individual information sources;
- Validation and regulatory acceptance status;
- Potential role in the IATA;
- Description;
- Scientific basis including Mode of Action (MoA);
- Protocol available;
- Strengths and weaknesses;
- Applicability domain and limitations;
- Predictive capacity, e.g., expressed as sensitivity, specificity and accuracy;
- Reliability, e.g., expressed as within- and between-laboratory reproducibility.

3. INTEGRATION OF IN VITRO TESTS INTO TOP-DOWN AND BOTTOM-UP TESTING APPROACHES

18. ~~It is generally acknowledged that a single in vitro test method is not able to cover the criteria for injury and inflammation addressed by the regulatory adopted in vivo animal test method, i.e. the in vivo rabbit eye test as described in OECD TG 405. Therefore, in order to replace or to reduce the use of the in vivo rabbit eye test, it~~ is recommended to make use of testing strategies that combine the strengths of individual in vitro test methods to address the required ranges of irritation potential and/or chemical classes (Scott et al., 2010). In particular, two tiered testing approaches as shown in Figure 2.1 are recommended for serious eye damage and ~~eye irritation hazard identification~~ **no classification needed**:

- **A Top-Down approach**, starting with in vitro test methods that can identify test chemicals causing serious and/or irreversible eye damage (UN GHS Cat. 1) with low false positive predictions and the highest possible accuracy.
- **A Bottom-Up approach**, starting with in vitro test methods that can identify test chemicals not requiring classification for eye hazard (UN GHS No Cat.) with low false negative predictions and the highest possible accuracy.
- **For suspected eye irritation (UN GHS Cat. 2), see OECD UN GHS Cat. 2 methods.**

19. These tiered testing approaches can be considered as Defined Approaches (DAs) to Testing and Assessment and can be used as a component within the IATA. According to the OECD GD 255 a Defined Approach to testing and assessment is a standardised strategy consisting of a defined set of information sources (in silico and/or in vitro) and a fixed Data Interpretation Procedure (DIP) that is applied to the combined data obtained from the information sources to derive predictions of toxicological effects that can be used either on their own, or together with other information sources within an IATA, to satisfy a specific regulatory need (OECD, 2016). The concept of DIP is taken from the OECD guidance document 34 (OECD, 2005), and is defined in this context as any algorithm for interpreting data from one or more information sources. The defined approach to testing and assessment can be used to support the hazard identification, hazard characterisation and/or safety assessment of chemicals and can be used either on its own to reach a conclusion, or together with other sources of information within an IATA (OECD, 2016). With a view to facilitating the evaluation of IATA in regulatory decision-making, the OECD GD 255 provides with a set of principles and a template for reporting defined approaches to testing and assessment. Such template enables a transparent, structured and harmonised approach to document the defined approaches to testing and assessment. These templates should be used alongside the reporting formats for other IATA components, such as QSARs (OECD, ~~2007~~ **2014a**), grouping and read-across strategies (OECD, 2014b) and non-guideline test methods (OECD, 2014c).

20. All available information and the WoE assessment should be used to formulate a hypothesis of the most likely eye hazard potential of the test chemical, e.g. likelihood to induce serious eye damage (UN GHS Cat. 1) **or likelihood to induce eye irritation (UN GHS Cat. 2)** or likelihood of no need for classification

for eye hazard (UN GHS No Cat.). This hypothesis and the regulatory context under which a decision must be taken should then guide the choice of the prospective testing approach and test methods to be used. The Bottom-Up approach should be followed only when all available collected information and the WoE assessment result in a high a priori probability that the test chemical does not require classification for eye hazard (UN GHS No Cat.). The Top-Down approach, on the other hand should be used when all available collected information and the WoE assessment result in a high a priori probability that the test chemical may induce serious eye damage (UN GHS Cat. 1) ~~or a likelihood for the test chemical to be an eye irritant (UN GHS Cat. 2).~~ Independently of the strategy undertaken, further in vitro testing will depend on the results obtained in the first test following the strategies as shown in Figure 2.1. Only in case of a high likelihood for the test chemical to be an eye irritant (UN GHS Cat. 2) but not to induce serious eye damage (non-Cat. 1), the initial in vitro test method in a top-down approach for identification of UN GHS Cat. 1 may be waived. In this case ~~though~~, the next step in the tiered strategy should be undertaken (i.e., testing with an OECD adopted test method for the identification of UN GHS No Cat. ~~or testing with an OECD adopted test method for the identification of UN GHS Cat. 2, test chemical~~), ~~followed in case of a positive result by a second WoE analyses to determine the most likely eye hazard classification, or the further testing with additional in vitro test methods~~ (see paragraphs 24 to 26).

21. Recommended testing options include the OECD adopted in vitro test methods as described in Module 3 (OECD TG 437 on the BCOP test method, OECD TG 438 on the ICE test method, OECD TG 460 on the FL test method, OECD TG 491 on the STE test method, ~~and~~ OECD TG 492 on the RhCE test methods, ~~OECD TG 467 on the Defined Approaches test methods, OECD TG 492B on the HCE TTT test method, OECD TG 494 on the Vitrigel test method, and OECD TG 496 on the Macromolecular test methods~~). It is generally acknowledged that when the applicability and limitations of the in vitro test methods adopted by the OECD are adequately considered, these methods can, irrespective of the starting point, be used to identify chemicals i) inducing serious eye damage (UN GHS Cat. 1); ~~or ii) inducing eye irritation (UN GHS Cat. 2) or iii) chemicals not requiring classification for eye hazard (UN GHS No Cat.)~~. Note that some test methods such as OECD TG 437 on BCOP, TG 438 on ICE, ~~and~~ TG 491 on STE, ~~and~~ ~~TG 496 on the Macromolecular~~ may be used to initiate the top-down and the bottom-up approaches at the same time, because they are able to provide both UN GHS Cat. 1 and No Cat. predictions, so that the two tiers of the strategy could be covered with one single in vitro assay, provided the test chemical fits the applicability domain and does not fall within the limitations of the test method for each tier (see Table 4.1). However, a test chemical that is neither predicted as UN GHS Cat. 1 nor as UN GHS No Cat. in the bottom-up or top-down approach would require further testing with ~~an OECD Cat. 2 testing method optimised in vitro methods not yet adopted by the OECD (Module 5) as described e.g. in paragraphs 24 to 26. If results obtained with these optimised in vitro methods not yet adopted by the OECD may be used to identify UN GHS Cat. 1 test chemicals, other outcomes can only be used in a new WoE evaluation to be conducted with the newly generated in vitro data together with the existing information (see Figure 2.1).~~ In vivo testing is to be used only as a last resort if still required e.g. by regulators to establish a definitive classification (UN GHS Cat. 1, Cat. 2 (Cat. 2A or Cat. 2B if applicable) or No Cat.).

22. The currently adopted in vitro test methods (OECD TGs 437, 438, 460, 491, ~~and~~ 492, 492B, 494, ~~and~~ 496) are applicable to both substances and mixtures. ~~TG 467 DAL-1 is not applicable to mixtures but is applicable to non-surfactant neat liquids. TG 467 DAL-2 is applicable to non-surfactant neat liquids, liquids and solids dissolved in water.~~ Indeed, OECD TGs 437 (BCOP), 438 (ICE), 491 (STE) and 492 (RhCE) have undergone evaluation studies conducted on both substances and mixtures (OECD ~~2013a, 2013b, 2013c, 2013d, 2013e, 2013f, 2013g, 2013h, 2013i, 2013j, 2013k, 2013l, 2013m, 2013n, 2013o, 2013p, 2013q, 2013r, 2013s, 2013t, 2013u, 2013v, 2013w, 2013x, 2013y, 2013z, 2014a, 2014b, 2014c, 2014d, 2014e, 2014f, 2014g, 2014h, 2014i, 2014j, 2014k, 2014l, 2014m, 2014n, 2014o, 2014p, 2014q, 2014r, 2014s, 2014t, 2014u, 2014v, 2014w, 2014x, 2014y, 2014z, 2015a, 2015b, 2015c, 2015d, 2015e, 2015f, 2015g, 2015h, 2015i, 2015j, 2015k, 2015l, 2015m, 2015n, 2015o, 2015p, 2015q, 2015r, 2015s, 2015t, 2015u, 2015v, 2015w, 2015x, 2015y, 2015z, 2016a, 2016b, 2016c, 2016d, 2016e, 2016f, 2016g, 2016h, 2016i, 2016j, 2016k, 2016l, 2016m, 2016n, 2016o, 2016p, 2016q, 2016r, 2016s, 2016t, 2016u, 2016v, 2016w, 2016x, 2016y, 2016z, 2017a, 2017b, 2017c, 2017d, 2017e, 2017f, 2017g, 2017h, 2017i, 2017j, 2017k, 2017l, 2017m, 2017n, 2017o, 2017p, 2017q, 2017r, 2017s, 2017t, 2017u, 2017v, 2017w, 2017x, 2017y, 2017z, 2018a, 2018b, 2018c, 2018d, 2018e, 2018f, 2018g, 2018h, 2018i, 2018j, 2018k, 2018l, 2018m, 2018n, 2018o, 2018p, 2018q, 2018r, 2018s, 2018t, 2018u, 2018v, 2018w, 2018x, 2018y, 2018z, 2019a, 2019b, 2019c, 2019d, 2019e, 2019f, 2019g, 2019h, 2019i, 2019j, 2019k, 2019l, 2019m, 2019n, 2019o, 2019p, 2019q, 2019r, 2019s, 2019t, 2019u, 2019v, 2019w, 2019x, 2019y, 2019z, 2020a, 2020b, 2020c, 2020d, 2020e, 2020f, 2020g, 2020h, 2020i, 2020j, 2020k, 2020l, 2020m, 2020n, 2020o, 2020p, 2020q, 2020r, 2020s, 2020t, 2020u, 2020v, 2020w, 2020x, 2020y, 2020z, 2021a, 2021b, 2021c, 2021d, 2021e, 2021f, 2021g, 2021h, 2021i, 2021j, 2021k, 2021l, 2021m, 2021n, 2021o, 2021p, 2021q, 2021r, 2021s, 2021t, 2021u, 2021v, 2021w, 2021x, 2021y, 2021z, 2022a, 2022b, 2022c, 2022d, 2022e, 2022f, 2022g, 2022h, 2022i, 2022j, 2022k, 2022l, 2022m, 2022n, 2022o, 2022p, 2022q, 2022r, 2022s, 2022t, 2022u, 2022v, 2022w, 2022x, 2022y, 2022z, 2023a, 2023b, 2023c, 2023d, 2023e, 2023f, 2023g, 2023h, 2023i, 2023j, 2023k, 2023l, 2023m, 2023n, 2023o, 2023p, 2023q, 2023r, 2023s, 2023t, 2023u, 2023v, 2023w, 2023x, 2023y, 2023z, 2024a, 2024b, 2024c, 2024d, 2024e, 2024f, 2024g, 2024h, 2024i, 2024j, 2024k, 2024l, 2024m, 2024n, 2024o, 2024p, 2024q, 2024r, 2024s, 2024t, 2024u, 2024v, 2024w, 2024x, 2024y, 2024z, 2025a, 2025b, 2025c, 2025d, 2025e, 2025f, 2025g, 2025h, 2025i, 2025j, 2025k, 2025l, 2025m, 2025n, 2025o, 2025p, 2025q, 2025r, 2025s, 2025t, 2025u, 2025v, 2025w, 2025x, 2025y, 2025z, 2026a, 2026b, 2026c, 2026d, 2026e, 2026f, 2026g, 2026h, 2026i, 2026j, 2026k, 2026l, 2026m, 2026n, 2026o, 2026p, 2026q, 2026r, 2026s, 2026t, 2026u, 2026v, 2026w, 2026x, 2026y, 2026z, 2027a, 2027b, 2027c, 2027d, 2027e, 2027f, 2027g, 2027h, 2027i, 2027j, 2027k, 2027l, 2027m, 2027n, 2027o, 2027p, 2027q, 2027r, 2027s, 2027t, 2027u, 2027v, 2027w, 2027x, 2027y, 2027z, 2028a, 2028b, 2028c, 2028d, 2028e, 2028f, 2028g, 2028h, 2028i, 2028j, 2028k, 2028l, 2028m, 2028n, 2028o, 2028p, 2028q, 2028r, 2028s, 2028t, 2028u, 2028v, 2028w, 2028x, 2028y, 2028z, 2029a, 2029b, 2029c, 2029d, 2029e, 2029f, 2029g, 2029h, 2029i, 2029j, 2029k, 2029l, 2029m, 2029n, 2029o, 2029p, 2029q, 2029r, 2029s, 2029t, 2029u, 2029v, 2029w, 2029x, 2029y, 2029z, 2030a, 2030b, 2030c, 2030d, 2030e, 2030f, 2030g, 2030h, 2030i, 2030j, 2030k, 2030l, 2030m, 2030n, 2030o, 2030p, 2030q, 2030r, 2030s, 2030t, 2030u, 2030v, 2030w, 2030x, 2030y, 2030z, 2031a, 2031b, 2031c, 2031d, 2031e, 2031f, 2031g, 2031h, 2031i, 2031j, 2031k, 2031l, 2031m, 2031n, 2031o, 2031p, 2031q, 2031r, 2031s, 2031t, 2031u, 2031v, 2031w, 2031x, 2031y, 2031z, 2032a, 2032b, 2032c, 2032d, 2032e, 2032f, 2032g, 2032h, 2032i, 2032j, 2032k, 2032l, 2032m, 2032n, 2032o, 2032p, 2032q, 2032r, 2032s, 2032t, 2032u, 2032v, 2032w, 2032x, 2032y, 2032z, 2033a, 2033b, 2033c, 2033d, 2033e, 2033f, 2033g, 2033h, 2033i, 2033j, 2033k, 2033l, 2033m, 2033n, 2033o, 2033p, 2033q, 2033r, 2033s, 2033t, 2033u, 2033v, 2033w, 2033x, 2033y, 2033z, 2034a, 2034b, 2034c, 2034d, 2034e, 2034f, 2034g, 2034h, 2034i, 2034j, 2034k, 2034l, 2034m, 2034n, 2034o, 2034p, 2034q, 2034r, 2034s, 2034t, 2034u, 2034v, 2034w, 2034x, 2034y, 2034z, 2035a, 2035b, 2035c, 2035d, 2035e, 2035f, 2035g, 2035h, 2035i, 2035j, 2035k, 2035l, 2035m, 2035n, 2035o, 2035p, 2035q, 2035r, 2035s, 2035t, 2035u, 2035v, 2035w, 2035x, 2035y, 2035z, 2036a, 2036b, 2036c, 2036d, 2036e, 2036f, 2036g, 2036h, 2036i, 2036j, 2036k, 2036l, 2036m, 2036n, 2036o, 2036p, 2036q, 2036r, 2036s, 2036t, 2036u, 2036v, 2036w, 2036x, 2036y, 2036z, 2037a, 2037b, 2037c, 2037d, 2037e, 2037f, 2037g, 2037h, 2037i, 2037j, 2037k, 2037l, 2037m, 2037n, 2037o, 2037p, 2037q, 2037r, 2037s, 2037t, 2037u, 2037v, 2037w, 2037x, 2037y, 2037z, 2038a, 2038b, 2038c, 2038d, 2038e, 2038f, 2038g, 2038h, 2038i, 2038j, 2038k, 2038l, 2038m, 2038n, 2038o, 2038p, 2038q, 2038r, 2038s, 2038t, 2038u, 2038v, 2038w, 2038x, 2038y, 2038z, 2039a, 2039b, 2039c, 2039d, 2039e, 2039f, 2039g, 2039h, 2039i, 2039j, 2039k, 2039l, 2039m, 2039n, 2039o, 2039p, 2039q, 2039r, 2039s, 2039t, 2039u, 2039v, 2039w, 2039x, 2039y, 2039z, 2040a, 2040b, 2040c, 2040d, 2040e, 2040f, 2040g, 2040h, 2040i, 2040j, 2040k, 2040l, 2040m, 2040n, 2040o, 2040p, 2040q, 2040r, 2040s, 2040t, 2040u, 2040v, 2040w, 2040x, 2040y, 2040z, 2041a, 2041b, 2041c, 2041d, 2041e, 2041f, 2041g, 2041h, 2041i, 2041j, 2041k, 2041l, 2041m, 2041n, 2041o, 2041p, 2041q, 2041r, 2041s, 2041t, 2041u, 2041v, 2041w, 2041x, 2041y, 2041z, 2042a, 2042b, 2042c, 2042d, 2042e, 2042f, 2042g, 2042h, 2042i, 2042j, 2042k, 2042l, 2042m, 2042n, 2042o, 2042p, 2042q, 2042r, 2042s, 2042t, 2042u, 2042v, 2042w, 2042x, 2042y, 2042z, 2043a, 2043b, 2043c, 2043d, 2043e, 2043f, 2043g, 2043h, 2043i, 2043j, 2043k, 2043l, 2043m, 2043n, 2043o, 2043p, 2043q, 2043r, 2043s, 2043t, 2043u, 2043v, 2043w, 2043x, 2043y, 2043z, 2044a, 2044b, 2044c, 2044d, 2044e, 2044f, 2044g, 2044h, 2044i, 2044j, 2044k, 2044l, 2044m, 2044n, 2044o, 2044p, 2044q, 2044r, 2044s, 2044t, 2044u, 2044v, 2044w, 2044x, 2044y, 2044z, 2045a, 2045b, 2045c, 2045d, 2045e, 2045f, 2045g, 2045h, 2045i, 2045j, 2045k, 2045l, 2045m, 2045n, 2045o, 2045p, 2045q, 2045r, 2045s, 2045t, 2045u, 2045v, 2045w, 2045x, 2045y, 2045z, 2046a, 2046b, 2046c, 2046d, 2046e, 2046f, 2046g, 2046h, 2046i, 2046j, 2046k, 2046l, 2046m, 2046n, 2046o, 2046p, 2046q, 2046r, 2046s, 2046t, 2046u, 2046v, 2046w, 2046x, 2046y, 2046z, 2047a, 2047b, 2047c, 2047d, 2047e, 2047f, 2047g, 2047h, 2047i, 2047j, 2047k, 2047l, 2047m, 2047n, 2047o, 2047p, 2047q, 2047r, 2047s, 2047t, 2047u, 2047v, 2047w, 2047x, 2047y, 2047z, 2048a, 2048b, 2048c, 2048d, 2048e, 2048f, 2048g, 2048h, 2048i, 2048j, 2048k, 2048l, 2048m, 2048n, 2048o, 2048p, 2048q, 2048r, 2048s, 2048t, 2048u, 2048v, 2048w, 2048x, 2048y, 2048z, 2049a, 2049b, 2049c, 2049d, 2049e, 2049f, 2049g, 2049h, 2049i, 2049j, 2049k, 2049l, 2049m, 2049n, 2049o, 2049p, 2049q, 2049r, 2049s, 2049t, 2049u, 2049v, 2049w, 2049x, 2049y, 2049z, 2050a, 2050b, 2050c, 2050d, 2050e, 2050f, 2050g, 2050h, 2050i, 2050j, 2050k, 2050l, 2050m, 2050n, 2050o, 2050p, 2050q, 2050r, 2050s, 2050t, 2050u, 2050v, 2050w, 2050x, 2050y, 2050z, 2051a, 2051b, 2051c, 2051d, 2051e, 2051f, 2051g, 2051h, 2051i, 2051j, 2051k, 2051l, 2051m, 2051n, 2051o, 2051p, 2051q, 2051r, 2051s, 2051t, 2051u, 2051v, 2051w, 2051x, 2051y, 2051z, 2052a, 2052b, 2052c, 2052d, 2052e, 2052f, 2052g, 2052h, 2052i, 2052j, 2052k, 2052l, 2052m, 2052n, 2052o, 2052p, 2052q, 2052r, 2052s, 2052t, 2052u, 2052v, 2052w, 2052x, 2052y, 2052z, 2053a, 2053b, 2053c, 2053d, 2053e, 2053f, 2053g, 2053h, 2053i, 2053j, 2053k, 2053l, 2053m, 2053n, 2053o, 2053p, 2053q, 2053r, 2053s, 2053t, 2053u, 2053v, 2053w, 2053x, 2053y, 2053z, 2054a, 2054b, 2054c, 2054d, 2054e, 2054f, 2054g, 2054h, 2054i, 2054j, 2054k, 2054l, 2054m, 2054n, 2054o, 2054p, 2054q, 2054r, 2054s, 2054t, 2054u, 2054v, 2054w, 2054x, 2054y, 2054z, 2055a, 2055b, 2055c, 2055d, 2055e, 2055f, 2055g, 2055h, 2055i, 2055j, 2055k, 2055l, 2055m, 2055n, 2055o, 2055p, 2055q, 2055r, 2055s, 2055t, 2055u, 2055v, 2055w, 2055x, 2055y, 2055z, 2056a, 2056b, 2056c, 2056d, 2056e, 2056f, 2056g, 2056h, 2056i, 2056j, 2056k, 2056l, 2056m, 2056n, 2056o, 2056p, 2056q, 2056r, 2056s, 2056t, 2056u, 2056v, 2056w, 2056x, 2056y, 2056z, 2057a, 2057b, 2057c, 2057d, 2057e, 2057f, 2057g, 2057h, 2057i, 2057j, 2057k, 2057l, 2057m, 2057n, 2057o, 2057p, 2057q, 2057r, 2057s, 2057t, 2057u, 2057v, 2057w, 2057x, 2057y, 2057z, 2058a, 2058b, 2058c, 2058d, 2058e, 2058f, 2058g, 2058h, 2058i, 2058j, 2058k, 2058l, 2058m, 2058n, 2058o, 2058p, 2058q, 2058r, 2058s, 2058t, 2058u, 2058v, 2058w, 2058x, 2058y, 2058z, 2059a, 2059b, 2059c, 2059d, 2059e, 2059f, 2059g, 2059h, 2059i, 2059j, 2059k, 2059l, 2059m, 2059n, 2059o, 2059p, 2059q, 2059r, 2059s, 2059t, 2059u, 2059v, 2059w, 2059x, 2059y, 2059z, 2060a, 2060b, 2060c, 2060d, 2060e, 2060f, 2060g, 2060h, 2060i, 2060j, 2060k, 2060l, 2060m, 2060n, 2060o, 2060p, 2060q, 2060r, 2060s, 2060t, 2060u, 2060v, 2060w, 2060x, 2060y, 2060z, 2061a, 2061b, 2061c, 2061d, 2061e, 2061f, 2061g, 2061h, 2061i, 2061j, 2061k, 2061l, 2061m, 2061n, 2061o, 2061p, 2061q, 2061r, 2061s, 2061t, 2061u, 2061v, 2061w, 2061x, 2061y, 2061z, 2062a, 2062b, 2062c, 2062d, 2062e, 2062f, 2062g, 2062h, 2062i, 2062j, 2062k, 2062l, 2062m, 2062n, 2062o, 2062p, 2062q, 2062r, 2062s, 2062t, 2062u, 2062v, 2062w, 2062x, 2062y, 2062z, 2063a, 2063b, 2063c, 2063d, 2063e, 2063f, 2063g, 2063h, 2063i, 2063j, 2063k, 2063l, 2063m, 2063n, 2063o, 2063p, 2063q, 2063r, 2063s, 2063t, 2063u, 2063v, 2063w, 2063x, 2063y, 2063z, 2064a, 2064b, 2064c, 2064d, 2064e, 2064f, 2064g, 2064h, 2064i, 2064j, 2064k, 2064l, 2064m, 2064n, 2064o, 2064p, 2064q, 2064r, 2064s, 2064t, 2064u, 2064v, 2064w, 2064x, 2064y, 2064z, 2065a, 2065b, 2065c, 2065d, 2065e, 2065f, 2065g, 2065h, 2065i, 2065j, 2065k, 2065l, 2065m, 2065n, 2065o, 2065p, 2065q, 2065r, 2065s, 2065t, 2065u, 2065v, 2065w, 2065x, 2065y, 2065z, 2066a, 2066b, 2066c, 2066d, 2066e, 2066f, 2066g, 2066h, 2066i, 2066j, 2066k, 2066l, 2066m, 2066n, 2066o, 2066p, 2066q, 2066r, 2066s, 2066t, 2066u, 2066v, 2066w, 2066x, 2066y, 2066z, 2067a, 2067b, 2067c, 2067d, 2067e, 2067f, 2067g, 2067h, 2067i, 2067j, 2067k, 2067l, 2067m, 2067n, 2067o, 2067p, 2067q, 2067r, 2067s, 2067t, 2067u, 2067v, 2067w, 2067x, 2067y, 2067z, 2068a, 2068b, 2068c, 2068d, 2068e, 2068f, 2068g, 2068h, 2068i, 2068j, 2068k, 2068l, 2068m, 2068n, 2068o, 2068p, 2068q, 2068r, 2068s, 2068t, 2068u, 2068v, 2068w, 2068x, 2068y, 2068z, 2069a, 2069b, 2069c, 2069d, 2069e, 2069f, 2069g, 2069h, 2069i, 2069j, 2069k, 2069l, 2069m, 2069n, 2069o, 2069p, 2069q, 2069r, 2069s, 2069t, 2069u, 2069v, 2069w, 2069x, 2069y, 2069z, 2070a, 2070b, 2070c, 2070d, 2070e, 2070f, 2070g, 2070h, 2070i, 2070j, 2070k, 2070l, 2070m, 2070n, 2070o, 2070p, 2070q, 2070r, 2070s, 2070t, 2070u, 2070v, 2070w, 2070x, 2070y, 2070z, 2071a, 2071b, 2071c, 2071d, 2071e, 2071f, 2071g, 2071h, 2071i, 2071j, 2071k, 2071l, 2071m, 2071n, 2071o, 2071p, 2071q, 2071r, 2071s, 2071t, 2071u, 2071v, 2071w, 2071x, 2071y, 2071z, 2072a, 2072b, 2072c, 2072d, 2072e, 2072f, 2072g, 2072h, 2072i, 2072j, 2072k, 2072l, 2072m, 2072n, 2072o, 2072p, 2072q, 2072r, 2072s, 2072t, 2072u, 2072v, 2072w, 2072x, 2072y, 2072z, 2073a, 2073b, 2073c, 2073d, 2073e, 2073f, 2073g, 2073h, 2073i, 2073j, 2073k, 2073l, 2073m, 2073n, 2073o, 2073p, 2073q, 2073r, 2073s, 2073t~~

applicability of the Test Guideline to a specific category of mixture, based on the chemistry and/or physico-chemical property, the Test Guideline should not be used for that specific category. While agrochemical formulations have successfully been tested using the EpiOcular EIT (OECD TG 492) for the identification of UN GHS No Cat., the BCOP [OP-KIT](#) (OECD TG 437) was found to be under-predictive for identification of UN GHS Cat. 1 agrochemical mixtures (Kolle et al., 2015). This could be due to the fact that the majority of the tested agrochemical mixtures (n=19 out of 21) in this study were classified in vivo based on persistence of effects only, which is a type of effect known not to be identified per se by the currently OECD adopted in vitro methods aiming at the identification of UN GHS Cat. 1 (see chapter 4.3).

23. The applicability domain and performance of the OECD adopted individual test methods are described in their respective Modules in chapter 4 and are summarized in Table 4.1. When using adopted in vitro test methods, it is critical to ensure using the most appropriate OECD TG for the specific purpose and chemical to be tested. In particular, the applicability domain plays an important role in the choice of the test method to be used. For example, test methods having the highest possible accuracy for the chemical class tested should be preferentially used. Similar care should be taken in case optimised in vitro test methods not yet adopted by the OECD are used and information on applicability domain is available on these test methods. In addition, it is important [that the test methods employed reflect mechanisms of eye irritation that are relevant to humans, and do not necessarily need to reflect or cover the mechanisms that take place in the rabbit test \(apart from where there is overlap\) \(Clippinger et al., 2021\). Furthermore, it is important to take into account the mechanistic insights provided by in vitro test methods, and how those cover the mechanisms taking place in the in vivo test method \(see paragraph 24 and chapter 4.2.2\).](#) Finally, when using two or more test methods (see paragraphs 24 and 25), the conditional independence of these test methods should be considered (Adriaens et al. 2017a; Hoffman et al., 2008). This can help to decide which test methods to be included in the Top-Down / Bottom-Up approaches and optimise the overall performance of the approach chosen.

24. One of the problems associated with the originally proposed two-tier Top-Down/Bottom-Up testing strategy (Scott et al., 2010) is that a default UN GHS Cat. 2 classification after only testing in two test methods would generate a significant number of false negative (Cat. 1 underclassified as Cat. 2) and false positive (No Cat. overclassified as Cat. 2) results (see Table 4.1). Currently accepted methods for identifying UN GHS Cat. 1, like BCOP and ICE, underpredict 14-48% of the in vivo Cat. 1 chemicals, mostly those inducing persistent effects without occurrence of initial high level injuries (classified in vivo based only on persistence of effects). Therefore, since the single in vitro test methods aiming at the identification of UN GHS Cat. 1 may not cover all mechanisms of action resulting in serious eye damage (e.g. persistence of effects) and can produce a certain amount of false negatives (see chapter 4.3), the use of additional in vitro test methods suitable for identifying UN GHS Cat. 1 based, if possible, on different mechanism of actions, may be considered in case a negative result is obtained with a first in vitro test method used for this purpose. Moreover, it is clear that due to the very high sensitivity required by regulatory authorities for accepting the use of in vitro test methods to identify chemicals not requiring hazard classification and labelling for serious eye damage/eye irritation (UN GHS No Cat.), their specificity will never go beyond 60-80% (the highest the specificity, the more limited the applicability). RhCE test methods, ICE and STE are those showing the best accuracy for identifying UN GHS No Cat. chemicals and their specificity is only 63-81% with already a few false negatives being obtained (sensitivity around 95%). In such a scenario, several methods capable of identifying UN GHS No Cat. chemicals with very high sensitivity will need to be combined to increase the overall specificity of the testing strategy to acceptable values. Therefore, since the currently OECD adopted in vitro test methods aiming at the identification of UN GHS No Cat. produce a significant amount of false positives (see chapter 4.3), the use of additional OECD adopted in vitro test methods for identifying UN GHS No Cat. may be considered in case a positive result is obtained with a first in vitro test method used for this purpose. In addition to the OECD adopted in vitro test method, the use of optimised non-OECD adopted in vitro test methods and/or endpoints, as described in chapter 4.5, may be used to identify UN GHS Cat. 1 test chemicals, or to be

considered as complementary information in a WoE evaluation for the identification of other eye hazard categories.

25. Some examples on the use of the proposed testing strategy approach have been reported. In particular for antimicrobial and cleaning products, the US Environmental Protection Agency (EPA) recommends the use of a testing approach for determining the appropriate eye hazard classification and labelling. The strategy, which represents a replacement of the in vivo data requirement, utilizes a decision tree involving the use of the BCOP, EpiOcular time-to-toxicity (ET50) and Cytosensor Microphysiometer test methods (US EPA, 2015). Other potential ways of combining in vitro tests methods in testing strategies based on the concept of the Bottom-up and Top-down approaches have been investigated by Kolle et al. (2011), and Hayashi et al. (2012a, 2012b) and Adriaens et al (2017a). Both ~~These studies showed that~~ combinations of methods in Defined Approaches (DAs) can lead to better predictions as compared to each individual test method on its own. Kolle et al. (2011) combined EpiOcular™ EIT and BCOP in a two-tier Bottom-up/Top-Down test strategy and Hayashi et al. (2012b) combined EpiOcular™ EIT, BCOP, STE and HET-CAM in a two-stage Bottom-Up tiered approach. In Adriaens et al. (2017a) two-tiered and three-tiered strategies combined an RhCE test method (EpiOcular™ EIT or SkinEthic™ EIT) at the bottom (identification No Cat) in combination with the BCOP LLBO (two-tiered strategy) or BCOP and SMI (three-tiered strategy) at the top (identification Cat 1).

26. Similar performance was obtained for the Top-down and Bottom-up approach. Based on the data presented in these two publications, Schaeffer and co-workers (2014) showed that specificity for identifying UN GHS No Cat. chemicals can increase substantially by combining in a test strategy several methods able to identify UN GHS No Cat. test chemicals (including both OECD adopted and non-adopted test methods). This occurs as a result of multiple methods complementing each other by correctly identifying different sets of UN GHS No Cat. chemicals. Interestingly the authors show that the increase in specificity of the test strategy as compared to the individual methods is not accompanied by a significant decrease in sensitivity due to the very high sensitivity already displayed by all of these methods on their own. Furthermore, the accuracy for the identification of UN GHS Cat. 2 by default at the end of the strategy would be significantly improved (Schaeffer et al., 2014). One of the aspects that should be considered when combining different test methods in a tiered strategy, is the dependence between the test methods. Ideally, the test methods that are combined in a testing strategy should be independent to improve the predictive performance. Hoffmann and colleagues (2008) and Adriaens et al (2017a) demonstrated that when two similar methods and thus highly conditional dependent methods are included in a testing strategy, the predictive performance of the strategy will not improve.

27. ~~Indeed, it is generally recognized that w~~When using the Top-Down and Bottom-Up approaches, the main difficulty lies in predicting the middle category of irritancy (e.g. UN GHS Cat. 2, Cat. 2A or Cat. 2B). ~~There now exists OECD test methods for identifying UN GHS Cat. 2 (but not Cat. 2A or Cat. 2B). The optional use of additional in vitro test method(s) may be helpful in improving the prediction of UN GHS Cat. 2. This could be due to an increased accuracy of a default Cat. 2 prediction by decreasing the number of false positives when identifying No Cat. and by decreasing the number of false negatives when identifying Cat. 1. Nevertheless further work and data are needed to reach an acceptable level of predictivity for UN GHS Cat. 2 chemicals. For example, conduct of statistical modelling (taking into consideration the conditional independence of the test methods as described in paragraph 23) may allow to define the desirable performances of the in vitro test methods that may, when combined in e.g., Defined Approaches to testing and assessment, and used within the appropriate applicability domain and regulatory context, be used to derive a default UN GHS Cat. 2 prediction if neither a UN GHS Cat. 1 nor a UN GHS No Cat. prediction can be made.~~

28. The in vivo rabbit eye test (OECD TG 405) should be conducted only as a last resort after all the existing information in Part 1 of the IATA has been considered, and after the in vitro testing in Part 3 has been conducted and evaluated in an additional WoE

Commented [EA3]: Or delete these paragraph as suggested by US FDA

evaluation together with the existing data. The in vivo animal test, if e.g. required by regulators, should be considered after in vitro testing only when:

- i) the test chemical is not directly identified as UN GHS Cat. 1, [UN GHS Cat. 2](#) or as UN GHS No Cat. by the in vitro test methods and WoE assessment cannot conclude with high enough confidence if the test chemical is Cat. 1, Cat. 2 (or Cat. 2A or Cat. 2B, if applicable), or No Cat. Depending on country-specific regulatory requirements, test methods not yet adopted by the OECD should also be considered both prospectively and in the WoE evaluation.
- ii) the test chemical cannot be tested with the in vitro test methods due to the limitations of the test methods or when falling outside of the applicability domain of the test method.

4. DESCRIPTION OF THE ELEMENTS OF THE IATA FOR SERIOUS EYE DAMAGE AND EYE IRRITATION

29. The individual sources of information to be used in Modules 1 to 8 (Table 2.1) and the elements of the weight of evidence evaluation of the collected information to be conducted in Module 9, within the IATA for the hazard identification of serious eye damage and eye irritation potential of test chemicals (or the absence thereof), have been characterised and are described below.

4.1. Module 1: Existing human data on serious eye damage and eye irritation

30. Existing human data include historical data that should be taken into account when evaluating intrinsic hazards of test chemicals. New testing in humans for hazard identification purposes is not acceptable for ethical reasons. Existing data can be obtained from single or repeated exposure(s) from case reports, poison information centres, medical clinics, occupational experience, epidemiological studies and volunteer studies. Note however, that the availability of the epidemiological studies for this endpoint is likely to be rare and the quality often questionable. The quality and relevance for hazard assessment of the existing human data should be critically reviewed. For example, in occupational studies with mixed exposure it is important that the test chemical causing serious eye damage or eye irritation is accurately identified. There may also be a significant level of uncertainty in human data due to poor reporting and lack of specific information on exposure. However, well-documented existing human data from various sources can provide useful information on serious eye damage and eye irritation hazard potential of a test chemical, sometimes for a range of exposure levels. For example, the MAGAM study, first conducted by a retrospective collection of data from poison control centres (in Germany, Austria and Switzerland) between 1998 and 2007 (Stürer et al., 2010), led to the MAGAM II prospective study conducted in 2013-2015, in which the criteria for data collection were defined prior to the start of the study to ensure high quality of the collected data from the poison centres. MAGAM II represents a multicentre study aimed at collecting and evaluating data on human eye exposures to detergents and maintenance products from a number of poison control centres, which includes, among other, information on severity of effects, duration and outcome.

31. Good quality and relevant human data can be used to determine serious eye damage or eye irritation potential of a test chemical and have precedence over other data. However, absence of reported ocular incidents in humans is no evidence in itself for no classification. The usefulness of the human data on adverse ocular effects will depend on the extent to which the effect, and its magnitude, can be reliably attributed to the test chemical of interest. Examples of how existing human data can be used in hazard classification for ocular effects have been reported (MAGAM II study; ECETOC, 2002). In humans, an ophthalmic examination by a physician would reveal a decay of vision. If it is not transient but persistent it implies classification in Category 1. If the discrimination

between Category 1 and Category 2 is not obvious, then Category 1 might be chosen; however, other types of information may be generated e.g. by performing in vitro testing, to support the final hazard classification conclusion.

Module 1 – Existing human data on serious eye damage and eye irritation	
General description	
Regulatory use (UN GHS classification)	Human data from accident (e.g. from hospitals) or poison control centre databases can provide evidence for UN GHS Cat. 1 and Cat. 2 classification. However, absence of incidents is not in itself evidence for no classification as exposures are generally unknown or uncertain.
Validation & regulatory acceptance status	Existing human data include historical data that should be taken into account when evaluating intrinsic hazards of test chemicals. New testing in humans for hazard identification purposes is not acceptable for ethical reasons.
Potential role in the IATA	Good quality and relevant human data would be expected to have precedence over other data when the adverse ocular effect and its magnitude can be reliably attributed to the test chemical of interest, however most often such information is not available so that human data on eye hazard effects are generally rather used in a WoE approach. Furthermore, absence of incidence in humans does not necessarily overrule in vitro data or existing animal data of good quality that are positive. Finally, if the discrimination between Category 1 and Category 2 is not obvious other types of information may be generated e.g. by performing in vitro testing, to support the final hazard classification conclusion.
Description	Ophthalmic examination by a physician revealing a decay of vision, which if not transient but persistent, implies classification in Category 1. If the discrimination between Category 1 and Category 2 is not obvious, then Category 1 might be chosen.
Scientific basis incl. MoA	All MoA are potentially covered.
Protocol available	No standard protocol is available. However, efforts have been undertaken to standardize collection of data from poison centres (e.g., MAGAM II study). Existing human data might be derived (e.g., in occupational, consumer, transport, or emergency response scenarios) from single or repeated exposure(s) from case reports, poison information centres, medical clinics, occupational experience, epidemiological studies and volunteer studies. Note however, that the availability of the epidemiological studies for this endpoint is likely to be rare and the quality often questionable.

Strengths and weaknesses	<p><u>Strengths</u></p> <ul style="list-style-type: none"> - Relevant data as obtained directly from the species of interest (humans). - Examples available on how existing human data can be used (MAGAM II study; ECETOC, 2002). <p><u>Weaknesses</u></p> <ul style="list-style-type: none"> - Not standardised. - Mostly based on accidental/uncontrolled exposure often in combination with co-exposure, leading to a high level of uncertainty. - Sufficient data to evaluate the actual exposure (duration and dose) might not be always available. - Data might be incomplete, insufficient or inaccurate. - Data on the reversibility of the effect might not be always available. - Data on additional, potentially confounding factors (e.g., purity, health status of the affected person, additional exposures) might not be available. - No UN GHS criteria for C&L based on human data are available.
Identification of UN GHS Cat. 1 and Cat. 2	
Applicability domain and limitations	<p><u>Applicability domain</u></p> <ul style="list-style-type: none"> - All test chemicals for which a clear and direct effect on the eye can be concluded from the available data (note that the exposure scenario and chemical identity (needed for concluding on a direct effect) are often not clearly defined in data obtained from accidental exposure). <p><u>Limitations</u></p> <ul style="list-style-type: none"> - Rarely available and, if available, not often with the necessary quality to be used on its own for C&L decisions, so that it is most often used in a WoE evaluation with other existing data to make C&L decisions.
Predictive capacity	The usefulness of human data will depend on the amount and quality of the available information. It is often associated with a high level of uncertainty due to lack of critical information such as chemical identity and purity, exposure scenario (dose and duration), health status of the persons exposed and/or the reported symptoms.
Reliability	Difficult to assess due to uncontrolled exposures (dose and timings) and reporting, although efforts exist to improve quality of data collection from poison centres (e.g., MAGAM II study).

4.2. Module 2: In vivo animal data according to OECD TG 405 on serious eye damage and eye irritation

4.2.1. Description and use of the in vivo rabbit eye test method (OECD TG 405) within the IATA

32. The OECD TG 405 (OECD, ~~2012~~2023a) on in vivo Acute Eye Irritation/Corrosion testing recommends the use of rabbits as preferred species. It was originally adopted in 1981, and revised in 2002 to include i) a supplement on a sequential testing and evaluation strategy for eye hazard identification, ii) use of dermal irritation/corrosion test data to predict eye corrosion prior to considering the conduct of an in vivo animal test and iii) the possibility to rinse solid materials from the eyes 1 hour after treatment (instead of the previous 24 hours). In 2012 the TG was further revised to include the possibility to use topical anaesthetics, systemic analgesics, and humane endpoints during in vivo animal testing to avoid most or all pain and distress without affecting the outcome of the test. The revision of 2017 includes that topical anaesthetics, systemic analgesics, and humane endpoints should be routinely used during in vivo animal testing.

33. In vivo animal testing should not be considered until all available data relevant to the eye hazard potential (or absence thereof) of a test chemical have been evaluated in a WoE analysis according to the present IATA, and the necessary prospective in vitro testing conducted as described in chapter 3 (see also Figure 2.1). This includes conducting a study on the skin corrosion potential of the test chemical before the in vivo animal test on serious eye damage and eye irritation. In cases where the in vivo animal test is required, it is recommended that it is performed in a sequential manner using initially one animal. If the results of this initial test with one animal indicate the test chemical to induce serious eye damage, further testing should not be performed. If serious eye damage is not observed in the initial test, the irritant or negative response should be confirmed using up to two additional animals. However, if an irritant effect was observed in the initial test the confirmatory test should be conducted in one animal at a time, rather than exposing the two additional animals simultaneously. It may not be necessary to test a total of three animals if classification of the test chemical can be achieved using only two animals. Finally, due consideration should be made to the intrinsic characteristics of the in vivo rabbit eye test method as described in chapter 4.2.2.

Module 2 – In vivo animal data on serious eye damage and eye irritation according to OECD TG 405	
General description	
Regulatory use (UN GHS classification)	Classification decision on serious eye damage (UN GHS Cat. 1), eye irritation (UN GHS Cat. 2, Cat. 2A and Cat. 2B), and no need for classification (UN GHS No Cat.).
Validation & regulatory status	The animal test method adopted in OECD TG 405 was never formally validated but has been the historical regulatory test method for testing serious eye damage and eye irritation hazard potential of test chemicals.
Potential role in the IATA	<p>In case in vivo animal test data of adequate quality are available, these should carry a certain intrinsic weight in the context of a WoE analysis, taking into consideration the critical appraisal of the intrinsic characteristics (e.g., uncertainty, variability, drivers of classification) of the in vivo rabbit test method as described in chapter 4.2.2.</p> <p>The in vivo animal test should be conducted only as a last resort after i) considering results from the in vivo and/or in vitro skin corrosion test method, ii) considering and evaluating all available information relevant to the serious eye damage and eye irritation hazard potential of the test chemical in a WoE analysis (Parts 1 and 2 of the IATA as described in Table 2.1 and Figure 2.1), and iii) considering the results obtained with prospective in vitro testing (Modules 3 and 5 of Part 3 of the IATA as described in Table 2.1 and Figure 2.1).</p>
Description	The test chemical is applied in a single dose (0.1 mL for liquids or an amount corresponding to a volume of 0.1 mL or a weight of not more than 100 mg for solids, pastes and particulate substances) to the conjunctival sac of one of the eyes of the experimental animal (albino rabbit is the preferred animal species) whereas the untreated eye serves as control. Degree of serious eye damage and eye irritation is assessed by scoring lesions to cornea (opacity), iris and conjunctiva (redness and oedema) at specific time intervals and the duration of the study should be sufficient to evaluate the reversibility or irreversibility of the effects. The UN GHS classification is based on the mean tissue scores obtained (as recorded per animal) at 24, 48 and 72 hours after exposure, and on the reversibility or irreversibility of effects observed for up to 21 days. Other effects in the eye and possible adverse systemic effects are also assessed to provide a complete evaluation of the effects.
Scientific basis incl. MoA	<p>The test method allows assessing:</p> <ul style="list-style-type: none"> - Serious eye damage, i.e. the production of tissue damage in the eye, or serious physical decay of vision, which is not fully reversible within 21 days of application, and

	<p>- Eye irritation, i.e. the production of changes in the eye, which are fully reversible within 21 days of application.</p> <p>The mechanisms by which such effects are produced and detected in the in vivo animal test method are multiple and depend on the type of chemicals tested. Regarding the cornea, these may include as a first step cell disruption, denaturation and swelling of collagen. This is followed in a second step by the production and release of intermediates that initiate the process of inflammation, causing the oedema in corneal stroma and invasion of leukocytes. In a third step, regeneration of epithelium may gradually occur resulting in decreased corneal opacity. Finally in some cases as a fourth step, destruction of cornea and stromal ulceration may occur 2 to 3 weeks after injury, mediated by hydrolytic enzymes coupled with inadequate collagen synthesis. When the cornea has reepithelialised or when the corneal stroma becomes totally vascularised, corneal ulceration ceases (Berta, 1992, Pfister, 1983; McCulley, 1987; Lemp, 1974).</p> <p>Other mechanisms of injury detected by the test method include i) inflammation of the conjunctivae in which the dilation of blood vessels can cause redness, and the increased effusion of water can cause oedema/chemosis, and ii) secretion of mucous leading to an increase in discharge. Iritis can also occur either as a direct effect or as a secondary reaction due to the corneal injury. Once iris is inflamed, infiltration of fluids can follow which can affect visual acuity accompanied by symptoms of itching, burning and stinging. Finally, other possible mechanisms of injury covered by the in vivo animal test method include: i) loss of corneal innervations, ii) tear film abnormalities due to injury to the lacrimal glands, iii) intense pain, lacrimation, and blepharospasm due to direct stimulation of free nerve endings located in the epithelium of the cornea and conjunctival lining, iv) neurogenic inflammation.</p> <p>Irreversible effects may occur when the damage extends to and beyond the corneal endothelium causing corneal perforation that may cause permanent loss of vision. Other persistent effects include discolouration of the cornea by a dye chemical, adhesion, pannus, and interference with the function of the iris or any other effects that impair sight which do not reverse within the test period.</p>
<p>Protocol available</p>	<p>OECD TG 405 (2012/2023a) based on the scoring system developed by Draize and co-workers (1944).</p>
<p>Strengths and weaknesses</p>	<p><u>Strengths</u></p> <ul style="list-style-type: none"> - The in vivo animal test method reflects all possible modes of action of serious eye damage and eye irritation reactions present in rabbit eyes. - It formed the basis for the GHS classification system, and can therefore identify the entire spectrum of eye effects i.e., UN GHS No Cat., Cat. 2 (and the UN GHS Cat. 2A and 2B), and Cat. 1. - Reversibility and/or persistence of effects can be directly observed.

	<p><u>Weaknesses</u></p> <ul style="list-style-type: none"> - Not formally validated. - The possibility of concluding Cat. 1 on the basis of a single eye exposure, which, depending on the type of effect(s) observed, can be associated with a very high uncertainty. - Reproducibility compromised by e.g.: <ul style="list-style-type: none"> • Subjectivity in the allocation of the ocular tissue scores; • Unclear duration and amount of exposure of the test chemical in the rabbit eyes which can vary depending on the properties of the test chemical (solid, paste or liquid) as well as the blinking and tear reflex from the animal (Prinsen, 2006); • Differences in animal behaviour (e.g., lacrimation, blinking, etc) which can lead to differences in reactions even before scoring of effects takes place (Prinsen, 2006);; • Absence (or presence) of post-treatment care. - For certain test chemicals (e.g., solid, sticky), blinking can result in mechanical damage, contributing to a higher degree of irritation (Prinsen, 2006). - Enclosure of test materials in the conjunctival cul-de-sac in combination with mechanical damage can lead to exacerbation of effects and secondary inflammation not directly caused by the test chemical (Prinsen, 2006). - The animal type of exposure does not reflect human accidental exposure scenarios (Wilhelmus, 2001). - There are differences in physiology and sensitivity to test chemicals between rabbit and human eyes. - Poor correlation was found between rabbit and human mean time to clear (Freeberg et al., 1986b) - The testing can be very painful to the rabbits.
Identification of UN GHS Cat. 1, Cat. 2 (A and B) and No Cat.	
Applicability domain and limitations	The test method is applicable to substances, mixtures and aerosols.
Predictive capacity	Differences in physiology and sensitivity exist between rabbit and human eyes, and the in vivo rabbit test has been shown to be in general more sensitive to hazard chemicals than the eyes of humans (Roggeband et al., 2000; Gershbein and McDonald, 1977; Wilhelmus, 2001; ILSI, 1996). More recently this has been shown to be particularly the case for test chemicals inducing serious eye damage (Ishii et al., 2013).

Reliability	<p>Taking into account the animal within-test variability only, at least 11% of chemicals classified in vivo as UN GHS Cat. 1 could be equally identified as Cat. 2 by the in vivo rabbit eye test itself, and about 12% of the Cat. 2 chemicals could be equally identified as non-classified chemicals (Adriaens et al., 2014).</p> <p>If variability between repeat studies were taken into account, the observed concordance of UN GHS classifications when considering a unified Cat. 2 classification was found to be of 65.2 % (15/23) (Barroso et al., 2017). If Cat. 2A and Cat. 2B are considered as different classifications, the observed concordance of UN GHS classifications was found to be 56.5 % (13/23). Finally concordance of the same main driver of classification (see chapter 4.2.2) was found to occur for 39.1 % (9/23) of the chemicals (Barroso et al., 2017).</p> <p>Furthermore, evaluation of public data from ECHA online dossiers on 9,782 in vivo rabbit eye studies on 3,420 unique substances, showed that the most reproducible outcomes were for the negative results (94% reproducible) and for chemicals inducing serious eye damage (73% reproducible), whereas there was a 10% chance of a non-irritant evaluation be given after a prior severe-irritant result based on the UN GHS classification criteria (Luechtefeld et al., 2016).</p>
-------------	--

4.2.2. Considerations on the intrinsic characteristics of the in vivo rabbit eye test method

34. In a recent study by Adriaens et al. (2014), co-sponsored by the European Commission and Cosmetics Europe, statistical resampling of in vivo rabbit test data (according to OECD TG 405) on 2134 chemicals demonstrated an overall probability of at least 11% that chemicals classified as UN GHS Cat. 1 by the in vivo rabbit eye test could be equally identified as UN GHS Cat. 2 and of about 12% for UN GHS Cat. 2 chemicals to be equally identified as UN GHS No Cat. simply due to the test method's inherent within-test variability. On the other hand, the chances for UN GHS No Cat. and UN GHS Cat. 2 test chemicals to be predicted in a higher UN GHS Category - was found to be negligible (< 1%). Altogether, these observations suggest that the classification criteria of the in vivo rabbit eye test are highly sensitive on their own (Adriaens et al., 2014). Taking into account the variability between repeat studies, an overall concordance of 65.2 % (15/23) was found for the UN GHS Cat. 1, a unified Cat. 2 and No Cat. classifications (Barroso et al., 2017). If Cat. 2A and Cat. 2B were considered as different classifications, an overall concordance of 56.5 % (13/23) was found for the UN GHS Cat. 1, Cat. 2A, Cat. 2B and No Cat. classifications (Barroso et al., 2017). An evaluation of public data from ECHA online dossiers on 9,782 in vivo rabbit eye studies on 3,420 unique substances, further showed that the most reproducible outcomes were for negative results (94% reproducible) and chemicals inducing serious eye damage (73% reproducible), whereas there was a 10% chance of a non-irritant evaluation be given after a prior serious eye damage result based on the UN GHS classification criteria (Luechtefeld et al., 2016). Considering these results, it is probably not achievable to develop in vitro test methods with no false negatives.

35. The results of the study by Adriaens and colleagues (Adriaens et al., 2014) also indicate that the persistence and severity of corneal opacity play an equally important role in the classification of a chemical as UN GHS Cat. 1, whereas corneal opacity and conjunctival redness are the most important tissue effects that determine the classification of UN GHS Cat. 2 eye irritants. In a study co-sponsored by the European Commission and

Cosmetics Europe, a further evaluation was performed to establish which of the in vivo rabbit eye test drivers of classification are most important from a regulatory point of view for driving UN GHS classification (Barroso et al., 2017). For this purpose a in vivo rabbit eye test Reference Database was compiled containing 681 independent in vivo studies on 634 individual chemicals representing a wide range of chemical classes. The analyses confirmed the previous results from Adriaens et al. (2014) by showing that corneal opacity is the most important tissue effect driving Cat. 1 classification (including corneal opacity mean ≥ 3 (days 1-3, severity) and corneal opacity persistence on day 21 in the absence of severity), whereas Cat. 2 classification was found to be mostly driven by corneal opacity mean ≥ 1 and conjunctival redness mean ≥ 2 . Based on the evidence presented in the manuscript, the authors identified a number of key criteria that should be taken into consideration when selecting reference chemicals for the development, evaluation and/or validation of alternative methods and/or strategies for serious eye damage/eye irritation testing. Such understanding is critical for properly assessing their predictive capacity and limitations. Furthermore, a critical revision of the UN GHS decision criteria for the classification of chemicals from the in vivo rabbit eye test data was proposed by Adriaens et al. (2014) and Barroso et al. (2017) based on the results of their analyses of historical in vivo data.

4.3. Module 3: In vitro data from OECD adopted test methods on serious eye damage and eye irritation

36. The present chapter provides a description of the in vitro information sources and their use within the IATA for serious eye damage and eye irritation. A number of in vitro test methods have been adopted since 2009 to identify i) test chemicals inducing serious eye damage (UN GHS Cat. 1), ii) [test chemicals inducing eye irritation \(UN GHS Cat 2\)](#) and/or iii) test chemicals not requiring classification for eye irritation or serious eye damage (UN GHS No Cat.). An overview of the regulatory use, applicability, limitations and performance of the OECD adopted in vitro test methods for eye hazard identification is given in Table 4.1 and Table 4.2. When using classification systems other than the UN GHS, the appropriate regulatory authorities should be consulted.

37. As compared to the in vivo rabbit eye test (OECD TG 405, [2012/2023a](#)), the currently available in vitro information sources do not directly assess effects on the iris although it should be noted that effects on the iris are of lesser importance for classification of test chemicals according to UN GHS (Adriaens et al., 2014; Barroso et al., 2017). Furthermore, the neurogenic components that drive tear film production are usually not present in the in vitro test methods. As a consequence, when compared with an in vivo rabbit eye study, application of a test chemical in the absence of this protective barrier might be expected to cause an increase in false positive outcomes. [Current in vitro test methods also lack limbal stem cells which can differentiate into corneal epithelial cells that are important for the reversibility of damage. The absence of limbal epithelium in a rabbit model resulted in corneal erosions and vascularization \(Huang and Tseng, 1991\)](#). Finally, the adopted in vitro test methods do not allow for an assessment of the potential for systemic toxicity associated with ocular exposure. However, these effects are typically predicted from other acute toxicity test methods, and may not be relevant for the many consumer products that are formulated with well characterized raw materials that have been already characterised for the presence/absence of systemic toxicity effects.

Table 4.1. Regulatory use, applicability, limitations and performance of the OECD adopted in vitro test methods for eye hazard identification – identification of UN GHS Category 1 and/or UN GHS No Category.

	BCOP (OECD TG 437)	ICE (OECD TG 438)	STE (OECD TG 491)	RhCE (OECD TG 492)	FL (OECD TG 460)	Vitrigel (OECD TG 494)	Macromolecular (OECD TG 496)
<u>Identification of UN GHS Category 1</u>							
Applicability	Substances and mixtures	Substances and mixtures	Substances, multi-constituent substances and mixtures that are dissolved or uniformly suspended for at least 5 minutes	Not applicable	Water soluble substances and mixtures	<u>Not applicable</u>	<u>Solid and liquid chemicals whose 10% solution/dispersion has a pH range of 4 ≤ pH ≤ 9</u>
Limitations	Alcohols and ketones risk overprediction	Alcohols risk <u>overprediction</u>	No other specific limitation reported	Not applicable	Strong acids and bases, cell fixatives, highly volatile test chemicals, coloured and viscous test chemicals, solid chemicals suspended in liquid that have tendency to precipitate	<u>Not applicable</u>	<u>Intensely coloured chemicals, chemicals which caused salting-out precipitation, high concentrations of some surfactants, and highly volatile chemicals</u>
Accuracy*	<u>OP-KIT: 79% (150/191)</u>	<u>86% (120/140)</u> <u>83% (142/172)</u>	83% (104/125)	Not applicable	77% (117/151)	<u>Not applicable</u>	<u>75%^a (66.5/89)</u>

Unclassified

	<u>LLBO: 78%^a (n=145)</u>						
False positive rate* (1-specificity)	<u>OP-KIT: 25% (32/126)</u> <u>LLBO: 21%^a (n=89)</u>	<u>7% (9/127)</u> 6% (7/113)	1% (1/86)	Not applicable	7% (7/103)	<u>Not applicable</u>	<u>19%^a (13.2/69)</u>
False negative rate* (1-sensitivity)	<u>OP-KIT: 14% (9/65)</u> <u>LLBO: 24%^a (n=56)</u>	<u>47% (21/45)</u> 48% (13/27)	51% (20/39)	Not applicable	56% (27/48)	<u>Not applicable</u>	<u>46%^a (9.3/20)</u>

DRAFT 6 Nov 2023

	<u>BCOP</u> (OECD TG 437)	<u>ICE</u> (OECD TG 438)	<u>STE</u> (OECD TG 491)	<u>RhCE</u> (OECD TG 492)	<u>FL</u> (OECD TG 460)	<u>Vitrigel</u> (OECD TG 494)	<u>Macromolecular</u> (OECD TG 496)
<u>Identification of UN GHS No Category</u>							
Applicability	Substances and mixtures	Substances and mixtures	Substances and mixtures Substances, multi-constituent substances and mixtures that are dissolved or uniformly suspended for at least 5 minutes	Substances and mixtures. Test chemicals interfering with MTT measurement (by i.e., colour interference or reduction of MTT) require the use of appropriate controls or HPLC-UPLC analysis if colour incompatibility with MTT higher than 60% is reported.	Not applicable	<u>Mono-constituent and multi-constituent substances, substances of unknown or variable composition, complex reaction products or biological materials (UVCBs)</u> <u>Any test chemical showing pH > 5 and keeping dissolution or homogeneous dispersion for at least three minutes in a 2.5% w/v concentration in culture medium.</u> <u>Test chemicals that do not dissolve readily can be tested after using one of the following techniques: a) mix mechanically using a vortex mixer, b)</u>	<u>Substances and mixtures</u> <u>Solid and liquid chemicals whose 10% solution/dispersion with 4 ≤ pH ≤ 9</u>

						sonication, and/or c) heating to a maximum temperature of 70°C.	
Limitations	Due to high false positive rates, BCOP should not be the first choice method to initiate a Bottom-up approach	Anti-fouling organic solvent-containing paints may be under-predicted For solid materials leading to a GHS No Cat. outcome, a second testing run is recommended	Highly volatile substances with vapour pressure > 6 kPa (at 25°C) Solid chemicals (substances and mixtures) other than surfactants and mixtures of surfactants only Mixtures containing substances with vapour pressure > 6kPa <u>that do not dissolve in mineral oil, or that do not form stable suspensions for at least 5 minutes</u> may risk underpredictions		Not applicable	<u>Test chemical preparations of both solids and liquids showing acidity (pH ≤ 5) and rapid phase separation are not in the applicability domain</u>	<u>intensely coloured chemicals, chemicals which caused salting-out precipitation, high concentrations of some surfactants, and highly volatile chemicals</u>
Accuracy*	<u>OP-KIT: 69% (135/196)</u> <u>LLBO: 83%^a (n=145)</u>	88% (161/184) <u>82% (125/152)</u>	90% (92/102)	<u>VRM1: 80% (n=112)</u> <u>VRM2: 84% (n=200)</u>	Not applicable	<u>78% (73/93)</u>	<u>75%^a (67.0/89)</u>
False positive rate* (1-specificity)	<u>OP-KIT: 69% (61/89)</u> <u>LLBO: 45%^a (n=41)</u>	24% (20/83) <u>33% (26/79)</u>	19% (9/48)	<u>VRM1: 37% (n=55)</u> <u>VRM2: 28% (n=103)</u>	Not applicable	<u>30% (10/33)</u>	<u>41%^a (17.7/43)</u>

False negative rate* (1-sensitivity)	OP-KIT: 0% (0/107) LLBO: 6% ^a (n=104)	3% (3/101) 4% (1/73)	2% (1/54)	VRM1: 4% (n=57) VRM2: 5% (n=97)	Not applicable	17% (10/60)	9% ^a (4.3/46)
---	---	------------------------------------	-----------	------------------------------------	----------------	-------------	--------------------------

* As reported in the respective Test Guidelines.

^a The proportion in the tables are based on weighted calculation. For each chemical, all results were taken into account and a correction factor was applied so that all chemicals had the same weight (weight of 1).

~~* As reported in the respective Test Guidelines. # VRM1: EpiOcular™ EIT and VRM2: SkinEthic™ HCE EIT, respectively~~

BCOP: Bovine Corneal Opacity and Permeability; FL: Fluorescein Leakage; ICE: Isolated Chicken Eye; RhCE: Reconstructed human Cornea-like Epithelium; STE: Short Time Exposure.

Table 4.2. Regulatory use, applicability, limitations and performance of the OECD adopted in vitro test methods for eye hazard identification – full replacement defined approaches and test method

	<u>Defined Approach DAL-1</u> (OECD TG 467)	<u>Defined Approach DAL-2</u> (OECD TG 467)	<u>SkinEThic™ HCE TTT</u> (OECD TG 492B)
<u>Applicability</u>	<u>Applicable to neat liquids.</u>	<u>Applicable to neat liquids, liquids and solids dissolved in water.</u>	<u>Substances and mixtures</u> <u>Test chemicals interfering with MTT measurement (by i.e., colour interference or reduction of MTT) require the use of appropriate controls or HPLC-UPLC analysis.</u>
<u>Limitations</u>	<u>Not applicable for surfactants, solids and liquid mixtures, liquid UVCBs and liquid multi-constituent substances.</u>	<u>Not applicable for surfactants and solids dispersed in water.</u>	
<u>Balanced accuracy</u>	<u>DAL-1a: 68.7% (n=94)</u> <u>DAL-1b: 75.0% (n=86)</u>	<u>74.3% (n=164)</u>	<u>74.4% (n=151)</u>
<u>Performance UN GHS Cat. 1</u>	<u>DAL-1a: 76.5% TP, 23.5% UP, 0.0% FN (n=17)</u> <u>DAL-1b: 76.5% TP, 23.5% UP, 0.0% FN (n=17)</u>	<u>81.2% TP, 17.6% UP, 1.2% FN (n=17)</u>	<u>79.2% TP, 20.8% UP, 0.0% FN (n=50)</u>
<u>Performance UN GHS Cat. 2</u>	<u>DAL-1a: 59.1% TP, 27.3% OP, 13.6% FN (n=22)</u> <u>DAL-1b: 68.7% TP, 30.4% OP, 0.9% FN (n=23)</u>	<u>56.3% TP, 30.2% OP, 13.5% FN (n=24)</u>	<u>69.2% TP, 18.3% OP, 12.5% FN (n=44)</u>
<u>Performance UN GHS No Cat.</u>	<u>DAL-1a: 70.5% TN, 29.5% FP (n=55)</u> <u>DAL-1b: 79.7% TN, 20.3% FP (n=46)</u>	<u>85.3% TN, 14.7% FP (n=123)</u>	<u>74.9% TN, 25.1% FP (n=57)</u>

TP: True Positive; UP: Under-Prediction; OP: Over-Prediction;

^a DAL-1 with VRM1, ^b DAL-1 with VRM2

Note: the performance statistics are based on weighted calculations. For each chemical, all results were taken into account and a correction factor was applied so that all chemicals had the same weight (weight of 1).

DRAFT 6 Nov 2023

4.3.1. Bovine Corneal Opacity and Permeability (BCOP) test method (OECD TG 437)

38. The OECD TG 437 on the BCOP test method was originally adopted in 2009 and updated in 2013 (OECD, 2013a; OECD, 2013e2013b2023b). In 2020, the TG was updated to allow the use of a laserlight-based opacitometer (LLBO) which was shown to have similar performance as the OP-KIT opacitometer used in the validation of the BCOP test method (OECD, 2023b). The BCOP OP-KIT test method underwent two retrospective validation studies by the US Interagency Coordinating Committee on the Validation of Alternative Methods (ICCVAM), in conjunction with the European Union Reference Laboratory for Alternatives to Animal Testing (EURL ECVAM) and the Japanese Center for the Validation of Alternative Methods (JaCVAM), in 2006 and 2010 (ICCVAM, 2006; ICCVAM, 2010b). In the two evaluations, the BCOP was endorsed as a scientifically valid test method for use as a screening test to identify UN GHS Cat. 1 substances and mixtures (ICCVAM, 2006, 2010b; ESAC, 2007). Furthermore the second evaluation study and a further retrospective evaluation of the in vitro and in vivo dataset used in the validation study concluded that the BCOP test method can also be used to identify UN GHS No Category substances and mixtures (ICCVAM 2010b; OECD, 2013e2013b). The data set was enlarged in the CON4E1 project with 80 chemicals, 67/80 chemicals were unique to this project (Verstraelen et al., 2017; Adriaens et al., 2017b). From these evaluations it was concluded that the BCOP test method can correctly identify test chemicals (both substances and mixtures) inducing serious eye damage (UN GHS Cat. 1) as well as those not requiring classification for eye hazard (UN GHS No Category), and it was therefore endorsed as scientifically valid for both purposes.

Commented [EA4]: OECD No 189

Module 3 – In vitro data: Bovine Corneal Opacity and Permeability (BCOP) test method (OECD TG 437)	
General Description	
Regulatory use (UN GHS classification)	Identification of i) test chemicals inducing serious eye damage (UN GHS Cat. 1), and ii) test chemicals not requiring classification for eye hazard (UN GHS No Cat.).
Validation & regulatory acceptance status	Validated and adopted as OECD TG 437-.
Potential role in the IATA	While the BCOP test method is not considered valid as a full-replacement replacement for the in vivo rabbit eye test, it can be used for regulatory classification and labelling (Figure 2.1) to identify, without further testing: <ul style="list-style-type: none"> - test chemicals inducing serious eye damage (UN GHS Cat. 1); and - test chemicals that do not require classification for eye hazard (UN GHS No Cat.). BCOP can be used to initiate a Top-Down approach. However, since it has a high overprediction rate for test chemicals that do not require

	<p>classification for eye hazard (69% <u>for OP-KIT and 45% for LLBO</u>), it should not be the first choice to initiate a Bottom-Up approach. Other validated and accepted in vitro methods with similar high sensitivity but higher specificity should be used instead as first tier of a Bottom-Up approach (e.g., OECD TG 492).</p> <p>A test chemical that is neither predicted as UN GHS Cat. 1 nor as UN GHS No Cat. with the BCOP test method would require additional <u>testing with a UN GHS Cat. 2 method and/or</u> WoE evaluation with other existing information and if still needed additional testing (in vitro and/or in vivo) as a last resort to establish a definitive classification (see Figure 2.1).</p>												
<p>Description</p>	<p>The BCOP test method is an organotypic ex vivo assay that makes use of isolated corneas from the eyes of freshly slaughtered cattle placed on corneal holders. Test chemicals are applied to the epithelial surface of the cornea by addition to the anterior chamber of the corneal holder. Damage by the test chemical is assessed by quantitative measurements of:</p> <ul style="list-style-type: none"> - Corneal opacity changes, measured as the amount of light transmission through the cornea with the help of an opacitometer ; and - Permeability, measured as the amount of sodium fluorescein dye that passes from the medium in the anterior chamber of the corneal holder, across the full thickness of the cornea, to the medium in the posterior chamber, detected with the help of a visible light spectrophotometer. <p>Both measurements are used to calculate an In vitro Irritancy Score (IVIS) <u>or LLBO Irritancy Score (LIS)</u>.</p> <table border="1" data-bbox="427 1227 1046 1512"> <thead> <tr> <th><u>Prediction</u></th> <th><u>OP-KIT</u></th> <th><u>LLBO</u></th> </tr> </thead> <tbody> <tr> <td><u>UN GHS No Cat.</u></td> <td><u>IVIS ≤ 3</u></td> <td><u>LIS ≤ 3</u></td> </tr> <tr> <td><u>NPCM</u></td> <td><u>3 < IVIS ≤ 55</u></td> <td><u>LIS > 30 and lux/7 ≤ 145 and OD₄₉₀ ≤ 2.5</u></td> </tr> <tr> <td><u>UN GHS Cat. 1</u></td> <td><u>IVIS > 55</u></td> <td><u>Lux/7 > 145 and/or OD₄₉₀ > 2.5</u></td> </tr> </tbody> </table> <p><u>NPCM : No stand-alone prediction can be made</u></p> <p>An IVIS score higher than (>) 55 leads to a UN GHS Category 1 prediction; an IVIS score smaller than or equal to (≤) 3 leads to UN GHS No Category prediction. In contrast 3 < IVIS ≤ 55, no prediction can be made on the UN GHS classification.</p>	<u>Prediction</u>	<u>OP-KIT</u>	<u>LLBO</u>	<u>UN GHS No Cat.</u>	<u>IVIS ≤ 3</u>	<u>LIS ≤ 3</u>	<u>NPCM</u>	<u>3 < IVIS ≤ 55</u>	<u>LIS > 30 and lux/7 ≤ 145 and OD₄₉₀ ≤ 2.5</u>	<u>UN GHS Cat. 1</u>	<u>IVIS > 55</u>	<u>Lux/7 > 145 and/or OD₄₉₀ > 2.5</u>
<u>Prediction</u>	<u>OP-KIT</u>	<u>LLBO</u>											
<u>UN GHS No Cat.</u>	<u>IVIS ≤ 3</u>	<u>LIS ≤ 3</u>											
<u>NPCM</u>	<u>3 < IVIS ≤ 55</u>	<u>LIS > 30 and lux/7 ≤ 145 and OD₄₉₀ ≤ 2.5</u>											
<u>UN GHS Cat. 1</u>	<u>IVIS > 55</u>	<u>Lux/7 > 145 and/or OD₄₉₀ > 2.5</u>											

Scientific basis incl. MoA	The BCOP test method addresses corneal effects, which are one of the major drivers of classification in vivo when considering the UN GHS classification (Adriaens et al., 2014; Barroso et al., 2017). Furthermore, it addresses the following Modes of Action for eye irritation: (i) cell membrane lysis (breakdown of membrane integrity as might occur from exposure to membrane active materials, e.g., surfactants), (ii) saponification (breakdown of lipids by alkaline action), and (iii) coagulation (precipitation/denaturation of macromolecules, particularly protein, characteristic of acid, alkali, or organic solvent exposure). If histopathological information is available, it may also address (iv) actions on macromolecules (chemicals that react with cellular constituents/organelles that may or may not lead to overt lysis or coagulation, e.g., alkylation, oxidative attack on macromolecules such as essential proteins or nucleic acids) (OECD, 2013 e2013b).
Protocol available	OECD Guidance Document 160 (OECD, 20112018e2018). BCOP LLBO
General reproducibility	Evaluation of the BCOP OP-KIT reliability showed a median coefficient of variation (CV) for IVIS for replicate corneas (n=3) within individual experiments ranging from 11.8% to 14.2% in one study, and median CV values for IVIS for replicate corneas (n=4) within individual experiments of 35%, respectively, in a second study (ICCVAM, 2006). The between experiment mean CV values of IVIS for 16 chemicals tested two or more times in three laboratories ranged from 12.6% to 14.8%, while the median CV values ranged from 6.7% to 12.4% (ICCVAM, 2006).
Strengths and weaknesses	<p><u>Strengths</u></p> <ul style="list-style-type: none"> - Officially validated test method. - Quantitative and objective measurements of opacity and permeability. - Controlled exposure conditions, including test chemical concentration and exposure duration. - Histological evaluation of the exposed eyes may provide additional information about e.g., the depth and type of injury (Furukawa et al., 2015; Maurer et al., 2002; OECD, 20112018) <p><u>Weaknesses</u></p> <ul style="list-style-type: none"> - The BCOP test method is not recommended for the identification of test chemicals that should be classified as irritant to eyes (UN GHS Cat. 2 or Cat. 2A) or test chemicals that should be classified as mildly irritant to eyes (UN GHS Cat. 2B) due to the considerable number of UN GHS Cat. 1 chemicals underclassified as UN GHS Cat. 2, 2A or 2B and UN GHS No Cat. chemicals overclassified as UN GHS Cat. 2, 2A or 2B. For this purpose, further testing with another suitable method may be required.

	<ul style="list-style-type: none"> - The reversibility of tissue lesions cannot be evaluated per se in the BCOP test method, although use of histological evaluations could aid predictions as to e.g., whether damage is irreversible (Furukawa et al., 2015; Maurer et al., 2002). - The BCOP OP-KIT was found to be under-predictive for identification of UN GHS Cat. 1 agrochemical formulations (Kolle et al., 2015). - Gases and aerosols have not been assessed yet in a validation study.
Identification of UN GHS Category 1	
Applicability domain and limitations	<p><u>Applicability</u></p> <ul style="list-style-type: none"> - The BCOP test method can be used for the testing of substances and mixtures (OECD, 2013a2023b, 2013c2013b, US EPA, 2015). <p><u>Limitations</u></p> <ul style="list-style-type: none"> - Positive results obtained with alcohols and ketones should be interpreted cautiously due to potential overprediction. However, since not all alcohols and ketones are overpredicted by the BCOP test method and some are correctly predicted as UN GHS Cat. 1, these two organic functional groups are not considered to be out of the applicability domain of the test method. - Solids and chemicals inducing persistent, non severe effects may risk underprediction (OECD, 2013a; Barroso et al., 2017). However, none of the false negatives identified in the ICCVAM validation database (ICCVAM, 2010, OECD, 2013a) or in the LLBO evaluation study (Adriaens et al., 2020), in the context of identifying chemicals inducing serious eye damage (UN GHS Category 1), were predicted as UN GHS No Category, resulted in IVIS ≤ 3 (criterion triggering UN GHS No Cat. prediction). Moreover, BCOP false negatives in this context are not critical since all test chemicals for which no stand-alone prediction can be made according to the UN GHS that produce an 3 < IVIS ≤ 55 would be subsequently tested and evaluated following the sequential testing strategy as described in chapter 3 and Figure 2.1. Finally, given the fact that some solid chemicals are correctly predicted by the BCOP test method as UN GHS Cat. 1, this physical state is also not considered to be out of the applicability domain of the test method. - Increased corneal permeability in the absence of corneal opacity, or in the presence of low grade corneal opacity, e.g. as observed following exposure of the bovine corneas to some types of substances (such as some surfactants and detergent products), should be carefully considered, possibly along with histopathological data, as this might indicate potential for eye hazard effects (OECD, 201120182018; ICCVAM, 2006)
Predictive capacity	When used for identification of UN GHS Cat. 1 test chemicals, the BCOP OP-KIT test method showed an overall accuracy of 79% (150/191), a false positive rate of 25% (32/126), and a false negative

	<p>rate of 14% (9/65) when compared to results obtained with the in vivo rabbit eye test method (OECD TG 405) classified according to the UN GHS classification system (OECD, 2013a, 2023b).</p> <p><u>When used for identification of UN GHS Cat. 1 test chemicals, the BCOP LLBO test method showed an overall accuracy of 78% (n=145), a false positive rate of 21% (n=89), and a false negative rate of 24% (n=56) when compared to results obtained with the in vivo rabbit eye test method (OECD TG 405) classified according to the UN GHS classification system (OECD, 2023b).</u></p>
Reliability	<p>When distinguishing UN GHS Cat. 1 chemicals from the other UN GHS eye hazard categories, 72% (91/127) of the test chemicals were found to have 100% of agreement of classification between laboratories (ICCVAM, 2010b).</p>
Identification UN GHS No Category	
Applicability domain and limitations	<p><u>Applicability</u></p> <ul style="list-style-type: none"> - The BCOP test method can be used for the testing of substances and mixtures (OECD, 2013a, 2013c, 2013b, 2023b, Kolle et al., 2016). <p><u>Limitations</u></p> <ul style="list-style-type: none"> - Since the BCOP test method can only identify correctly 31% (OP-KIT) and 55% (LLBO) of the test chemicals that do not require classification for eye irritation or serious eye damage, this test method should not be the first choice to initiate a Bottom-Up approach. Other validated and accepted in vitro methods with similar high sensitivity but higher specificity should be used instead as first tier of a Bottom-Up approach (e.g., RhCE test methods falling within OECD TG 492). Nevertheless, although the false positive rate obtained with BCOP is considerably high (69% for OP-KIT and 45% for LLBO), it is not considered critical since all test chemicals for which no stand-alone prediction can be made according to the UN GHS that produce an $3 < IVIS \leq 55$ would be subsequently tested and evaluated following the sequential testing strategy as described in chapter 3 and Figure 2.1
Predictive capacity	<p>When used for the identification of UN GHS No Cat. test chemicals, the BCOP OP-KIT test method showed an overall accuracy of 69% (135/196), a false positive rate of 69% (61/89), and a false negative rate of 0% (0/107), when compared to the in vivo rabbit eye test method (OECD TG 405) data classified according to the UN GHS classification system (OECD, 2013a, 2023b).</p> <p><u>When used for the identification of UN GHS No Cat. test chemicals, the BCOP LLBO test method showed an overall accuracy of 83% (n=145), a false positive rate of 45% (n=41), and a false negative rate of 6% (n=104), when compared to the in vivo rabbit eye test method (OECD TG 405) data classified according to the UN GHS classification system (OECD, 2023b).</u></p>

Reliability	When distinguishing UN GHS No Cat. chemicals from chemicals classified for eye hazard (UN GHS Cat. 1 and 2), 80% (103/128) of the test chemicals were found to have 100% agreement of classification between laboratories (ICCVAM, 2010b).
-------------	--

4.3.2. Isolated Chicken Eye (ICE) test method (OECD TG 438)

39. The OECD TG 438 on the ICE test method was originally adopted in 2009 and updated in 2013 and in 2018 (OECD, 2013b, 2018a, 2023c; OECD, 2013d, 2018b). The ICE test method underwent two retrospective validation studies by the US ICCVAM in conjunction with EURL ECVAM and JaCVAM, in 2006 and 2010 (ICCVAM, 2006; ICCVAM, 2010b). In the two evaluations, the ICE was endorsed as a scientifically valid test method for use as a screening test to identify UN GHS Cat. 1 substances and mixtures (ICCVAM, 2006, 2010b; ESAC, 2007). A further retrospective evaluation of the in vitro and in vivo dataset used in the validation study concluded that the ICE test method can also be used to identify UN GHS No Category substances and mixtures (OECD, 2013d, 2018b, 2023c). From these evaluations it was concluded that the ICE test method can correctly identify test chemicals (both substances and mixtures) inducing serious eye damage (UN GHS Cat. 1) as well as those not requiring classification for eye hazard (UN GHS No Category). Furthermore, histopathology has been shown to be a useful additional endpoint to identify UN GHS Category 1 non-extreme pH (2 < pH < 11.5) detergents and surfactants (Cazelle et al., 2014; OECD GD 188, 2018b, 2019; OECD GD 160, 2018a, e).

Module 3 – In vitro data: Isolated Chicken Eye (ICE) test method (OECD TG 438)	
General description	
Regulatory use (UN GHS classification)	Identification of i) test chemicals inducing serious eye damage (UN GHS Cat. 1), and ii) test chemicals not requiring classification for eye hazard (UN GHS No Cat.).
Validation & regulatory acceptance status	Validated and adopted as OECD TG 438.
Potential role in the IATA	<p>While the ICE test method is not considered valid as a full-replacement for the in vivo rabbit eye test, it can be used to initiate either the Top-Down or the Bottom-Up approach for regulatory classification and labelling (Figure 2.1. to identify, without further testing:</p> <ul style="list-style-type: none"> - test chemicals inducing serious eye damage (UN GHS Cat. 1); and - test chemicals that do not require classification for eye hazard (UN GHS No Cat.). <p>A test chemical that is neither predicted as UN GHS Cat. 1 nor as UN GHS No Cat. with the ICE test method would require further testing with a UN GHS Cat. 2 method and/or additional WoE evaluation with other existing information and if still needed additional testing (in vitro</p>

	and/or in vivo) as a last resort to establish a definitive classification (Figure 2.1).
Description	<p>The ICE test method is an organotypic ex vivo assay based on the short-term maintenance of chicken eyes in vitro. In this test method, damage by the test chemical is assessed Toxic effects to the cornea are measured by (i) a quantitative measurement of increased corneal thickness (swelling), (ii) a qualitative assessment of corneal opacity, (iii) a qualitative assessment of damage to epithelium based on application of fluorescein to the eye (fluorescein retention), and (iv) a qualitative evaluation of macroscopic morphological damage to the surface. Furthermore, histopathology can be used to increase the sensitivity of the method for identifying UN GHS Category 1 non-extreme pH ($2 < \text{pH} < 11.5$) detergents and surfactants. In particular, If histopathological information is available, it may also address depth of injury and predict reversibility of effects (OECD, 20112018e2019; Maurer et al., 2002; Cazelle et al., 2014), depth of injury (Maurer et al., 2002) as well as possible actions on macromolecules (chemical effects on cellular constituents/organelles that may or may not lead to overt lysis or coagulation due to e.g., alkylation, oxidative attack on macromolecules such as essential proteins or nucleic acids) (Scott et al., 2010)."</p> <p>The corneal swelling, opacity and damage assessments following exposure to a test chemical are assessed individually and assigned a qualitative categorization, that are then combined together to derive an in vitro eye hazard classification, either as UN GHS Cat. 1 or as UN GHS No Cat. However, no decision on classification can be made for test chemicals not predicted to be UN GHS Cat. 1 or UN GHS No Cat. with the ICE test method.</p>
Scientific basis incl. MoA	<p>The ICE test method addresses corneal effects, which are one of the major drivers of classification in vivo when considering the UN GHS classification (Adriaens et al., 2014; Barroso et al., 2017). Furthermore, it addresses the following Modes of Action for eye irritation: (i) cell membrane lysis (breakdown of membrane integrity as might occur from exposure to membrane active materials, e.g., surfactants), (ii) saponification (breakdown of lipids by alkaline action), and (iii) coagulation (precipitation/denaturation of macromolecules, particularly protein, characteristic of acid, alkali, or organic solvent exposure). If histopathological information is available, it may also address (iv) (ir)reversibility of effects and (v) actions on macromolecules (chemicals that react with cellular constituents/organelles that may or may not lead to overt lysis or coagulation, e.g., alkylation, oxidative attack on macromolecules such as essential proteins or nucleic acids) (Scott et al., 2010; OECD, 2018b2018a; OECD, 2018e2019).</p>
Protocol available	OECD Guidance Document 160 (OECD, 20172018e2018a).
General reproducibility	Evaluation of the ICE reliability (without histopathology) showed coefficient of variation (CV) values for the corneal thickness

	<p>measurement, when results were compared within experiments, varying from 1.8% to 6.3% (OECD, 2013d2018b2019). The other endpoints evaluated produced larger ranges of CV values due to the relatively small values that were produced by test chemicals not requiring classification. Regarding the between-laboratory reproducibility of the ICE test method (without histopathology), the EC/HO international validation study on alternatives to the in vivo rabbit eye test showed inter-laboratory correlations of 82.9, 84.9 and 84.4% (OECD, 2013d2018b2019).</p> <p>Regarding histopathology, appropriate reproducibility was found between pathologists and peer-reviewers from three independent laboratories of (10/12 or 83%) and over time (17/18 for non-extreme pH detergents and 6/6 for surfactants) for the ICE histopathological derived predictions (OECD, SD 188, 2019). However, to ensure such reproducibility, there is a need for (i) an internal peer-review system to be in place; (ii) assessment of the original slides in order to enable the evaluation of three dimensional effects; and (iii) appropriate training & proficiency appraisal.</p>
Strengths and weaknesses	<p><u>Strengths</u></p> <ul style="list-style-type: none"> - Officially validated test method. - Measurements are performed both quantitatively and qualitatively with the help of a slit-lamp. - Controlled exposure conditions, including test chemical concentration and exposure duration. - Histological evaluation of the exposed eyes may provide additional information about e.g., the depth and type of injury and reversibility of effects allows identification of UN GHS Cat. 1 non-extreme pH (2 < pH < 11.5) detergents and surfactants (Maurer et al., 2002; Cazelle et al., 2014; OECD, 2018a,b; OECD, 20112018c2019; Cazelle et al., 2014) <p><u>Weaknesses</u></p> <ul style="list-style-type: none"> - The ICE test method is not recommended for the identification of test chemicals that should be classified as irritant to eyes (UN GHS Cat. 2 or Cat. 2A) or test chemicals that should be classified as mildly irritant to eyes (UN GHS Cat. 2B) due to the considerable number of UN GHS Cat. 1 chemicals underclassified as UN GHS Cat. 2, 2A or 2B and UN GHS No Cat. chemicals overclassified as UN GHS Cat. 2, 2A or 2B. For this purpose, further testing with another suitable method may be required. - The reversibility of tissue lesions cannot be evaluated per se in the ICE test method. However, histological evaluation could aid predictions as to e.g., whether damage is irreversible (OECD, 2018b; 2018c2019; Cazelle et al., 2014; Maurer et al., 2002). - Gases and aerosols have not been assessed yet in a validation study.

Identification of UN GHS Category 1	
Applicability domain and limitations	<p><u>Applicability</u></p> <ul style="list-style-type: none"> - The ICE test method can be used for the testing of substances and mixtures (OECD, 2013b2018a, 2013d2018b2019, 2023c; Cazelle et al., 2014, 2015). - It is applicable to solids, liquids, emulsions and gels. Liquids may be aqueous or non-aqueous and solids may be soluble or insoluble in water. <p><u>Limitations</u></p> <ul style="list-style-type: none"> - Positive results obtained with alcohols should be interpreted cautiously due to potential overprediction. However, since not all alcohols are overpredicted by the ICE test method and some are correctly predicted as UN GHS Cat. 1, this organic functional groups is not considered to be out of the applicability domain of the test method. - Solids, surfactants and chemicals inducing persistent, non severe effects may risk underprediction (OECD, 2019, 2013b2018a2023c; Barroso et al., 2017). However, false negative rates in this context (UN GHS Cat. 1 identified as not being UN GHS Cat. 1) are not critical since all test chemicals that come out negative would be subsequently tested and evaluated following the sequential testing strategy as described in chapter 3 and Figure 2.1. Furthermore use of histopathology may help to decrease the under-prediction of non-extreme pH detergents ($2 < \text{pH} < 11.5$) and surfactants (OECD, 2018^{ab}; OECD, 2018e2019; Cazelle et al., 2014).
Predictive capacity	<p>When used for identification of UN GHS Cat. 1 test chemicals, the ICE test method showed an overall accuracy of 8683% (142/172420/440), a false positive rate of 76% (9/1277/113) and a false negative rate of 4748% (21/4513/27) when compared to in vivo rabbit eye test method (OECD TG 405) classified according to the UN GHS classification system (OECD, 2013b2018a2023c).</p> <p>When histopathology is considered as an additional endpoint to identify UN GHS Category 1 non-extreme pH ($2 < \text{pH} < 11.5$) detergents and surfactants, the false negative rate of the ICE test method is decreased and its accuracy is increased (from 64% to 27% false negatives (n=22) and from 53% to 77% accuracy (n=30)), whilst an acceptable false positive rate is maintained (from 0% to 12.5% false positives (n=8)) (OECD, 2018b; OECD, 2018e2023c).</p>
Reliability	When distinguishing UN GHS Cat. 1 from the other UN GHS eye hazard categories, a between-laboratories reproducibility of 75% (44/59) was observed (ICCVAM, 2006).
Identification UN GHS No Category	

Applicability domain and limitations	<p><u>Applicability</u></p> <p>The ICE test method can be used for the testing of substances and mixtures (OECD, 2013b, 2013d2019, 2023c)</p> <p><u>Limitations</u></p> <p>Anti fouling organic solvent-containing paints may be underpredicted (OECD 2013d2019, 2023c)</p> <p>In the case of solid materials leading to GHS No Cat. Outcome , a second run of three eyes is recommended to confirm or discard the negative outcome.</p>
Predictive capacity	<p>When used for the identification of UN GHS No Cat. test chemicals, the ICE test method showed an overall accuracy of 8288% (161/184125/152), a false positive rate of 2433% (20/8326/79), and a false negative rate of 31% (3/1014/73), when compared to in vivo rabbit eye test method (OECD TG 405) classified according to the UN GHS (OECD, 2013b2018a2023c). When anti-fouling organic solvent containing paints are excluded from the database, the accuracy of the ICE test method was found to be 8388% (159/181123/149), the false positive rate 2433% (20/8326/78), and the false negative rate of 20% (2/990/74) for the UN GHS classification system (OECD, 2013b2018a2023c).</p>
Reliability	<p>When distinguishing UN GHS No Cat. from chemicals classified for eye hazard (UN GHS Cat. 1 and 2), 75% (44/59) of the tested chemicals were found to have 100% agreement of classification between laboratories (ICCVAM, 2010b).</p>

4.3.3. Short Time Exposure (STE) test method (OECD TG 491)

40. The OECD TG 491 on the STE test method was adopted in 2015 [and updated in 2018 and 2020](#) (OECD, [2015a](#)[2023e](#)). The STE test method underwent two prospective validation studies, one conducted by the Validation Committee of the Japanese Society for Alternative to Animal Experiments (JSAE) (Sakaguchi et al., 2011) and another by JaCVAM (Kojima et al., 2013). A peer review was conducted by the US ICCVAM (ICCVAM, 2013), and from these evaluations it was concluded that the STE test method can correctly identify test chemicals (both substances and mixtures) inducing serious eye damage (UN GHS Cat. 1) as well as chemicals (excluding highly volatile substances and all solid chemicals other than surfactants) not requiring classification for eye hazard (UN GHS No Category). The data set was enlarged in the CON4EI project with 80 chemicals, 49/80 chemicals were unique to this project (Adriaens et al. 2017a, 2017c). [Further work demonstrated that highly volatile substances can be correctly tested when using mineral oil instead of saline as a solvent \(Abo et al., 2018\).](#)

Module 3 – In vitro data: Short Time Exposure (STE) test method (OECD TG 491)

General description

Regulatory use (UN GHS classification)	Identification of i) test chemicals inducing serious eye damage (UN GHS Cat. 1), and ii) test chemicals (excluding highly volatile substances and all solid chemicals other than surfactants) not requiring classification for eye hazard (UN GHS No Cat.).
Validation & regulatory acceptance status	Validated and adopted as OECD TG 491-.
Potential role in the IATA	<p>While the STE test method is not considered valid as a full-replacement for the in vivo rabbit eye test, it can be used to initiate either the Top-Down or the Bottom-Up approach for regulatory classification and labelling (Figure 2.1) to identify, without further testing:</p> <ul style="list-style-type: none"> - test chemicals inducing serious eye damage (UN GHS Cat. 1); and - limited types of test chemicals (excluding highly volatile substances and solid substances and mixtures other than surfactants) that do not require classification for eye hazard (UN GHS No Cat.). <p>A test chemical that is neither predicted as UN GHS Cat. 1 nor as UN GHS No Cat. with the STE test method would require <u>further testing with a UN GHS Cat. 2 method and/or</u> additional WoE evaluation with other existing information and if still needed additional testing (in vitro and/or in vivo) as a last resort to establish a definitive classification (Figure 2.1).</p>
Description	<p>The STE test method is a cytotoxicity-based in vitro assay that is performed on a confluent monolayer of Statens Seruminstitut Rabbit Cornea (SIRC) cells, cultured on a 96-well polycarbonate microplate. Each test chemical is tested at both 5% and 0.05% concentrations. After five-minute exposure to the test chemical, cell viability is assessed by the quantitative measurement, after extraction from cells, of the blue formazan salt produced by the living cells by enzymatic conversion of the vital dye MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide), also known as Thiazolyl Blue Tetrazolium Bromide (Mosmann, 1983).</p> <p>The relative cell viability of the treated SIRC cells (compared to the solvent control) is used to estimate the potential eye hazard of the test chemical. A test chemical is classified as UN GHS Cat. 1 when both the 5% and 0.05% concentrations result in a relative cell viability smaller than or equal to (\leq) 70%. Conversely, a test chemical is predicted as UN GHS No Cat. when both 5% and 0.05% concentrations result in a relative cell viability higher than ($>$) 70%.</p>
Scientific basis incl. MoA	It has been reported that 80% of a solution dropped into the eye of a rabbit is excreted through the conjunctival sac within three to four minutes, while greater than 80% of a solution dropped into the human eye is excreted within one to two minutes. The STE test

	method attempts to approximate these exposure times using the five-minute exposure to the test chemical. Decrease in cell viability is then used to predict potential adverse effects leading to ocular damage. The cytotoxic effects of test chemicals on corneal epithelial cells is an important mode of action leading to corneal epithelium damage and eye irritation.
Protocol available	The Test Guideline is based on a protocol developed by Kao Corporation (Takahashi et al., 2008). <u>STE Protocol available at:</u> https://ntp.niehs.nih.gov/whatwestudy/niceatm/test-method-evaluations/ocular/ste
General reproducibility	Evaluation of the STE reliability showed CV values for within-laboratory variability for test chemicals classified as UN GHS No Cat. spanning from 0.3% to 23.5% in four studies evaluated. Test chemicals classified in vitro tended to have greater CV values, as expected, because the cell viability for these test chemicals was often quite low. The mean viability for the positive control, 0.01% sodium lauryl sulfate, was 41.7% (N = 71) with a CV of 24.7%. A between-laboratory reproducibility of 83-100% was observed (ICCVAM, 2013).
Strengths and weaknesses	<u>Strengths</u> <ul style="list-style-type: none"> - Officially validated test method. - Quantitative measurements of cell viability. - Controlled exposure conditions, including test chemical concentration and exposure duration. <u>Weaknesses</u> <ul style="list-style-type: none"> - The STE test method is not recommended for the identification of test chemicals that should be classified as irritant to eyes (UN GHS Cat. 2 or Cat. 2A) or test chemicals that should be classified as mildly irritant to eyes (UN GHS Cat. 2B) due to the considerable number of UN GHS Cat. 1 chemicals underclassified as UN GHS Cat. 2, 2A or 2B and UN GHS No Cat. chemicals overclassified as UN GHS Cat. 2, 2A or 2B. For this purpose, further testing with another suitable method may be required. - The reversibility of tissue lesions cannot be evaluated per se in the STE test method. - Gases and aerosols have not been assessed yet in a validation study. <p style="text-align: center;">Identification of UN GHS Category 1</p>
Applicability domain and limitations	<u>Applicability</u>

	<p>- Test chemicals (substances and mixtures) that are dissolved or uniformly suspended for at least 5 minutes in physiological saline, 5% dimethyl sulfoxide (DMSO) in saline, or mineral oil.</p> <p><u>Limitations</u></p> <p>- The high false negative rate observed (51%), is not critical in the present context, since all test chemicals that induce a cell viability of $\leq 70\%$ at a 5% concentration and $> 70\%$ at 0.05% concentration would be subsequently would be subsequently tested and evaluated following the sequential testing strategy as described in chapter 3 and Figure 2.1.</p>
Predictive capacity	When used for identification of UN GHS Cat. 1 test chemicals, the STE test method showed an overall accuracy of 83% (104/125), a false positive rate of 1% (1/86), and a false negative rate of 51% (20/39) as compared to the in vivo rabbit eye test method (OECD TG 405) classified according to the UN GHS (OECD, 2015a2023e).
Identification UN GHS No Category	
Applicability domain and limitations	<p><u>Applicability</u></p> <p>- Test chemicals (substances and mixtures) that are dissolved or uniformly suspended for at least 5 minutes in physiological saline, 5% dimethyl sulfoxide (DMSO) in saline, or mineral oil.</p> <p><u>Limitations</u></p> <p>- Mixtures containing Hhighly volatile substances with a vapour pressure over 6 kPa (at 25°C25°C) that do not dissolve in mineral oil, or that do not form stable suspensions for at least 5 minutes, are excluded from the applicability domain of the STE test method for the identification of UN GHS No Cat. due to the high false negative rate. Results obtained with mixtures containing substances with vapour pressure higher than 6kPa should be interpreted cautiously due to potential underprediction, and should be justified on a case-by-case basis.</p> <p>- Solid chemicals (substances and mixtures) other than surfactants and mixtures composed only of surfactants are also excluded from the applicability domain of the STE test method for the identification of UN GHS No Cat. due to high false negative rates observed.</p> <p>- Chemicals should be assessed for direct reduction of MTT as advised in the STE test protocol (NICEATM, 2012)</p>
Predictive capacity	When used for the identification of UN GHS No Cat. test chemicals, the STE test method showed an overall accuracy of 85% (110/130), a false negative rate of 12% (9/73), and a false positive rate of 19% (11/57) as compared to the in vivo rabbit eye test method (OECD TG 405) classified according to the UN GHS (OECD, 2015a2023e). If highly volatile substances and solid chemicals (substances and mixtures) other than surfactants are excluded from the dataset, the

	<p>overall accuracy improves to 90% (92/102), the false negative rate to 2% (1/54), and the false positive to 19% (9/48). Further work demonstrated that highly volatile substances can be correctly tested when using mineral oil instead of saline as a solvent (Abo et al., 2018). The accuracy of the STE test for highly volatile substances (i.e. vapour pressure > 6kPa) was then 95% (19/20), the false negative rate was 0% (0/7), and the false positive rate was 8% (1/13). In addition in-house data on 40 mixtures, showed an accuracy of 88% (35/40), a false positive rate of 50% (5/10), and a false negative rate of 0% (0/30) for predicting UN GHS No Cat. when compared to the in vivo rabbit eye test (Saito et al., 2015).</p>
--	---

4.3.4. Reconstructed human Cornea-like Epithelium Eye Irritation Test (RhCE EIT) (OECD TG 492)

41. The OECD TG 492 on the RhCE Test Methods was adopted in 2015 and revised in 2017, 2019, and 2023 (OECD, 2017, 2019, 2023). The in vitro test methods currently covered by this Test Guideline are the EpiOcular™ Eye Irritation Test (EIT), ~~which makes use of the commercially available EpiOcular™ OCL-200 RhCE tissue construct, and the SkinEthic™ HCE Eye Irritation Test, which makes use of the commercially available SkinEthic™ Human Corneal Epithelium (HCE/S) tissue construct~~ [the LabCyte CORNEA-MODEL24 EIT, and the MCTT HCE™ EIT. Each of these methods makes use of commercially available RhCE tissue constructs as test system \(EpiOcular™ OCL-200, SkinEthic™ Human Corneal Epithelium \(HCE/S\), LabCyte CORNEA-MODEL24, and MCTT HCE™\).](#) The ~~EpiOcular™ EIT and SkinEthic™ HCE EIT two EIT~~ test methods underwent a prospective validation study conducted by EURL ECVAM and Cosmetics Europe (Barroso et al., 2014) and by industry (Alépée et al., 2016a, 2016b) respectively, and the outcome was peer-reviewed by the EURL ECVAM Scientific Advisory Committee (ESAC, 2014, 2016a). ~~They are referred to in the as the Validated Reference Methods – VRM1 (EpiOcular™ EIT) and VRM 2 (SkinEthic™ HCE EIT). The LabCyte CORNEA-MODEL24 EIT method was validated in a performance standard based study and peer-reviewed by JaCVAM (OECD, 2018b). The MCTT HCE™ underwent a me-too validation study (Lim et al., 2019).~~ From these evaluations it was concluded that the ~~both four~~ EIT test methods can correctly identify test chemicals (substances and mixtures) not requiring classification for eye hazard (UN GHS No Category). In contrast to the in vitro methods described earlier (BCOP, ICE and STE), the RhCE EIT ~~test methods is~~ are not applicable for the identification of test chemicals inducing serious eye damage (UN GHS Cat. 1). The data set was enlarged in the CON4EI project with 80 chemicals, 27 up to 32 chemicals were unique to this project (Adriaens et al. 2017a, 2017b; Kandarova et al., 2017a; Van Rompay et al., 2017).

Module 3 – In vitro data: Reconstructed human Cornea-like Epithelium – Eye Irritation Test (OECD TG 492)	
General description	
Regulatory use (UN GHS classification)	Identification of test chemicals not requiring classification for eye hazard (UN GHS No Cat.).

Validation & regulatory acceptance status	Validated and adopted as OECD TG 492.
Potential role in the IATA	<p>While the EpiOcular™ EIT, and SkinEthic™ HCE EIT, <u>LabCyte CORNEA-MODEL24 EIT, and MCTT HCE™ EIT</u> is are not considered valid as a full-replacement for the in vivo rabbit eye test, it they can be used within the Top-Down and Bottom-Up approaches and in particular to initiate the Bottom-Up approach for regulatory classification and labelling (Figure 2.1) to identify, without further testing:</p> <ul style="list-style-type: none"> - test chemicals that do not require classification for eye hazard (UN GHS No Cat). <p>The EpiOcular™ EIT, and SkinEthic™ HCE EIT, <u>LabCyte CORNEA-MODEL24 EIT, and MCTT HCE™ EIT</u> is are not intended to differentiate between UN GHS Cat. 1 (serious eye damage) and UN GHS Cat. 2 (eye irritation). This differentiation will need to be addressed by another tier of a test strategy (Figure 2.1). A test chemical that is not predicted as not requiring classification for eye hazard (UN GHS No Cat.) with EpiOcular™ EIT, or SkinEthic™ HCE EIT, <u>LabCyte CORNEA-MODEL24 EIT or MCTT HCE™ EIT</u> will thus require additional in vitro testing and/or additional WoE evaluation with other existing information before progressing with further testing within the IATA in order to establish a definitive classification (Figure 2.1).</p>
Description	<p>Three-dimensional RhCE tissues are reconstructed from primary human cells, which have been cultured for several days to form a stratified, highly differentiated squamous epithelium morphologically similar to that found in the human cornea. The EpiOcular™, <u>LabCyte CORNEA-MODEL24, and MCTT HCE™</u> RhCE tissue construct consists of at least 3 viable layers of cells and a non-keratinized surface, showing a cornea-like structure analogous to that found in vivo. The SkinEthic™ HCE tissue construct consists of at least 4 viable cell layers, including columnar cells and wing cells, with the presence of intermediate filaments, mature hemi-desmosomes and desmosomes, and specific human corneal cytokeratins. The test chemical is applied topically to a minimum of two RhCE tissue constructs. Following the exposure and post-treatment incubation periods, tissue viability is assessed by the enzymatic conversion of tetrazolium dye [MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; Thiazolyl blue tetrazolium bromide for EpiOcular™ EIT and SkinEthic™ HCE EIT; WST-8 [2-(2-methoxy-4-nitrophenyl)-3-(4-nitrophenyl)-5-(2,4-disulfophenyl)-2H-tetrazolium, monosodium salt for LabCyte CORNEA-MODEL24 EIT or WST-1 [4-[3-(4-Iodophenyl)-2-(4-nitrophenyl)-2H-5-tetrazol-1,3-benzene disulfonate for MCTT HCE™ EIT] in viable cells of the vital dye MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; Thiazolyl blue tetrazolium bromide) into a <u>blue coloured formazan dye (blue MTT formazan, or yellow WST-8 and WST-1 formazan) MTT formazan salt</u> which is extracted from the tissues and quantitatively measured (Mosmann, 1983). Test chemicals</p>

	not requiring classification and labelling according to UN GHS are identified as those having a tissue viability higher than (>) 60% for EpiOcular™ EIT and SkinEthic™ HCE EIT liquids' protocol, or > 50% for SkinEthic™ HCE EIT solids' protocol, <u>or > 40% for LabCyte CORNEA-MODEL24 EIT or > 35% for liquids and > 60% for solids in MCTT HCE™ EIT.</u>
Scientific basis incl. MoA	<p>Reconstructed human cornea-like epithelium (RhCE) closely mimics the histological, morphological, biochemical and physiological properties of the human corneal epithelium.</p> <p>The use of viability of the RhCE tissues after topical exposure to a test chemical to discriminate UN GHS No Cat. chemicals from those requiring classification and labelling (UN GHS Cat. 1 and 2) is based on the assumption that all chemicals inducing serious eye damage or eye irritation will induce cytotoxicity in the corneal epithelium and/or conjunctiva. Indeed, it has been shown that cytotoxicity plays an important mechanistic role in determining the overall serious eye damage and eye irritation response of a chemical regardless of the physicochemical processes underlying tissue damage (Jester et al., 1998; Maurer et al., 2002).</p>
Protocol available	<p>DB-ALM protocol no. 164 on the EpiOcular™ EIT test method (DB-ALM, 2015).</p> <p>DB-ALM protocol no. 190 on the SkinEthic™ HCE Eye Irritation Test Liquid (EITL) (DB-ALM, 2017).</p> <p>DB-ALM protocol no. 191 on the SkinEthic™ HCE Eye Irritation Test Solid (EITS) (DB-ALM, 2017).</p> <p><u>LabCyte CORNEA-MODEL24 EIT SOP, Version 2.5.6. (February, 2017). LabCyte CORNEA-MODEL24 eye irritation test operation protocol. Available at: http://www.iacvam.jp/files/doc/06_11/06_11_E1.pdf.</u></p> <p><u>MCTT HCE™ EIT SOP, Version 1.7. (August, 2018). MCTT HCE™ eye irritation test operation protocol. Available at: http://www.keraskin.co.kr/eng/product/mucosalmodel.asp</u></p>
Strengths and weaknesses	<p><u>Strengths</u></p> <ul style="list-style-type: none"> - Officially validated test methods. - Human-based 3D tissue models. - Quantitative measurements of cell viability. - Controlled exposure conditions, including test chemical concentration and exposure duration. <p><u>Weaknesses</u></p> <ul style="list-style-type: none"> - The EIT RhCE test methods does not allow discrimination between eye irritation/reversible effects on the eye (UN GHS Cat. 2) and serious eye damage/irreversible effects on the eye (UN GHS Cat. 1), nor

	<p>between eye irritants (UN GHS Cat. 2A) and mild eye irritants (UN GHS Cat. 2B). For these purposes, further testing with other suitable test methods is required.</p> <ul style="list-style-type: none"> - Gases and aerosols have not been assessed yet in a validation study. - The reversibility of tissue lesions cannot be evaluated per se in the EIT RhCE test methods.
Identification UN GHS No Category	
Applicability domain and limitations	<p><u>Applicability</u></p> <ul style="list-style-type: none"> - The RhCE test methods can be used for the testing of substances and mixtures (OECD, 2015b72023f; Kolle et al., 2015, 2016; Kandárová et al., 2017a; Alépée et al., 2016a, 2016b; Van Rompay et al., 2017). - It is applicable to solids, liquids, semi-solids and waxes. Liquids may be aqueous or non-aqueous and solids may be soluble or insoluble in water (OECD, 20172023f). <p><u>Limitations</u></p> <ul style="list-style-type: none"> - test chemicals presenting non-specific interactions with MTT (i.e., absorbing light in the same range as MTT formazan (naturally or after treatment) or able to directly reduce the vital dye MTT (to MTT formazan)) that are $\geq 60\%$ should be taken with caution when OD is used to measure the extracted MTT formazan. However, use of HPLC/UPLC-spectrophotometry as an alternative procedure to measure MTT formazan allows circumventing this, and is especially useful for those test chemicals strongly absorbing in the same wavelength as MTT formazan which are not compatible with the standard optical density (OD) measurement (Alépée et al., 2015).
Predictive capacity	<p>When used for the identification of UN GHS No Cat. test chemicals, the EpiOcular™ EIT test method showed an overall accuracy of 80% (based on 112 chemicals), a false negative rate of 4% (based on 57 chemicals), and a false positive rate of 37% (based on 55 chemicals) when compared to in vivo rabbit eye test data (OECD TG 405) classified according to the UN GHS (OECD, 2015b72023a). When used for the identification of UN GHS No Cat. test chemicals, the SkinEthic™ HCE EIT test method showed an overall accuracy of 84% (based on 200 chemicals), a false negative rate of 5% (based on 97 chemicals), and a false positive rate of 28% (based on 103 chemicals) (Alépée et al., 2016a, 2016b). In addition, a study on agrochemical formulations using the EpiOcular™ EIT RhCE test method according to OECD TG 492, showed an overall accuracy of 82% (based on 97 formulations), a false negative rate of 9% (based on 54 formulations) and a false positive rate of 28% (based on 43 formulations) for predicting UN GHS No Cat. when compared to the in vivo rabbit eye test (Kolle et al., 2015).</p>
Reliability	<p>The concordance of predictions obtained with the EpiOcular™ EIT RhCE test method was found to be in the order of 95% within laboratories and 93% between laboratories. The concordance of</p>

	predictions obtained with the SkinEthic™ HCE EIT test method was found to be in the order of 92% within laboratories and 95% between laboratories (based on 120 chemicals).
--	---

4.3.5. Fluorescein Leakage (FL) test method (OECD TG 460)

42. The OECD TG 460 on the FL test method was adopted in 2012 [and revised in 2017](#) (OECD, [2012b](#), [2012c](#)[2023a](#)). The FL test method has been evaluated in a retrospective validation study coordinated by EURL ECVAM in collaboration with US ICCVAM and JaCVAM (EURL ECVAM, 2008a, b), followed by peer review by the ESAC (ESAC, 2009b). From these evaluations it was concluded that the test method can correctly identify water-soluble test chemicals (both substances and mixtures) inducing serious eye damage (UN GHS Cat. 1). In contrast to the in vitro methods described earlier (BCOP, ICE, STE, RhCE), the FL assay is not applicable for the identification of test chemicals not requiring classification for eye hazard (UN GHS No Category).

Module 3 – In vitro data: Fluorescein Leakage (FL) test method (OECD TG 460)	
General description	
Regulatory use (UN GHS classification)	Identification of test chemicals inducing serious eye damage (UN GHS Cat. 1).
Validation & regulatory acceptance status	Validated and adopted as OECD TG 460.
Potential role in the IATA	<p>While the FL test method is not considered valid as a full-replacement for the in vivo rabbit eye test, it can be used within the Top-Down and Bottom-Up approaches and in particular to initiate the Top-Down approach for regulatory classification and labelling (Figure 2.1) to identify, without further testing:</p> <ul style="list-style-type: none"> - limited types of test chemicals (water soluble substances and mixtures), inducing serious eye damage (UN GHS Cat. 1). <p>A test chemical that is not predicted to be UN GHS Cat. 1 with the FL test method will require additional in vitro testing and/or additional WoE evaluation with other existing information before progressing with further testing within the IATA in order to establish a definitive classification (see Figure 2.1).</p>
Description	<p>The FL test method is a cell-function based in vitro assay that is performed on a confluent monolayer of Madin-Darby Canine Kidney (MDCK) CB997 tubular epithelial cells cultured on permeable inserts. The toxic effects of a test chemical are measured after a short exposure time (1 minute) by an increase in permeability of sodium fluorescein through the epithelial monolayer of MDCK cells. The amount of fluorescein leakage that occurs is proportional to the chemical-induced</p>

	<p>damage to the tight junctions, desmosomal junctions and cell membranes, and is used to estimate the ocular toxicity potential of a test chemical. The concentration of test chemical (mg/mL) causing 20% FL relative to the value recorded for the untreated confluent monolayer and inserts without cells (FL20), is used to predict UN GHS Cat. 1 classification (i.e., FL20 \leq 100 mg/ml: UN GHS Cat. 1).</p>
<p>Scientific basis incl. MoA</p>	<p>The potential for a test chemical to induce serious eye damage is assessed by its ability to induce damage to an impermeable confluent epithelial monolayer. The MDCK cell line model the non-proliferating state of the in vivo corneal epithelium and forms tight junctions and desmosomal junctions similar to those found on the apical side of conjunctival and corneal epithelia.</p> <p>The short exposure period allows water-based substances and mixtures to be tested neat, if they can be easily removed after the exposure period, which allows more direct comparisons of the results with the chemical effects in humans.</p> <p>The integrity of trans-epithelial permeability is a major function of an epithelium such as that found in the conjunctiva and the cornea. Trans-epithelial permeability is controlled by various tight junctions. Tight and desmosomal junctions in vivo prevent solutes and foreign materials penetrating the corneal epithelium. Loss of trans-epithelial impermeability, due to damaged tight junctions and desmosomal junctions, is one of the early events in chemical-induced ocular irritation. Increasing permeability of the corneal epithelium in vivo has been shown to correlate with the level of inflammation and surface damage observed as eye irritation develops (OECD, 2012/2023d).</p>
<p>Protocol available</p>	<p>DB-ALM Protocol no 71 on the FL test method (DB-ALM, 2013).</p>
<p>General reproducibility</p>	<p>Based on the data acquired in the validation study for 60 chemicals according to INVITTOX protocol 71, 43/60 materials (71.7%) had 100% agreement among all 4 participating laboratories. When concordance between 3 of the 4 laboratories was investigated, 59/60 materials (98.3%) had 100% agreement among 3 of the 4 laboratories. Moreover, data from INVITTOX protocol 120 were used as weight of evidence to further assess the Reproducibility of the FL test method. A good agreement of classification was obtained with 7/9 materials (77.8%) having 100% agreement among 3 laboratories, and 26/29 materials (89.7%) having 100% agreement among 2 laboratories (OECD, 2012/2023a).</p>
<p>Strengths and weaknesses</p>	<p><u>Strengths</u></p> <ul style="list-style-type: none"> - Officially validated test method. - Quantitative measurements. - Controlled exposure conditions, including test chemical concentration and exposure duration.

	<p>- The FL test method may also assess recovery. Preliminary analyses indicated that recovery data (up to 72 h following exposure to the test chemical) could potentially increase the predictive capacity of the FL test method, although further evaluation is needed and would benefit from additional data preferably acquired by further testing (OECD, 20122023d).</p> <p><u>Weaknesses</u></p> <p>- The FL test method is not recommended for the identification of test chemicals that should be classified as mild/moderate irritants (UN GHS Cat. 2 or UN GHS Cat. 2A and 2B), or of test chemicals which should not be classified for ocular irritation (UN GHS No Cat.), as demonstrated by the validation study (EURL ECVAM, 2008).</p> <p>- Gases and aerosols have not been assessed yet in a validation study.</p>
Identification of UN GHS Category 1	
Applicability domain and limitations	<p><u>Applicability</u></p> <p>- The test method is applicable to water soluble test chemicals (substances and mixtures) and/or where the toxic effect is not affected by dilution.</p> <p><u>Limitations</u></p> <p>- Strong acids and bases, cell fixatives and highly volatile test chemicals are excluded from the applicability domain as these chemicals have mechanisms that are not measured by the FL test method, e.g. extensive coagulation, saponification or specific reactive chemistries.</p> <p>- Coloured and viscous test chemicals are difficult to remove from the monolayer following the short exposure period but predictivity of the test method could be improved if a higher number of washing steps was used.</p> <p>- The final concentration to cells of solid test chemicals suspended in liquid that have the propensity to precipitate can be difficult to determine.</p>
Predictive capacity	<p>When used for identification of UN GHS Cat. 1 test chemicals, data obtained with the FL test method showed an accuracy of 77% (117/151), a false positive rate of 7% (7/103) and a false negative rate of 56% (27/48) when compared to in vivo rabbit eye test method (OECD TG 405) classified according to the UN GHS classification system (OECD, 2012b2023d).</p>

4.3.6. Defined Approaches for Serious Eye Damage and Eye Irritation (OECD TG 467)

43. The OECD TG 467 on Defined Approaches for Serious Eye Damage and Eye Irritation was adopted in 2022 (OECD, 2022b). Two rule-based DAs for non-surfactant liquids (DAL-1 and DAL-2) are currently included in this TG. The DAL-1 is based on the use of a combination of test methods described in OECD TG 437 and TG 492 as well as the physicochemical properties of the test chemical (Alépée et al., 2019a). DAL-2 in contrast, is based on the use of a combination of test methods described in the OECD TG 437 and TG 491 (Alépée et al., 2019b). The methods used in DAL-1 and DAL-2 encompass the following validated test methods: the Bovine Corneal Opacity and Permeability (BCOP) using the laser light-based opacitometer (LLBO) according to the OECD TG 437, the Reconstructed human Cornea-like Epithelium (RhCE) according to the OECD TG 492 and the Short Time Exposure in vitro (STE) according to the OECD TG 491.

<u>Module 3 – Defined Approaches for Serious Eye Damage and Eye Irritation (OECD TG 467)</u>	
<u>General description</u>	
<u>Regulatory use (UN GHS classification)</u>	<u>Eye Hazard identification.</u>
<u>Validation & regulatory acceptance status</u>	<u>Adopted as OECD TG 467.</u>
<u>Potential role in the IATA</u>	<u>Full-replacement for the in vivo rabbit eye test for regulatory classification and labelling (Figure 2.1) to identify, without further testing:</u> <ul style="list-style-type: none"> - <u>test chemicals inducing serious eye damage (UN GHS Cat. 1); and</u> - <u>test chemicals inducing eye irritation (UN GHS Cat. 2); and</u> - <u>test chemicals that do not require classification for eye hazard (UN GHS No Cat).</u>
<u>Description</u>	<u>The DAL-1 describes the combination of one and/or three physicochemical properties with the results of two in vitro test methods (RhCE and BCOP LLBO) for the identification of the eye hazard potential of non-surfactant liquids. The RhCE models that are part of DAL-1 are the EpiOcular™ Eye Irritation Test (EIT) and the SkinEthic™ Human Corneal Epithelium (HCE) EIT (OECD TG 492, 2023f, see also section 4.3.4). The BCOP LLBO according to the OECD TG 437 is used (OECD, 2023b, see also section 4.3.1). The DIP uses the readout of the prediction models of each of the individual test method as defined by the TGs and/or information on the physicochemical properties. Physicochemical property exclusion rules based on water solubility (WS) or a combination of octanol-water partition coefficient (LogP), vapour pressure (VP) and surface tension (ST) of the neat liquid are used in a first step to identify UN GHS No. Cat. Liquids that are not identified as No Cat. according to the physicochemical property-based</u>

	<p><u>exclusion rules, are then evaluated based on a RhCE test method in Step 2. Liquids that result in a tissue viability > 60% are classified UN GHS No Cat. Liquids that result in a tissue viability ≤ 60% are then evaluated based on the BCOP LLBO test method in a third step. Liquids that result in an opacity > 145 are predicted UN GHS Cat. 1 and the remaining liquids are classified UN GHS Cat. 2 (OECD, 2022a).</u></p> <p><u>The DAL-2 describes the combination of two in vitro test methods (STE: OECD TG 491 and BCOP LLBO: OECD TG 437) for the identification of the eye hazard potential of non-surfactant neat liquids, liquids and solids dissolved in water. The DIP uses the readout of the prediction models of each of the individual test method as defined by the TGs. The STE test method is used to identify UN GHS No Cat. liquid chemicals (liquids that result in a mean cell viability > 70% at a 5% and 0.05% concentration) or to identify UN GHS Cat. 1 liquids (Cat. 1: liquids that result in a mean cell viability ≤ 70% at a 5% and 0.05% concentration). For liquids that result in a mean cell viability ≤ 70% at 5% concentration but > 70% at 0.05%, the BCOP LLBO is needed. Liquids that result in an opacity > 145 are predicted as UN Cat. 1 and the remaining liquids are assigned to UN GHS Cat. 2.</u></p>
<p><u>Scientific basis incl. MoA</u></p>	<p><u>DAL-1</u></p> <p><u>The scientific basis of the individual test methods is described in section 4.3.1. (BCOP), and 4.3.4. (RhCE) of this document.</u></p> <p><u>The use of physicochemical exclusion rules resulted in an increase of the specificity (correct identification of No Cat.). Some liquids were predicted NPCM by the RhCE models but were correctly identified based on the physicochemical properties [(water solubility <0.02mg/mL or a combination of octanol–water partition coefficient (LogP >1), vapor pressure (>3mm Hg), and surface tension (ST <30 dyne/cm)]</u></p> <p><u>DAL-2</u></p> <p><u>The scientific basis of the individual test methods is described in section 4.3.1. (BCOP), and 4.3.3. (STE) of this document.</u></p>
<p><u>Protocol available</u></p>	<p><u>BCOP LLBO</u></p> <p><u>DB-ALM protocol no. 164 on the EpiOcular™ EIT test method (DB-ALM, 2015).</u></p> <p><u>DB-ALM protocol no. 190 on the SkinEthic™ HCE Eye Irritation Test Liquid (EITL) (DB-ALM, 2017).</u></p> <p><u>STE Protocol available at:</u> https://ntp.niehs.nih.gov/whatwestudy/niceatm/test-method-evaluations/ocular/ste</p> <p><u>Analytical methods for measuring the physicochemical properties (GL 104, GL 105, GL 107, GL 115, GL 117, GL 123) + QSAR models with QMRF (Annex E, OECD SD 354, 2022b)</u></p>

<u>Strengths and weaknesses</u>	<p><u>Strengths of the test methods that are part of the DAL-1 and DAL-2 are discussed in the respective chapters of this GD.</u></p> <p><u>Weaknesses</u></p> <ul style="list-style-type: none"> - <u>Mixtures, gases and aerosols have not been assessed yet with the DAs.</u> - <u>The reversibility of tissue lesions cannot be evaluated per se with the DAs.</u>
<u>Identification UN GHS No Category, Category 2, and Category 1</u>	
<u>Applicability domain and limitations</u>	<p><u>Applicability</u></p> <ul style="list-style-type: none"> - <u>The DAL-1 is applicable to neat liquids (OECD, 2022b).</u> - <u>The DAL-2 is applicable to non-surfactant neat liquids, liquids and solids dissolved in water (OECD, 2022b).</u> <p><u>Limitations</u></p> <ul style="list-style-type: none"> - <u>DAL-1 is not applicable for surfactants, solids and liquid mixtures, liquid UVCBs and liquid multi-constituent substances (OECD, 2022b).</u> - <u>DAL-2 is not applicable for surfactants and solids dispersed in water (OECD, 2022b).</u> - <u>Users should refer to the limitations of the individual in vitro test methods as specified in their respective TGs.</u> - <u>Users should refer to the limitations of the individual methods for measuring the physicochemical properties as specified in their respective GLs.</u>
<u>Predictive capacity</u>	<p><u>The DAL-1 with VRM1 (EpiOcular™ EIT) has a balanced accuracy of 68.7% (based on 94 chemicals) with correct predictions of 76.5% for Cat. 1 (based on 17 chemicals), 59.1% for Cat. 2 (based on 22 chemicals) and 70.5% for No Cat. (based on 55 chemicals) (OECD, 2022b).</u></p> <p><u>The DAL-1 with VRM2 (SkinEthic™ HCE EIT) has a balanced accuracy of 75.0% (based on 86 chemicals) with correct predictions of 76.5% for Cat. 1 (based on 17 chemicals), 68.7% for Cat. 2 (based on 23 chemicals) and 79.7% for No Cat. (based on 46 chemicals) (OECD, 2022b).</u></p> <p><u>The DAL-2 has a balanced accuracy of 74.3% (based on 164 chemicals) with correct predictions of 81.2% for Cat. 1 (based on 17 chemicals), 56.3% for Cat. 2 (based on 24 chemicals) and 81.2% for No Cat. (based on 123 chemicals) (OECD, 2022b).</u></p>
<u>Reliability</u>	<p><u>Transferability, within- and between-laboratory reproducibility of these individual test methods have been assessed during their respective validation studies (see 4.3.1., 4.3.3., 4.3.4.)</u></p>

4.3.7. Reconstructed human Cornea-like Epithelium Time-to-Toxicity Test (RhCE EIT) (OECD TG 492B)

44. The OECD TG 492B on the SkinEthic™ Human Corneal Epithelium (HCE) Time-to-Toxicity (TTT) test method was adopted in 2022 (OECD, 2022a). The in vitro test method currently covered by this Test Guideline is the SkinEthic™ HCE TTT which makes use of the commercially available SkinEthic™ Human Corneal Epithelium (HCE/S) tissue construct. The HCE TTT test method underwent a prospective validation study conducted by industry (Alépée et al., 2020, 2021, 2022) and the outcome was peer-reviewed by an independent international peer-review panel (OECD, 2022c). From these evaluations it was concluded that the SkinEthic™ HCE TTT is able to correctly identify chemicals (both substances and mixtures) by discriminating the three UN GHS categories for serious eye damage/eye irritation, i.e. UN GHS Cat. 1, Cat. 2 and No Cat. chemicals.

Module 3 – In vitro data: Reconstructed human Cornea-like Epithelium - Time-to-Toxicity Test (OECD TG 492B)	
General description	
<u>Regulatory use (UN GHS classification)</u>	<u>Eye Hazard identification.</u>
<u>Validation & regulatory acceptance status</u>	<u>Validated and adopted as OECD TG 492B.</u>
<u>Potential role in the IATA</u>	<u>Full-replacement for the in vivo rabbit eye test for regulatory classification and labelling (Figure 2.1) to identify, without further testing:</u> <u>- test chemicals inducing serious eye damage (UN GHS Cat. 1); and</u> <u>- test chemicals inducing eye irritation (UN GHS Cat. 2); and</u> <u>- test chemicals that do not require classification for eye hazard (UN GHS No Cat).</u>
<u>Description</u>	<u>Three-dimensional RhCE tissues are reconstructed from primary human cells, which have been cultured for several days to form a stratified, highly differentiated squamous epithelium morphologically similar to that found in the human cornea. The SkinEthic™ HCE tissue construct consists of at least 4 viable cell layers, including columnar cells and wing cells, with the presence of intermediate filaments, mature hemi-desmosomes and desmosomes, and specific human corneal cytokeratins. The test chemical is applied topically to a minimum of two RhCE tissue constructs. Following the exposure and post-treatment incubation periods, tissue viability is assessed by the enzymatic conversion in viable cells of the vital dye MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; Thiazolyl blue tetrazolium bromide) into a blue MTT formazan salt which is extracted from the</u>

	<p>tissues and quantitatively measured (Mosmann, 1983). The SkinEthic™ HCE TTT Test is based on two protocols, one for liquids (SkinEthic™ HCE TTL) and one for solids (SkinEthic™ HCE TTS). The SkinEthic™ HCE TTT uses three exposure times for TTL and two exposure times for TTS. Liquid chemicals that results in a mean tissue viability $\leq 50\%$ within all three exposure times (5 minutes neat, 16 and 120 minutes with 20% diluted liquid) will be classified as a UN GHS Cat. 1, and liquids that results in a mean tissue viability strictly above 50% will be classified as a UN GHS No Cat. Any other combination of mean tissue viability values will lead to a UN Cat. 2 classification. Solid chemicals that results in a mean tissue viability $\leq 40\%$ after 30 minutes exposure and $\leq 60\%$ after 120 minutes exposure will be classified as a UN GHS Cat 1. A mean tissue viability strictly above these cut-offs within the two time treatments will classify the chemical as a UN GHS No Cat. Any other combination of mean tissue viability values will classify the chemical as a UN GHS Cat 2.</p>
<p><u>Scientific basis incl. MoA</u></p>	<p>See Section 4.3.4.</p>
<p><u>Protocol available</u></p>	<p>SkinEthic™ HCE TTL SOP, Version 1. (June 16, 2020) SkinEthic™ HCE TTS SOP, Version 1. (June 16, 2020)</p>
<p><u>Strengths and weaknesses</u></p>	<p><u>Strengths</u></p> <ul style="list-style-type: none"> - Officially validated test method. - Human-based 3D tissue model. - Quantitative measurements of cell viability. - Controlled exposure conditions, including test chemical concentration and exposure duration. <p><u>Weaknesses</u></p> <ul style="list-style-type: none"> - Gases and aerosols have not been assessed yet in a validation study. - The reversibility of tissue lesions cannot be evaluated per se in the HCE TTT test method.
<p><u>Identification UN GHS No Category, Category 2, and Category 1</u></p>	
<p><u>Applicability domain and limitations</u></p>	<p><u>Applicability</u></p> <ul style="list-style-type: none"> - The SkinEthic™ HCE TTT RhCE test method can be used for the testing of substances and mixtures (OECD, 2022a). - It is applicable to solids, liquids, semi-solids and waxes. Liquids may be aqueous or non-aqueous and solids may be soluble or insoluble in water (OECD, 2022a). <p><u>Limitations</u></p> <ul style="list-style-type: none"> - Test chemicals absorbing light in the same range as formazan dye (FD, naturally or after treatment) and test chemicals able to directly

	<u>reduce the vital dye MTT (to FD) may interfere with the tissue viability measurements and need the use of adapted controls for corrections. However, use of HPLC/UPLC-spectrophotometry as an alternative procedure to measure MTT formazan allows circumventing this, and is especially useful for those test chemicals strongly absorbing in the same wavelength as MTT formazan which are not compatible with the standard optical density (OD) measurement (OECD, 2022a).</u>
<u>Predictive capacity</u>	<u>The SkinEthic™ HCE TTT test has a balanced accuracy of 74.4% (based on 151 chemicals) with correct predictions of 79% for Cat. 1 (based on 50 chemicals), 69% for Cat. 2 (based on 44 chemicals) and 75% for No Cat. (based on 57 chemicals) (OECD, 2022a).</u>
<u>Reliability</u>	<u>The concordance of predictions obtained with the SkinEthic™ HCE TTT test method was 85-95% for TTL (20 chemicals) and 100% for TTS (20 chemicals) within laboratories and 90-100% between laboratories (based on 40 chemicals).</u>

4.3.8. Vitrigel®-Eye Irritancy Test Method (OECD TG 494)

45. The OECD TG 494 on the The Vitrigel®-EIT method was adopted in 2021 (OECD, 2021). The method is an in vitro assay using human corneal epithelium (hCE) models fabricated in a collagen Vitrigel® membrane (CVM) chamber (Yamauchi, et al. 2013). The protocol was developed by Yamauchi and Takezawa (2019), and subject to a validation study organized by JaCVAM in cooperation with the International Collaboration on Alternative Test Methods (ICATM) (Kojima et al., 2019). The validation report was evaluated by an independent peer-review panel composed of international experts (OECD, 2019b). From this evaluation the OECD Expert Group concluded that the Vitrigel®-EIT method is valid for use as an initial step in a bottom-up approach for identification of test chemicals not requiring classification for eye hazard (UN GHS No Cat.).

<u>Module 3 – In vitro data: Vitrigel®-Eye Irritancy Test Method (OECD TG 494)</u>	
<u>General description</u>	
<u>Regulatory use (UN GHS classification)</u>	<u>Identification of test chemicals not requiring classification for eye hazard (UN GHS No Cat.).</u>
<u>Validation & regulatory acceptance status</u>	<u>Validated and adopted as OECD TG 494.</u>
<u>Potential role in the IATA</u>	<u>While the Vitrigel®-EIT is not considered valid as a full-replacement for the in vivo rabbit eye test, it can be used within to initiate the Bottom-Up approach for regulatory classification and labelling (Figure 2.1) to identify, without further testing:</u>

	<p>- test chemicals that do not require classification for eye hazard (UN GHS No Cat).</p> <p>The Vitrigel®-EIT method is not intended to identify nor differentiate between UN GHS Cat. 1 and UN GHS Cat. 2. A test chemical that is not predicted as not requiring classification for eye hazard (UN GHS No Cat.) will thus require additional in vitro testing and/or additional WoE evaluation with other existing information before progressing with further testing within the IATA in order to establish a definitive classification (Figure 2.1).</p>
<u>Description</u>	<p>The Vitrigel®-EIT method uses the destructive activity of the test chemical against the barrier function of hCE models as an endpoint to assess the extent of damage to the hCE model.</p> <p>The test chemical is dissolved or suspended in the culture medium at a concentration of 2.5% w/v. Next, the test chemical is added to a minimum of three hCE models for 3 minutes. The transepithelial electrical resistance (TEER) value of the hCE model cultured in the CVM chamber is measured at intervals of 10 seconds for a period of three minutes after exposure to the test chemical preparation. The eye irritation potential of the test chemical is predicted by analyzing time-dependent changes in TEER values using the score of three indexes. The test chemical is identified as not requiring classification and labelling according to UN GHS (No Cat.) if the scores of the indexes are Time lag > 180 seconds and Intensity < 0.05 %/seconds and Plateau level ≤ 5.0 %.</p>
<u>Scientific basis incl. MoA</u>	<p>It is known that chemicals that are irritating to the eye first destroy tear film and epithelial barrier function of the eye, subsequently induce epithelial cell death, and finally produce stromal degeneration and endothelial cell death, resulting in corneal opacity (Reim, 2013; Movahedan, et al., 2013). Therefore, the change of the epithelial barrier function is a relevant endpoint for detecting eye irritation (Uematsu et al., 2007; Meloni et al., 2010).</p>
<u>Protocol available</u>	<p>Standard Protocol for the Vitrigel-EIT method, Version 1.82e (Yamauchi and Takezawa, 2019)</p>
<u>Strengths and weaknesses</u>	<p><u>Strengths</u></p> <ul style="list-style-type: none"> - Officially validated test methods. - Human corneal epithelium model. - Quantitative measurements of TEER. - Controlled exposure conditions, including test chemical concentration and exposure duration. <p><u>Weaknesses</u></p> <ul style="list-style-type: none"> - The Vitrigel®-EIT test method does not allow discrimination between eye irritation/reversible effects on the eye (UN GHS Cat. 2) and serious eye damage/irreversible effects on the eye (UN GHS Cat. 1), nor

	<p><u>between eye irritants (UN GHS Cat. 2A) and mild eye irritants (UN GHS Cat. 2B). For these purposes, further testing with other suitable test methods is required.</u></p> <p><u>- Mixtures, gases and aerosols, however, have not been assessed in the validation study.</u></p> <p><u>- The reversibility of tissue lesions cannot be evaluated per se in the Vitrigel®-EIT test method.</u></p>
Identification UN GHS No Category	
<u>Applicability domain and limitations</u>	<p><u>Applicability</u></p> <p><u>- The Vitrigel®-EIT test method technically applicable to mono-constituent substances, multi-constituent substances, substances of unknown or variable composition, complex reaction products or biological materials (UVCBs) (OECD, 2021).</u></p> <p><u>- Any test chemical satisfying the criteria of the pre-test (i.e., test chemical preparations showing pH > 5 and keeping dissolution or homogeneous dispersion for at least three minutes) in a 2.5% w/v concentration in culture medium can be tested with the Vitrigel®-EIT method (OECD, 2021).</u></p> <p><u>- Test chemicals that do not dissolve readily can be tested after using one of the following techniques: a) mix mechanically using a vortex mixer, b) sonication, and/or c) heating to a maximum temperature of 70°C (OECD, 2021).</u></p> <p><u>Limitations</u></p> <p><u>- Test chemical preparations of both solids and liquids showing acidity (pH ≤ 5) and rapid phase separation are not in the applicability domain.</u></p>
<u>Predictive capacity</u>	<p><u>The predictive capacity was evaluated based on validation and the developer's in-house data for 93 chemicals (Kojima et al., 2019). The Vitrigel®-EIT method achieved a sensitivity of 83% (50/60), a specificity of 70% (23/33), and an accuracy of 78% (73/93).</u></p>
<u>Reliability</u>	<p><u>The within-laboratory reproducibility was shown to be to be 80–100% at all three laboratories and the between-laboratory reproducibility was 92%.</u></p>

4.3.9. Macromolecular Test Method (OECD TG 496)

46. The OECD TG 496 on the Macromolecular method was adopted in 2019 (OECD, 2023g). The in vitro macromolecular test method Ocular Irritation (OI®) is a biochemical in vitro test method that underwent an independent validation study between 2009 and 2012 (Eskes et al., 2014), followed by an independent peer-review by EURL-ECVAM in 2016 (ESAC, 2016). The assessment of additional data, as recommended by ESAC, regarding the characterisation of the raw material (Macromolecular matrix powder) and its stability over time, was conducted by the OECD expert group. The Ocular Irritation® assay is the first validated macromolecular test assay to identify chemicals inducing serious eye

damage (i.e., UN GHS Cat. 1) and chemicals that do not require classification for eye irritation or serious eye damage as defined by the UN GHS (UN GHS No Cat.).

Module 3 – In vitro data: Macromolecular test method (OECD TG 496)	
General Description	
<u>Regulatory use (UN GHS classification)</u>	Identification of i) test chemicals inducing serious eye damage (UN GHS Cat. 1), and ii) test chemicals not requiring classification for eye hazard (UN GHS No Cat.).
<u>Validation & regulatory acceptance status</u>	Validated and adopted as OECD TG 496.
<u>Potential role in the IATA</u>	<p>While the in vitro macromolecular test method is not considered valid as a full-replacement replacement for the in vivo rabbit eye test, it can be used for regulatory classification and labelling (Figure 2.1) to identify, without further testing:</p> <ul style="list-style-type: none"> - test chemicals inducing serious eye damage (UN GHS Cat. 1); and - test chemicals that do not require classification for eye hazard (UN GHS No Cat.). <p>The in vitro macromolecular test method is recommended as an initial step of a Top-Down approach to positively identify chemicals inducing serious eye damage. (UN GHS Cat. 1) without further testing. The in vitro macromolecular test method is also recommended to identify chemicals that do not require classification for eye irritation or serious eye damage (UN GHS No Cat.), and may therefore be used as an initial step within a Bottom-Up approach.</p> <p>A test chemical that is neither predicted as UN GHS Cat. 1 nor as UN GHS No Cat. with the in vitro macromolecular test method would require additional testing with a UN GHS Cat. 2 method and/or WoE evaluation with other existing information and if still needed additional testing (in vitro and/or in vivo) as a last resort to establish a definitive classification (see Figure 2.1).</p>
<u>Description</u>	<p>The in vitro macromolecular test method is an acellular biochemical test system that consists of two components: a macromolecular matrix and a membrane disc for the controlled delivery of the test chemical to the matrix. The macromolecular matrix serves as the target for the test chemical and is composed of a mixture of proteins, glycoproteins, carbohydrates, lipids and low molecular weight components. Upon rehydration, a complex macromolecular matrix is formed which mimics the highly ordered structure of the transparent cornea (Gordon, 1992; Kelly, 1989). Test chemicals presenting an ocular hazard will produce turbidity of the macromolecular matrix that is quantified by measuring the changes in light scattering (Optical Density (OD) at 405 nm).</p>

	<p>After 24 hours of exposure, OD readings are performed and Irritation Draize Equivalent (IDE) Score for each tested dose/concentration of the test chemical are calculated. The highest IDE Score of the five tested doses/concentrations of a test chemical, namely Maximal Qualified Score (MQS), is then used to determine an UN GHS ocular hazard category.</p> <p>A MQS > 30.0 leads to a UN GHS Cat. 1 prediction; a MQS ≤ 12.5 leads to UN GHS No Cat. prediction. If the MQS result is > 12.5 – 30.0 No final Prediction Can be made (NPCM) on the UN GHS classification.</p>
<u>Scientific basis incl. MoA</u>	<p>Test chemicals causing ocular damage are known to produce denaturation of collagen and saponification of lipids (e.g., by alkalis), coagulation and precipitation of proteins (e.g., by acids) and/or dissolvance of lipids (e.g., by solvents) (Eskes et al., 2010). Test chemicals producing protein denaturation, unfolding and changes in conformation will lead to the disruption and disaggregation of the highly organized macromolecular reagent matrix, and produce turbidity of the macromolecular reagent.</p>
<u>Protocol available</u>	<p>DB-ALM protocol no. 157 on the Ocular Irritation® Assay System (DB-ALM, 2013).</p>
<u>Strengths and weaknesses</u>	<p><u>Strengths</u></p> <ul style="list-style-type: none"> - Officially validated test method. - Quantitative and objective measurements of OD. - Controlled exposure conditions, including test chemical concentration and exposure duration. <p><u>Weaknesses</u></p> <ul style="list-style-type: none"> - The in vitro macromolecular test method is not recommended for the identification of test chemicals that should be classified irritating to eyes (i.e., UN GHS Cat. 2 or Cat. 2A) or test chemicals that should be classified as irritating to eyes (UN GHS Cat. 2B) due to the considerable number of in vivo UN GHS Cat. 1 chemicals underclassified as UN GHS Cat. 2, 2A or 2B and of in vivo UN GHS No Cat. chemicals overclassified as UN GHS Cat. 2, 2A or 2B. For this purpose, further testing with another suitable method may be required with UN GHS Cat. 2 test methods. - The in vitro macromolecular test method does not address cytotoxicity and reversibility aspects of ocular toxicity. - Gases and aerosols have not been assessed yet in a validation study.
<u>Identification of UN GHS Category 1</u>	
<u>Applicability domain and limitations</u>	<p><u>Applicability</u></p>

	<p>- The in vitro macromolecular test method is applicable to solid and liquid chemicals whose 10% solution/dispersion (v/v or w/v as appropriate) has a pH in the range $4 \leq \text{pH} \leq 9$ (OECD, 2023g).</p> <p>- The in vitro macromolecular test method is applicable to substances and mixtures (OECD, 2023g).</p> <p><u>Limitations</u></p> <p>- Intensely coloured chemicals, chemicals which caused salting-out precipitation, high concentrations of some surfactants, and highly volatile chemicals, may interfere with the test system (DB-ALM, 2013).</p>
<u>Predictive capacity</u>	When used for identification of UN GHS Cat. 1 test chemicals, the in vitro macromolecular test method showed an overall accuracy of 75% (66.5/89), a specificity of 81% (55.8/69), and a sensitivity of 54% (10.7/20) calculated based on a weighted approach as compared to in vivo rabbit eye test method (OECD TG 405) data classified according to the UN GHS (OECD, 2023g).
<u>Reliability</u>	The concordance of predictions of the OI [®] test method based on 56 chemicals was found to be 83% between laboratories (Eskes et al., 2014).
<u>Identification UN GHS No Category</u>	
<u>Applicability domain and limitations</u>	<p><u>Applicability</u></p> <p>- The in vitro macromolecular test method is applicable to solid and liquid chemicals whose 10% solution/dispersion (v/v or w/v as appropriate) has a pH in the range $4 \leq \text{pH} \leq 9$ (OECD, 2023g).</p> <p>- The in vitro macromolecular test method is applicable to substances and mixtures (OECD, 2023g).</p> <p><u>Limitations</u></p> <p>- Intensely coloured chemicals, chemicals which caused salting-out precipitation, high concentrations of some surfactants, and highly volatile chemicals, may interfere with the test system (DB-ALM, 2013).</p>
<u>Predictive capacity</u>	When used for the identification of UN GHS No Cat. test chemicals, the in vitro macromolecular test method showed an overall accuracy of 75% (67.0/89), a sensitivity of 91% (41.7/46) and a specificity of 59% (25.3/43) calculated based on a weighted approach as compared to in vivo rabbit eye test method data classified according to the UN GHS (OECD, 2023g).
<u>Reliability</u>	The concordance of predictions of the OI [®] test method based on 56 chemicals was found to be 82% between laboratories (Eskes et al., 2014).

4.4. Module 4: Other existing animal data from non-OECD adopted test methods on serious eye damage and eye irritation

[43-47](#). Existing data from modified OECD TG 405 or in vivo animal test methods adopted by specific countries and/or regulatory authorities similar but not fully compliant with OECD TG 405, shall be considered. Although not fully following the recommendations from the OECD TG 405, existing data obtained from these in vivo animal studies may be useful in giving indication on the potential eye hazard effects of a test chemical. Examples of such in vivo animal test methods include the original Draize test method (Draize et al., 1944), the US FHSA method 16CFR 1500.42 (US CPSC 2015b) and eventual modifications to TG 405. An evaluation shall be made on the degree of similarity and differences of these test methods as compared to the OECD TG 405, and the results used in WoE assessment in Module 9 to support classification and labelling decisions.

[44-48](#). In addition to the above test methods, another non-OECD adopted in vivo animal test is the Low Volume Eye Test (LVET) which involves the application of 1/10th of the amount applied in OECD TG 405 (e.g., 10 µL instead of 100 µL for liquids) directly onto the cornea (instead of into the conjunctival sac) and uses the same scale and the data interpretation as those used in OECD TG 405. Such amount is based on anatomical and physiological considerations indicating that the tear volume in both rabbit and human eyes is approximately the same (~7-8 µL), and that after blinking, the volume capacity in the human eye is ~10 µL (A.I.S.E. 2006). Furthermore, the use of direct cornea exposure mimics human exposure scenarios that can be reasonably expected from e.g. accidental ocular exposure to household detergents and cleaning products. Indeed, the LVET has been mainly used for detergent and cleaning products (Freeberg et al., 1984; Freeberg et al., 1986a,b; Cormier et al., 1995; Roggeband et al., 2000). It was found to still overpredict the effects in man, but to a lesser extent as compared to the classical in vivo rabbit eye test described in OECD TG 405 (Freeberg et al., 1984, 1986b; Roggeband et al., 2000). Following a retrospective validation study and independent peer review, the LVET was not recommended for prospective use, i.e. to generate new data (ESAC, 2009a; ICCVAM, 2010a). Furthermore, although the LVET was considered to have a tendency to classify in lower hazard categories when compared to OECD TG 405 (ICCVAM, 2010a), it was acknowledged that retrospective LVET data may still be useful on a case-by-case basis (e.g. in a WoE approach) to identify potential ocular irritants for the limited use domain of detergent and cleaning products and their main ingredients (i.e., surfactants used in these products) (ESAC, 2009a; ICCVAM, 2010a).

4.5. Module 5: Other data from non-OECD adopted alternative test methods on serious eye damage and eye irritation

[45-49](#). In addition to the OECD adopted in vitro test methods, a number of promising alternative test methods and complementary endpoints exist that may provide with complementary and/or useful information for predicting eye hazard effects. These encompass: (i) histopathology as an additional in vitro endpoint recommended by the OECD GD 160 ([20172018c2018](#)); (ii) test methods that underwent validation studies according to e.g. the OECD GD 34 (2005); and (iii) promising optimized alternative methods for predicting e.g., irreversible effects and UN GHS Cat. 2 classification. [Table 4.3](#) [Table 4.2](#) provides with an overview of these test methods including a description of the endpoints assessed, their proposed application and their validation and regulatory status. Furthermore other non-OECD adopted alternative methods on serious eye damage and eye irritation may also include test methods derived or adapted from OECD adopted in vitro test methods that make use of e.g., i) the same endpoint but measured with a different technology, ii) a new endpoint (in)directly related to the endpoint addressed in the OECD adopted test method(s), and iii) an adapted methodology(s) using the adopted model.

4.5.1. OECD Guidance Document 160 on the use of histopathology as an additional endpoint

~~46-50.~~ Originally adopted in 2011 and further revised in 2017 and 2018, the OECD adopted the Guidance Document ~~No.~~ 160 which provides standard procedures for the collection, fixing and processing of tissues for histological evaluation as an additional endpoint to the BCOP and ICE test methods for eye hazard testing (OECD, ~~20172018e~~2018). The Guidance Document suggests that histopathological evaluation may be useful for (i) assessing histological damage of chemical classes or formulations that are not well characterized in these test methods; (ii) assisting with determination of a mode of action where it cannot be easily predicted; (iii) assisting with determination of the likelihood of delayed effects; (iv) evaluation of the depth of injury, which has been proposed as a measure of reversibility or irreversibility of ocular effects (Maurer et al., 2002); (v) further characterization of the severity or scope of the damage as needed (Harbell et al., 2006; Maurer et al., 2002); and (vi) assisting with discrimination of cases where the response falls along the borderline between two categories based on the standard test method decision criteria. GD 160 mainly addresses the use of histopathology as an additional endpoint to the BCOP and ICE (TG 437 and TG 438) based on the experiences gained so far with these test methods, however, it is conceivable that such endpoint may also be applicable to other tissue models such as the more recently adopted RhCE (TG 492) and the non-OECD adopted IRE test method (see chapter 4.5.2).

Table 4.332. Overview of non-OECD adopted test methods useful in supporting eye hazard identification. Note that this is likely to be non-exhaustive. Furthermore it is recommended to check latest status of those methods under discussions at the OECD level.

Test method	Endpoint(s) assessed	Proposed application by test developer	Validation & regulatory status	Comments
OECD Guidance Document 160 on the use of histopathology as an additional endpoint to the BCOP and ICE test methods				
Histopathology as an additional endpoint	<ul style="list-style-type: none"> - Assisting in determining mode of action, likelihood of delayed persistent effects, depth of injury, and borderline effects in standard ICE and BCOP - Further characterization of chemical classes / formulations not well characterized in BCOP and ICE - Further characterization of the severity or damage - May be useful to other tissue-based methods such as RhCE and IRE 	<p><u>ICE:</u> Identification of UN GHS Cat. 1 detergents and cleaning products.</p>	OECD GD 160	-
RhCE in vitro test methods that underwent validation				
LabCyte CORNEA-MODEL24-EIT	RhCE mo-to assay falling within TG 492	Identification of UN-GHS No Cat. test chemicals	Validated based on performance	Under discussions at the OECD level

Commented [EA5]: This is included in TG 438 as additional endpoint, should we remove this here?

			standards & under peer review	
Vitrigel-EIT method	Barrier function of human corneal epithelium cells cultured in a collagen vitrigel membrane chamber	Identification of UN-GHS No-Cat. test chemicals	Underwent validation & under peer review	Under discussions at the OECD level
EpiOcular time-to-toxicity (ET50) assay	Time of exposure to reduce tissue viability of 50% as compared to the control tissues, using a reconstructed human corneal-like epithelial model	Moderate to mild irritants	Underwent validation & peer review for specific applicability domain Accepted by certain regulatory authorities (US EPA, 2015)	Accepted for testing antimicrobial and cleaning products, when used in combination with BCOP and Cytosensor Microphysiometer
Macromolecular in vitro test methods that underwent validation				
Ocular Irritation	Denaturation of a macromolecular matrix composed of proteins, glycoproteins, lipids and low molecular weight components	Identification of: UN-GHS Cat. 1 test chemicals falling within the applicability domain of the test method UN-GHS No-Cat. test chemicals falling within the applicability domain of the test method, excluding test chemicals having the functional	Underwent validation & peer review Accepted by certain regulatory authorities for the identification of serious eye damage (ECHA, 2015)	Under discussions at the OECD level

		groups — acrylate, carboxamide and cycloalkene		
Cell-based in vitro test methods that underwent validation				
Cytosensor Microphysiometer	Metabolic rate of L929 fibroblasts	Identification of: UN GHS Cat. 1 water-soluble test chemicals UN GHS No Cat. water-soluble surfactants and surfactant-containing test chemicals	Considered scientific valid following peer review Accepted by certain regulatory authorities (ECHA 2015; USA EPA, 2015)	The original version of the apparatus is no longer commercially available at the time of redaction of this GD. A new OECD TG will be considered when new apparatuses showing similar performances as the original apparatus are available
Neutral Red Release (NRR)	Cytotoxicity, measured as release of neutral red dye in monolayer fibroblast cell cultures	Identification of UN GHS No Cat. water soluble test chemicals Proof-of-concept study with a modified prediction model showed the NRR to be useful also for identification of UN GHS Cat. 1 agrochemical formulations (Settivari et al., 2016)	Underwent validation & peer review	Further work was recommended before a statement on the scientific validity of the NRR could be made

Red Blood Cell (RBC) test	Haemolysis of red blood cells, oxyhaemoglobin denaturation (from e.g. calf blood from slaughterhouse, human blood, rabbit blood)	Identification of serious eye damage and no need for classification	Underwent validation & peer review	Further work was recommended before a statement on the scientific validity of the RBC could be made
Organotypic in vitro test methods that underwent validation				
Isolated Rabbit Eye (IRE)	Corneal effects of enucleated rabbit eyes (obtained e.g. from the food chain of from euthanized laboratory rabbits used for other purposes than ocular procedures, providing that no abnormalities are detected in the eyes prior to use)	Identification of UN GHS Cat. 1 test chemicals	Underwent validation & peer review Accepted by certain regulatory authorities for the identification of serious eye damage (ECHA, 2015)	Further work was recommended before a statement on the scientific validity of the IRE could be made
Hen's Egg Test on the Chorio-Allantoic Membrane (HET-CAM)	Haemorrhage, lysis and coagulation of blood vessels of the chorioallantoic membrane (CAM) of fertilized chicken eggs	Identification of: UN GHS Cat. 1 test chemicals (based on coagulation) UN GHS No Cat. test chemicals (based on coagulation, haemorrhage and lysis)	Underwent validation & peer review Accepted by certain regulatory authorities for the identification of serious eye damage (ECHA, 2015)	Further work was recommended before a statement on the scientific validity of the HET-CAM can be made Depending upon the regulatory context, this assay may be considered an animal test

Chorio-Allantoic Membrane Vascular Assay (CAMVA)	Haemorrhage, hyperaemia and constriction of blood vessels of the chorioallantoic membrane (CAM) of fertilized chicken eggs	Moderate to mild irritants	Underwent validation	Further work required to evaluate the scientific validity of the test method
Other promising assays				
Porcine Ocular Cornea Opacity/Reversibility Assay (PorCORA)	Reversibility of cornea injury in air-interface ex vivo porcine corneas cultured for 21 days	Proposed for identification of Serious Eye Damage based on persistence/reversibility of effects as well as severity of effects and a better discrimination between Cat. 1 and Cat. 2	Optimised	-
Ex Vivo Eye Irritation Test (EVEIT)	Reversibility of epithelial and stromal damage of isolated corneas from rabbit eyes (obtained from the food chain) cultured at air-liquid interface for 72 hours	Proposed for the assessment of both severity and persistence/reversibility of ocular lesions, allowing to discriminate between all UN GHS categories (No Cat., Cat. 2A/B and Cat. 1)"	Optimised	-

3D hemi-cornea	Cytotoxicity and/or depth of injury of a multilayered human-based epithelium and stroma with embedded keratocytes cultured in a collagenous matrix	Proposed for identification of all UN GHS categories (Cat. 1, Cat. 2A/2B and No Cat.)	Optimised	-
Slug mucosal irritation (SMI) assay	Mucus produced from the mucosal surface of slugs	Proposed for identification all UN GHS categories (No Cat., Cat. 2A/B and Cat. 1)	Optimised	Depending upon the regulatory context, this assay may be considered an animal test

DRAFT 6/10/2017

~~47-51.~~ In particular, the use of histopathology as an additional endpoint to the ICE test method was found to decrease the rate of false negatives (as observed with the ICE test method when used as a stand-alone) for the identification of UN GHS Cat. 1 for the limited applicability domain of non-extreme pH detergent and cleaning products (OECD, 2018b; Cazelle et al., 2014, 2015). Interestingly, these mixtures were mostly classified in vivo UN GHS Cat. 1 due to persistence of effects, i.e. mild ocular effects that persisted over the 21 day observation period in the tested rabbits. The authors developed a decision criteria for identification of UN GHS Cat. 1 based on semi-quantitative histopathological observations (Prinsen et al., 2011) in which epithelial vacuolation (in the mid and lower layers) and epithelial erosion (of at least moderate level) were found to be the most typical histopathological effects induced by UN GHS Cat. 1 non-extreme pH formulations ($2 < \text{pH} < 11.5$) detergents that were classified in vivo mainly due to persistence of effects (Cazelle et al., 2014). Use of such criteria for non-extreme pH ($2 < \text{pH} < 11.5$) detergents and surfactants detergent and cleaning formulations that were identified by the standard ICE test method as 'no prediction can be made', allowed to decrease the rate of Cat. 1 false negatives observed with the ICE test method alone whilst maintaining a good accuracy and an acceptable specificity (OECD, 2018b; Cazelle et al., 2014). Following demonstration of reproducibility between pathologists and peer-reviewers from three independent laboratories of (10/12 or 83%) and over time (17/18 for non-extreme pH detergents and 6/6 for surfactants), the use of ICE histopathological criteria was included within the OECD TG 438 (2018a) for the limited applicability domain of non-extreme pH ($2 < \text{pH} < 11.5$) detergents and surfactants. However, to ensure such reproducibility, there is a need for (i) an internal peer-review system to be in place; (ii) assessment of the original slides in order to enable the evaluation of three dimensional effects; and (iii) appropriate training & proficiency appraisal. Furthermore, appropriate and relevant data are needed to verify and expand the applicability of the ICE histopathology decision criteria to other chemistries.

~~48-52.~~ Furthermore, recent studies on the BCOP test method suggest that histopathology might be useful in predicting in vivo ocular irritation, particularly for test chemicals with $3 < \text{IVIS} \leq 25$ that would be classified as mild irritants (Cat. 2B) according to the UN GHS (Furukawa et al., 2015). The authors showed that corneal epithelial lesions caused by Cat. 2B test chemicals were localized on the border between the corneal epithelium and stroma.

4.5.2. In vitro test methods that underwent validation studies

~~49-53.~~ Methods that underwent validation studies according to e.g. the OECD GD 34 (2005), encompass reconstructed human tissue models, organotypic test methods, cell based assays and a macromolecular test method. These test methods as well as additional test methods may become available for addressing eye hazards, therefore it is advised to always check the latest status of these test methods on the OECD website .

4.5.2.1. Reconstructed human tissue models

~~50-54.~~ The **LabCyte CORNEA-MODEL** is a RhCE model that underwent a performance-based validation study according to the OECD GD-216 (2015cb). Furthermore, results obtained on 61 test chemicals showed good predictive capacity of the test method (Katoh et al., 2013). It has been proposed as a me-too assay to the RhCE test method falling within the OECD TG 492, and at the time of the redaction of this document, it is currently under peer-review.

~~54-55.~~ The **Vitrigel-eye irritancy test method** is a RhCE based assay which assesses the effects of test chemical on the barrier function of human corneal epithelium cells cultured in a collagen vitrigel membrane. Prediction of UN GHS No Cat. is based on a time-dependent profile of transepithelial electrical resistance assessed for 3 min after exposure to the test chemicals. A total of 118 chemicals have been tested, and when test chemicals having a $\text{pH} \leq 5$ are removed from the applicability domain, the assay showed

performances in line with the adopted test methods for the prediction of UN GHS No Cat. test chemicals (Yamaguchi et al., 2016). The assay underwent a formal validation study and at the time of the redaction of this document, is currently under peer-review.

52-56. Finally, **the EpiOcular time-to-toxicity (ET50) assay** is a RhCE assay in which the eye hazard effects are evaluated by the time necessary to reduce tissue viability to e.g. 50% (in contrast to the decrease in cell viability with a fixed exposure time recommended in OECD TG 492). The assay underwent validation studies focusing on surfactant ingredients and a limited number of formulations (Blazka et al., 2000, 2003). It further underwent a peer-review when used as a part of a testing strategy together with the BCOP and Cytosensor Microphysiometer test methods, to evaluate anti-microbial cleaning products (ICCVAM, 2010c). Such test strategy was accepted by the US EPA for determining the appropriate eye hazard classification for antimicrobial cleaning products (US EPA, 2015). Further work evaluated the usefulness of this assay, when combined with the NRR to evaluate the eye hazard potential of agrochemical formulations (Settivari et al., 2016). The data set was enlarged with 80 chemicals in the CON4EI project (Kandarova et al., 2017b).

4.5.2.2. Macromolecular assays

53. The Ocular Irritation (OI) assay is based on a macromolecular reagent produced from a biological extract that is composed of proteins, glycoproteins, lipids and low molecular weight components that self-associate to form a complex matrix. Eye hazard is assessed based on the premise that irritant test chemicals will lead to protein denaturation and disaggregation of the macromolecular matrix. The changes in protein structure result in changes in turbidity which are measured at an OD of 405 nm. The assay underwent a prospective and a retrospective validation study (Eskes et al., 2014), in which the test method is proposed to identify test chemicals falling within its applicability domain (both substances and mixtures) inducing serious eye damage (UN GHS Cat. 1) as well as those not requiring classification for eye hazard (UN GHS No Cat.). The test method showed good within-laboratory variability including transferability to a naïve laboratory, and between-laboratory based on concordance of classifications. When used for the identification of UN GHS Cat. 1 versus other categories, and for the identification UN GHS No Cat. versus classified materials, excluding the functional groups acrylate, carboxamide and cycloalkene, the test method showed accuracy, false negative and false positive rates which were in line with currently adopted test methods for that purpose (Eskes et al., 2014). The outcome of the validation study was subsequently evaluated by EURL ECVAM and peer reviewed by the ESAC (ESAC, 2016b), in which a few technical issues were identified, which are currently under discussion at the OECD level. Furthermore, the OI assay is accepted by certain countries for the prediction of serious eye damage (UN GHS Cat. 1) (ECHA, 2015).

4.5.2.3. Cell-based assays

54-57. A draft OECD Test Guideline has been proposed on the **Cytosensor Microphysiometer (CM)** test method (OECD, 2012d/2012b). The CM has been evaluated in a retrospective validation study coordinated by EURL ECVAM in collaboration with US ICCVAM and JaCVAM (EURL ECVAM, 2008b), followed by peer review by the ESAC (ESAC, 2009b). From these evaluations it was concluded that the test method can correctly identify water-soluble test chemicals (both substances and mixtures) inducing serious eye damage (UN GHS Cat. 1) as well as water-soluble surfactants and surfactant-containing test chemicals not requiring classification for eye hazard (UN GHS No Cat.). The assay is performed on a sub-confluent monolayer of adherent mouse L929 fibroblasts cultured in a sensor chamber using a pH-meter to detect changes in acidity (Harbell et al., 1997). The

rate of change in acidity (per unit time) measured during the assay serves as a read-out to determine the metabolic rate of the population of cells. If a test chemical causes cytotoxicity to this population of cells, it is assumed that the metabolic rate will fall. The concentration of a test chemical that leads to a 50 % decline (MRD50) in the basal metabolic rate of the population is the parameter used to indicate cytotoxic effects. Identification of water-soluble test chemicals inducing serious eye damage (UN GHS Cat. 1) is triggered by an MRD50 \leq 2 mg/ml whereas UN GHS No Cat. water-soluble surfactants and surfactant containing mixtures are identified by an MRD50 \geq 10 mg/ml (OECD, [2012a](#); [2012b](#)). The CM may also address questions of cell metabolism and recovery. However, the assay requires the use of a Cytosensor Microphysiometer instrument, and at the time of redaction of this GD the original version of this apparatus is no longer commercially available so that the implementation of the assay with newly acquired original apparatus is not possible. Nevertheless, similar me-too apparatus are being commercialised but these have not been validated yet. Adoption of an OECD TG on the CM will be considered when new apparatuses are available that show similar performances to the original version. However, the Cytosensor Microphysiometer is accepted by certain regulatory authorities (ECHA, 2015; US EPA, 2015).

55-58. The **Neutral Red Release (NRR)** is based on near-confluent monolayer cell cultures, and assesses the eye hazard effects of test chemicals by exposure to serial dilutions of test chemicals for 1 to 5 minutes. The concentration of test chemical producing a 50% release of pre-loaded neutral red dye is obtained by extrapolation from the dose-response curve and used to predict eye hazard. The NRR test method has been evaluated in a retrospective validation study coordinated by EURL ECVAM in collaboration with US ICCVAM and JaCVAM (EURL ECVAM, 2008b), followed by peer review by the ESAC (ESAC, 2009b). However, further work was recommended before a statement on the scientific validity of the NRR could be made including to test additional number and classes of chemicals, and to obtain more data on between-laboratory variability (ESAC, 2009b). Although not formally endorsed as scientifically valid, the NRR test method was considered promising by the Validation Management Group for the identification of UN GHS No Cat., water-soluble test chemicals (EURL ECVAM, 2008b; ESAC 2009c). Furthermore, a recent proof-of-concept study making use of a modified prediction model, suggested the test method to be useful also for the identification of UN GHS Cat. 1 agrochemical formulations (Settivari et al., 2016).

56-59. The **Red Blood Cell (RBC)** haemolysis test is based on the potential of a test chemical to disrupt cell membranes as assessed by measuring photometrically the leakage of haemoglobin from freshly-isolated red blood cells incubated with the test chemical under standard conditions (Muir et al., 1983; Pape et al., 1987, Pape & Hope 1990; Pape et al., 1999; Lewis et al., 1993). The denaturation (i.e. change in protein configuration) of oxyhaemoglobin is used as second toxicological endpoint. Mammalian erythrocytes might be obtained through e.g. slaughterhouse material. The RBC test method has been evaluated in a retrospective validation study coordinated by EURL ECVAM in collaboration with US ICCVAM and JaCVAM (EURL ECVAM, 2008b), followed by peer review by the ESAC (ESAC, 2009b). The evidence then available was considered insufficient to support a recommendation on the RBC's scientific validity for regulatory use. In particular a more consistent dataset was deemed necessary to improve confidence on the RBC's applicability domain (ESAC, 2009b, 2009c).

4.5.2.4. Organotypic assays

57-60. The **Isolated Rabbit Eye (IRE)** is based on the same principles as the ICE test method, but instead of chicken eyes it uses enucleated rabbit eyes (obtained from the food chain or from euthanized laboratory rabbits, providing that the animals have not previously been used for ocular procedures, and that no abnormalities are detected in the eyes prior to use in the IRE by e.g. slit-lamp examination), instead of chicken eyes. The effects of test chemicals are assessed by evaluating the corneal thickness (swelling), corneal opacity, area of corneal involvement, fluorescein penetration and morphological changes to the

corneal epithelium. Similar to BCOP and ICE, histopathology may be used as an additional endpoint. Furthermore, confocal microscopy may be used to determine the extent and depth of ocular injury. The IRE test method underwent retrospective validation by ICCVAM (ICCVAM 2006, 2010b) in which further work was recommended before a statement on its scientific validity could be made. The main reason was the fact that several endpoints and protocols for the IRE were applied and evaluated, each with insufficient data provided to make a sound conclusion (ESAC, 2007). Despite this, the IRE continues to be used (Guo et al., 2010) and is accepted by certain countries for the prediction of serious eye damage (UN GHS Cat. 1) (ECHA, 2015). However, depending upon the regulatory context, this assay may or may not be considered as an animal test.

58-61. The Hen's Egg Test on the Chorio-Allantoic Membrane (HET-CAM) is an assay that allows evaluating vascular effects. It makes use of the chorioallantoic membrane (CAM) of fertilized chicken eggs, a vascular foetal membrane composed of the fused chorion and allantois. The acute effects induced by a test chemical on the small blood vessels and proteins of this soft tissue membrane can be used as indicator of ocular effects induced by the test chemical (ICCVAM, 2010b). This characteristic makes the HET-CAM particularly suited to predict conjunctival injury and effects to the vascular system. The test chemical is applied directly to the CAM of fertilized hen eggs, and acute effects such as haemorrhage, lysis of blood vessels and coagulation are assessed. The test method is accepted by certain countries for the identification of serious eye damage (UN GHS Cat. 1) (ECHA, 2015) although further work was recommended before a statement on its scientific validity could be made (ICCVAM 2006, 2010b). One potential reason for such outcome is the existence of a variety of protocols and prediction models used for the same test method. A workshop organized in 2012 by the German Federal Institute for Risk Assessment (BfR), the European Partnership for Alternative Approaches to Animal Testing (EPAA) and Services and Consultation on Alternative Methods (SeCAM) have made recommendations on the most suitable endpoints and protocols to be used either for the identification of UN GHS Cat. 1 or for the identification on UN GHS No Cat. Briefly, for the identification of serious eye damage (UN GHS Cat. 1), coagulation was the recommended endpoint based either on the mean time to develop coagulation or on the severity of coagulation observed at a single time after exposure (Spielmann et al., 1991; Steiling et al., 1999). For the identification of test chemicals not requiring classification (UN GHS No Cat.), the evaluation of coagulation, haemorrhage and lysis at different fixed time points (0.5, 2 and 5 min) was recommended (Luepke, 1985), based on the IS(a) prediction model (ICCVAM, 2010b). The necessity of re-considering the validation status of the method was also raised during this workshop in order to re-analyze the HET-CAM predictive capacity (for the identification of both UN GHS Cat. 1 and UN GHS No Cat.) taking into account the new data generated since 2009. In particular a new validation study has been initiated in 2015 by the Brazilian Centre for the Validation of Alternative Methods (BraCVAM) to complete such dataset. Due to the fact that the HET-CAM method uses live chick embryos, depending upon the regulatory context, this assay may or may not be considered as an animal test. The HET-CAM is accepted by certain countries for the prediction of serious eye damage (UN GHS Cat. 1) (ECHA, 2015).

59-62. The Chorioallantoic Membrane Vascular Assay (CAMVA) is another assay that also assesses the potential hazard effects of test chemicals to the blood vessels of the CAM. In preparing for the test, a small opening is cut into the shell of the egg four days after fertilisation and a small amount of albumen is removed, to permit optimal growth of the CAM. On day 10, the test substance is applied directly onto a small area of the CAM, and after exposure for 30 minutes, the eggs are examined for any vascular change to the CAM, such as haemorrhaging or hyperaemia (capillary injection) or the occurrence of vessels devoid of blood flow (ghost vessels). The concentration of a test material eliciting such damaging effects in 50 % of the treated eggs is used to predict eye hazard. The CAMVA has been included in a number of validation studies (for review see Eskes et al., 1995), and has mostly been applied to the assessment of materials in the mild-to-moderate irritation range (Cerven and Moreno, 1998). However, the CAMVA has not been assessed in parallel by more than two or three laboratories; thus, larger-scale validation or a

retrospective validation based on the existing data is required, in order to further evaluate the scientific validity of the test method (Brantom et al., 1997; Bagley et al., 1992, 1999).

4.5.3. Promising optimized in vitro test methods

~~60-63.~~ A number of assays have been developed to address mechanisms of action not covered by the currently accepted test method. Perhaps the most important mechanism is the discrimination of reversible vs. irreversible effects. As described in chapter 4.2.2, persistence of effects appears as a major driver for UN GHS Cat. 1 classification that may not be directly predicted by the currently adopted ex vivo and in vitro test methods. Two test methods have been developed specifically to address this mechanism of action, the Porcine Cornea Opacity/Reversibility Assay and the Ex Vivo Eye Irritation Test. The **Porcine Cornea Opacity/Reversibility Assay (PorCORA)** assay makes use of an air-interface culture system to sustain ex vivo porcine corneas in culture for 21 days (similar to the in vivo observation period described in TG 405), and determines whether cornea injury once inflicted will reverse (Piehl et al., 2010). Corneal injury reversibility is measured using Sodium Fluorescein stain to detect compromised epithelial barrier function. The test method was shown to identify test chemicals causing both reversible and irreversible serious eye damage in the in vivo rabbit eye test based on 32 tested UN GHS Cat. 1 test chemicals (Piehl et al., 2011). The **Ex Vivo Eye Irritation Test (EVEIT)** in contrast is based on isolated corneas from rabbit eyes (slaughtered for food process), cultured in an air-liquid interface in conditions that allow maintenance of the normal physiological and biochemical functions of the entire rabbit cornea in vitro for 72 hours after sample application. Effects on cornea and reversibility of epithelial and stromal damage are assessed at 24h, 48h and 72h after test chemical application, by measuring corneal opacity (by macroscopic imaging in combination with fluorescein staining) as well as depth of damage, corneal thickness and structural changes assessed by the non-invasive Optical Coherence Tomography technique. A prediction model has been developed to identify the three UN GHS Categories of eye hazard (Spöler et al., 2015).

~~61-64.~~ In addition, two test methods have been suggested for the discrimination of the entire range of ocular hazards, including the UN GHS Cat. 2 classification i.e., the 3D hemi-cornea model, and the slug mucosal irritation (SMI) assay. The **3D hemi-cornea model** is a new model comprised of a multilayered RhCE and a stroma with embedded human corneal keratocytes in a collagenous matrix for which two different test approaches are proposed. The first approach quantifies the cytotoxicity within the epithelium and the stroma separately and uses both values obtained, based on pre-defined thresholds for each compartment, to predict the potential eye hazard (Bartok et al., 2015). The second approach quantifies the cytotoxicity by microscopically assessing the depth of injury within the hemi-corneal tissue (Zorn-Kruppa et al., 2014). Preliminary results showed the capacity of the two approaches to differentiate UN GHS Cat. 1 from UN GHS Cat. 2 test chemicals based on 30 chemicals tested with the first approach, 25 chemicals tested with the second approach, and 14 chemicals tested in both approaches covering the entire range of eye hazards (Bartok et al., 2015; Zorn-Kruppa et al., 2014; Tandon et al., 2015). The same studies support the initial approach to differentiate between all 3 GHS categories, although the selectivity of both methods still must be improved before they can be used as stand-alone methods. The successful method transfer has been demonstrated in a ring trial with both approaches (Mewes et al., 2017). The slug mucosal irritation assay in contrast predicts the eye hazard of test chemicals based on the protein release from the mucosal surface the amount of mucus produced by the of slugs (*Arion lusitanicus*). A pre-validation study was conducted with four participating laboratories and the testing of 20 chemicals covering the entire range of eye hazards. All UN GHS No Cat. were predicted correctly by the four laboratories. Furthermore, identification of both UN GHS Cat. 2 and UN GHS Cat. 1 showed good predictivity (Adriaens et al., 2005, 2008). The assay was also shown to be promising to predict ocular discomfort caused by shampoos (Lenoir et al., 2011). The data set was enlarged with 80 chemicals (Adriaens et al. 2017d) and this assay is incorporated in a three-tiered strategy using use an RhCE test method (EpiOcular™ EIT or SkinEthic™

HCE EIT) at the bottom (identification No Cat.) in combination with the BCOP and SMI at the top (identification Cat. 1) (Adriaens et al., 2017a). However, depending upon the regulatory context, this assay may or may not be considered as an animal test.

4.6. Module 6: Existing human, in vivo and in vitro data on skin corrosion

~~62-65~~. Existing human, in vivo and in vitro data generated on skin corrosion should be taken into account, such as those derived from an Integrated Approach on Testing and Assessment for Skin Corrosion and Irritation (OECD, 2014a). If sufficient and adequate quality data exists to assign Skin Corrosive Cat. 1, 1A, 1B or 1C, the risk of serious damage to eyes is considered implicit (UN GHS Cat. 1).

4.7. Module 7: Physico-chemical properties (existing, measured or estimated)

~~63-66~~. Test chemicals having $\text{pH} \leq 2.0$ or $\text{pH} \geq 11.5$ are predicted to be corrosive to skin or cause serious eye damage (UN GHS Cat. 1). However, where extreme pH is the only basis for classification as serious eye damage, it may also be important to take into consideration the acid/alkaline reserve i.e., a measure of the buffering capacity of a test chemical, especially for classification of mixtures containing acidic or alkaline substances (Young et al., 1988).

~~64-67~~. The determination of pH should be performed following OECD TG 122 (2013) ~~e2013e~~ ~~2013e~~ ~~2013e~~. This Test Guideline also describes procedures to determine acid reserve or alkali reserve for test chemicals that are acidic ($\text{pH} < 4$) or alkaline ($\text{pH} > 10$) by titration with standard sodium hydroxide or sulphuric acid solution using electrometric endpoint detection.

~~65-68~~. However, the pH or p_H in combination with buffering capacity should not be used alone to exonerate from serious eye damage classification. Indeed, when the pH or p_H in combination with acid/alkaline reserve suggest that the test chemical might not induce serious eye damage, further in vitro testing should be considered.

~~66-69~~. Other physico-chemical properties such as melting point, molecular weight, octanol-water partition coefficient, surface tension, vapour pressure, aqueous solubility and lipid solubility, may also be used to identify chemicals not likely to cause such adverse health effects (Gerner et al., 2005; Tsakovska et al., 2005). Such physico-chemical parameters may be measured or estimated using non-testing methods (see module 7), e.g., (Q)SARs, and may be used to help orient chemicals to a Top-Down or Bottom-Up approach in Part 3 of the IATA (Figure 2.1).

Module 7 – Data on physico-chemical properties: Extreme pH	
General description	
Regulatory use (UN GHS classification)	Prediction of serious eye damage (UN GHS Cat. 1)
Validation & regulatory status	Not formally validated but accepted as part of IATA.

Potential role in the IATA	Useful to identify test chemicals with potential to induce serious eye damage. However, the pH or pH in combination with buffering capacity should not be used alone to exonerate from serious eye damage classification. Indeed, when the pH or pH in combination with acid/alkaline reserve suggest that the test chemical might not induce serious eye damage, further in vitro testing should be considered.
Description	pH measurement (considering buffering capacity, if relevant).
Scientific basis incl. MoA	Test chemicals exhibiting extreme pH (either $\text{pH} \leq 2.0$ or $\text{pH} \geq 11.5$), with high buffering capacity when relevant, are likely to produce necrosis to the eyes.
Protocol available	OECD TG 122 (2013e2013e2013b) describes the procedure to determine pH, acidity and alkalinity of aqueous solutions or aqueous dispersions having a $\text{pH} \leq 14$.
Strengths and weaknesses	<p><u>Strengths</u></p> <ul style="list-style-type: none"> - Simplicity. - Low cost. <p><u>Weaknesses</u></p> <ul style="list-style-type: none"> - No information available on the test method reliability (reproducibility). - Predicts serious eye damage induced by pH effects but not by other mechanisms. - There are known cases of test chemicals with extreme pH that do not induce serious eye damage and therefore, use of pH information alone for deciding on Cat. 1 classification may lead to overclassification.
Identification of UN GHS Category 1	
Applicability domain and limitations	<p><u>Applicability</u></p> <ul style="list-style-type: none"> - Although OECD TG 122 allows pre-treatment with acetone to avoid plugging of the electrodes, some test chemical properties, such as low water solubility or rapid hydrolysis, might impair pH measurements. <p><u>Limitations</u></p> <ul style="list-style-type: none"> - For extreme pH mixtures having low or no buffering capacity suggesting the mixture may not be corrosive despite the low or high pH value, the non-corrosive classification still needs to be confirmed by other data (preferably by data from an appropriate validated in vitro test method).

4.8. Module 8: Non-testing data on serious eye damage and eye irritation

4.8.1. (Q)SAR, expert systems, grouping and read-across (substances)

~~67~~-70. Non-testing methods can be used if they provide adequate, relevant and reliable data for serious eye damage and eye irritation for the substance of interest. For substances, the non-testing methods can be divided into two different categories:

- -Read-across using grouping of substances , and
- -Qualitative and quantitative Structure-Activity-Relationships ((Q)SAR) as well as expert and other prediction systems that often incorporate multiple SARs, QSARs, expert rules and/or data.

~~68~~-71. With the introduction of the OECD (Q)SAR Toolbox in combination with the eChemPortal , useful tools are provided for:

- -Finding existing data on the substance under question (target),
- -Identifying analogues for potential read-across and grouping and finding existing data on these analogues,
- -Applying a number of SARs and other profilers for serious eye damage and irritation to the target structure,
- -Grouping and deriving simple (Q)SAR or trend relationships.

~~69~~-72. Guidance on how to apply (Q)SARs for regulatory use and on how to assess the validity and suitability of (Q)SAR models and adequacy of their predictions is provided in the OECD GD 69 (OECD, 2007) and is also available from the corresponding section of the OECD website-. Other useful guidance documents have also been published to aid in determining how and when to apply QSAR models-. Together, these resources can help inform a determination of whether a (Q)SAR result might be used to replace a test result. Furthermore, examples of how to build and report grouping of substances and read-across are also available-.

70-73. The mechanism of serious eye damage/eye irritation involves toxicodynamic and toxicokinetic parameters. Some (Q)SAR models predict serious eye damage and eye irritation based on toxicodynamic properties only (e.g. acidity or basicity, electrophilicity, other reactivity, surfactant activity, membrane destruction). These models have to be checked whether they also take into account, or have to be used in combination with models covering toxicokinetic parameters such as potential of a substance to cross relevant outer membranes of the eye (cornea) and to be active in the living tissue underneath. Conversely, some (Q)SAR models predict (the absence of) serious eye damage and/or eye irritation solely from e.g. physico-chemical properties considered to illustrate the toxicokinetic behaviour of a substance and have to be checked whether they also take into account, or to be used in combination with models relying on toxicodynamic properties. Ideally, such models would also take into account the potential for metabolism, autoxidation, or hydrolysis of the parent compound and how that might impact any effects on the eye.

74-74. For example, the BfR rule-base implemented in Toxtree and the OECD QSAR Toolbox contains both physico-chemical exclusion rules and structure-based inclusion rules (structural alerts). Evaluations of these rules for the prediction/exclusion of eye irritation have been carried out in accordance with the OECD principles for (Q)SAR validation (Tsakovska et al., 2005, on structural alerts; Tsakovska et al., 2007, on physico-chemical exclusion rules). However, inclusion and exclusion rules were evaluated separately, and not used in combination in these works.

72-75. When applied, these two sets of rules may sometimes provide contradictory information, i.e. a structural alert might indicate serious eye damage and/or eye irritation potential, while at the same time, based on physico-chemical properties, absence of effect is predicted. In such cases, it is recommended to consider additional information (e.g. on

the behaviour of chemically similar substances). In other cases, applicability of one (or more) of the physico-chemical exclusion rules might indicate absence of serious eye damage and/or eye irritation potential of the target substance, while no structural alert for serious eye damage and/or eye irritation is triggered. Given that the absence of any known structural alert is not equivalent to the absence of a potential effect, in such a situation the substance should still be examined for potentially reactive substructures (and examining the behaviour of chemical analogues would still be beneficial).

~~73-76~~. While these considerations apply to the use of the BfR rule-base for direct classification/non-classification, less certainty might be required for e.g. a decision on further in vitro testing i.e., where the exclusion rules suggest the absence of an effect, a Bottom-Up approach could be followed (see Figure 2.1).

4.8.1.1. SARs, grouping and read-across for serious eye damage and eye irritation

~~74-77~~. Read-across, SARs and Grouping/Category formation are treated together because they represent approaches based on the same basic concept. Note that, depending on the legal framework and Member Country, specific requirement may be associated to the read-across and grouping approaches. For example, under the EU REACH Regulation, read-across needs to be justified, documented, and supported by reliable data on the source(s), i.e. one or more substances (ECHA, 2015).

~~75-78~~. Toxicological data gaps for a chemical can be filled by prediction based on similar chemicals for which test data are available. While this has historically been accomplished based on structure and physico-chemical properties, mechanistic (biological) similarity is increasingly being used to add confidence to this process. Efforts are ongoing to develop consensus on applying these principles to facilitate their effective use in regulatory context (Ball et al., 2016; Zhu et al., 2016). Structural alerts are substructures in the substance that are considered to reflect chemical or biochemical reactivity underlining the toxicological effect. The occurrence of a structural alert for a substance suggests the presence of an effect, and structural analogues that have exhibited serious eye damage or irritation potential can be used to predict serious eye damage or eye irritation effects of the substance of interest, or be used to tailor further testing and assessment, as indicated in the Figure 2.1. Structural alerts for serious eye damage/eye irritation have been described in the literature, e.g. in Gerner et al. (2005).

~~76-79~~. The similarity of two substances can also be based for example on a common functional group, common precursors or common break-down products (analogue approach). Grouping requires that toxicological properties of the target substance may be predicted from the data of the source substances, basically by interpolation. Predictions based on read-across may therefore be possible for chemically similar substances if it can be shown that their similarity reflects reactive substructures able to react with ocular tissue, even if that substructure has so far not been coded into a structural alert in any of the available literature or software models. Indeed, knowledge on structural alerts for serious eye damage and irritation is always evolving (in particular where new classes of substances are introduced into the market).

~~77-80~~. While not typically useful for regulatory decisions in isolation, negative data can be useful in certain cases. In these cases it is helpful to consider both the structural and mechanism of action similarity along with applicability domain.

4.8.1.2. QSARs and expert systems for serious eye damage and eye irritation

~~78-81~~. An overview of the available (Q)SARs for serious eye damage/eye irritation is provided in [Table 4.4](#)[Table 4.43](#)[Table 4.3](#), and more details can be retrieved in published reviews (ECHA, 2015 – appendix R.7.2-3; Gallegos Saliner et al., 2006, 2008). Furthermore, in recent studies QSAR models based on multiple artificial neural network molecular descriptor selection functionalities were developed, to maximize the applicability domain of the battery for the assessment of both eye irritation and serious eye damage

potential (Verma and Matthews, 2015a, 2015b). The same authors developed an in silico expert system based upon exclusion rules of physico-chemical properties to facilitate the rapid screening and prioritization of test chemicals (Verma and Matthews, 2015c). Predictions from multiple QSAR models in a weight of evidence also allows enhancing the confidence in the prediction.

~~79-82~~ Expert systems are computer programs that guide hazard assessment by predicting toxicity endpoints of certain substance structures based on the available information. They can be based on automated rule-induction systems (e.g., TopKat and MultiCASE), or on a knowledge-based system (e.g. Derek).

~~80-83~~ The freely downloadable OECD QSAR Toolbox software contains two profilers relevant for serious eye damage/eye irritation based on the BfR rule-base. This rulebase is based on the combined use of two predictive approaches: a) physicochemical exclusion rules to identify chemicals with no eye irritation/serious eye damage potential; and b) structural inclusion rules (SARs) to identify chemicals with eye irritation/serious eye damage potential (Gerner et al., 2005). The use of a combination of profilers and data for analogues could allow for the prediction of serious eye damage/eye irritation for new substances through a read-across or category approach.

~~84-84~~ Not all of the models were developed for the purpose of UN GHS classification, so that it is important to assess in each case whether the endpoint or effect being predicted corresponds to the regulatory endpoint of interest.

4.8.2. Bridging principles and theory of additivity (mixtures)

~~82-85~~ Non-testing methods for mixtures can be divided into (UN, 2015):

- Bridging, when data are not available for the complete mixture, and
- Theory of additivity, when data are available for the ingredients of the mixture.

~~83-86~~ Bridging principles are used when the mixture itself has not been tested for serious eye damage and irritation, but there are sufficient data on both the individual ingredients and similar tested mixtures to adequately characterise the hazards of the mixtures. The following bridging principles may be used: dilution, batching, concentration of mixtures of the highest serious eye damage/eye irritation category, interpolation within one hazard category, substantially similar mixtures, and aerosols (see chapter 3.3.3.2 of UN, ~~2015~~2023).

~~84-87~~ The theory of additivity is used when data are available for all or only some of the ingredients, but not on the mixture as a whole. It assumes that each ingredient inducing serious eye damage or eye irritation contributes to the overall serious eye damage and/or irritation properties of the mixture in proportion to its potency and concentration. When applying such theory, considerations on the quality of the data of the ingredients is critical (e.g., data reported in Safety Data Sheets may be based on in vivo, in vitro or no test data). The mixture is classified as inducing serious eye damage and/or irritation when the sum of the concentrations of the relevant ingredients exceeds a pre-set cut-off value / concentration limit (see chapter 3.3.3.3 of UN, ~~2015~~2023).

Table 4.43. Overview of available (Q)SARs for serious eye damage and eye irritation. Note that this list is likely to be non-exhaustive and does not imply endorsement by OECD of any of the listed models for a particular prediction

Source	Applicability domain
Literature Models	
Basant et al. (2016)	
Verma and Matthews (2015a, 2015b)	Based on ADMET Predictor program
Solimeo et al. (2012)	N.A.
Gerner et al. (2005)	Based on physico-chemical values
Abraham et al. (2003)	Pure bulk liquids
Barratt (1995, 1997)	Neutral organic chemicals
Computerised Models – Freely available	
BfR rule base: included in the OECD QSAR Toolbox and Toxmatch, Toxtree, ToxPredict and Ambit*	EU New Chemicals (NONS) database, organic chemicals with no significant hydrolysis potential and purity > 95 %
PaDEL-DDPredictor (Liew and Yap, 2013)	Calculated by the model based on the range of descriptors
US FDA (Verma and Matthews, 2015c)	Based on physico-chemical properties
Computerised Models - Commercial	
ACD/Percepta	Organic chemicals
Derek Nexus	Organic chemicals and some metals
HazardExpert	Organic chemicals
Molcode	Organic chemicals
MultiCASE / Case Ultra	Organic chemicals
TopKat	Organic chemicals
Review papers	
Verheven et al. (2017)	N.A.
Bhatarai et al. (2016)	N.A.
Gallegos Saliner et al. (2006, 2008)	N.A.
Patlewicz et al. (2003)	N.A.

N.A. – Not Applicable. A detailed description of the above models is given in Appendix R.7.2-3 of the ECHA IR/CSA guidance 7a (ECHA, 2015, p. 252-257).* Underwent independent assessment.

4.8.3. Overview of non-testing data on serious eye damage and eye irritation

Module 8 – Non-testing data on serious eye damage and irritation	
Regulatory use (UN GHS classification)	<p><i>Substances</i></p> <p>Usually used as supporting information in a weight of evidence approach.</p> <p><i>Mixtures</i></p> <p>To be used for classification decision on serious eye damage (UN GHS Cat. 1), eye irritation (UN GHS Cat. 2 and UN GHS Cat. 2A and 2B), and no need for classification (UN GHS No Cat.).</p>
Validation & regulatory status	<p><i>Substances</i></p> <p>Validation and regulatory acceptance based on case-by-case.</p> <p><i>Mixtures</i></p> <p>Regulatory adopted approach.</p>
Potential role in the IATA	<p><i>Substances</i></p> <p>Non-testing methods are usually used as supporting information in a WoE approach, e.g., to support observations from available data from other in vivo test methods (Module 4) and to support in vitro results on serious eye damage and/or eye irritation (Modules 3 and 5). If further testing is required, information generated with this Module may be used for deciding how to address Part 3 i.e., to initiate a Top-Down or a Bottom-Up approach (Figure 2.1).</p> <p><i>Mixtures</i></p> <p>To be used when data are not available on the complete mixture or when data are available for all or some ingredients of the mixture. Furthermore, when validated in vitro test methods for serious eye damage and eye irritation are available, these may be used to generate data to classify the mixture instead of or in conjunction with the non-testing methods.</p>
Description	<p><i>Substances</i></p> <ul style="list-style-type: none"> - Analogue approaches (read-across, SARs, and grouping). - (Q)SARs. - Expert and other prediction systems that often include several (Q)SARs, expert rules and data. <p><i>Mixtures</i></p>

	<ul style="list-style-type: none"> - Bridging principles - Theory of additivity
Scientific basis incl. MoA	<p><i>Substances:</i></p> <p>Mainly correlative approaches based on the general assumption that substances with comparable structural properties have comparable serious eye damage and/or eye irritation properties. However this might change once the Adverse Outcome Pathway (AOP) project (OECD, 2013g) has made further progress or more (Q)SARs might become available built on mechanistically based high-throughput in vitro data.</p> <p><i>Mixtures:</i></p> <p>Bridging principles are used when there are sufficient data on both the individual ingredients and similar tested mixtures to adequately characterise the hazards of the mixtures. The following bridging principles may be used: based on dilution, batching, concentration of the highest corrosion/irritation category, interpolation within one hazard, substantially similar mixtures, and aerosols.</p> <p>The theory of additivity is used when data are available on the ingredients, but not on the mixture as a whole. It assumes that each ingredient inducing serious eye damage and/or eye irritation contributes to the overall serious eye damage and/or irritation properties of the mixture in proportion to its potency and concentration. The mixture is classified as inducing serious eye damage or eye irritation when the sum of the concentrations of the relevant ingredients exceeds a cut-off value / concentration limit (see chapter 3.3.3.3 of UN, 2015/2023).</p>
Strengths and weaknesses	<p><u>Strengths</u></p> <p><i>Substances and mixtures</i></p> <ul style="list-style-type: none"> - Ease of application. - Low cost. <p><u>Weaknesses</u></p> <p><i>Substances</i></p> <p>Results may be less relevant compared to experimental data, depending on the substance as well as the non-testing method and its underlying (model development/validation) dataset.</p> <p><i>Mixtures</i></p> <p>An impact assessment carried out by A.I.S.E. (Cazelle et al., 2014) showed that the use of the UN GHS theory of additivity for</p>

	<p>classification of detergent and cleaning products can result in the over-labelling of many products currently not requiring classification according to consistent animal, in vitro and human data. Similar findings were reported for agrochemical formulations (Corvaro et al., submitted2017)</p>
Applicability domain and limitations	<p><u>Applicability</u></p> <p><i>Substances</i></p> <p>Model-specific and needs to be defined in a (Q)SAR Model Reporting Format (QMRF). Also (Q)SAR Prediction Reporting Format (QPRF) are used to describe whether a prediction for a specific substance should be regarded as within the Applicability Domain or not.</p> <p>Application of these non-testing approaches is rather straight-forward for mono-constituent substances, whereas for multi-constituent substances, this only holds true if the composition of the substance is known (i.e. percentage of each of the discrete organic constituents) because then predictions can be performed on each constituent and the effect of the multi-constituent substance predicted by employing a dose addition approach.</p> <p>For Substances of Unknown and Variable Composition and Biologicals (UVCB), by definition, not all of the constituents are known with respect to their identity and/or their relative concentrations. (Q)SAR models and grouping approaches have, however, been employed on multi-constituent substances and UVCBs with partly unknown composition details for other endpoints than serious eye damage and irritation by accepting some uncertainty and assuming that all constituents of the considered UVCBs are represented by a few known constituents/groups of constituents, on which QSAR models or grouping approaches then could be employed.</p> <p><i>Mixtures</i></p> <p>The bridging principle is applicable to mixtures having data on both their individual ingredients and similar tested mixtures. The theory of additivity is applicable to mixtures that have data available for all or for some ingredients.</p> <p><u>Limitations</u></p> <p><i>Substances</i></p> <ul style="list-style-type: none"> - Limited applicability to the UN GHS classification scheme. - Applicability limited to the applicability domain of the model. <p><i>Mixtures</i></p> <p>Need to have sufficient data on similar tested mixtures as well as the ingredients of the mixture.</p>
Predictive capacity	<i>Substances</i>

	<p>Model-, domain- and context-specific. e.g. for ToxTree (rule-based) and MultiCase (statistics-based) computerized models, the prediction on the coverage of 80 substances was very low (reached 15 to 58%) (Geerts et al., 2017).</p> <p><i>Mixtures –</i></p> <p>Only limited data available. An impact assessment carried out by A.I.S.E. showed that the use of the UN GHS theory of additivity for classification of detergent and cleaning products can result in the over-labelling of many products currently not requiring classification according to consistent animal, in vitro and human experience data. Furthermore, a retrospective analysis of 225 agrochemical formulations indicated that, while overpredictive across categories, the use of the UN GHS theory of additivity for classification of agrochemical formulation can provide value for the identification of UN GHS No Cat. consistent with the classification based on in vivo animal test (Corvaro et al., submitted2017).</p>
--	---

4.9. Module 9: Phases and elements of Weight of Evidence (WoE) approaches

85-88. A weight of evidence determination means that all available and scientifically justified information bearing on the determination of hazard is considered together. In case of serious eye damage and eye irritation this includes structural information, information on physico-chemical parameters (e.g., pH, acid/alkaline reserve), information from category approaches (e.g., grouping) or read-across, (Q)SAR results, the results of suitable in vitro tests, relevant animal data, and human data. The quality and consistency of the data should be taken into account when weighing each piece of available information. Information such as study design, mechanism or mode of action, dose-effect relationships and biological relevance may be considered. Both positive and negative results can be assembled together in a single weight of evidence determination. Evaluation must be performed on a case-by-case basis and with expert judgement. In case of inconsistent data, the quality and relevance has to be carefully assessed in order to derive a conclusion. No formula can be presented for this analysis; a detailed explanation of the expert judgment used to overrule e.g. a single positive finding should accompany the derived conclusion.

86-89. A WoE approach may involve an assessment of the relative values/weights of different pieces of the available information that has been retrieved and gathered in previous steps (for an example see Hulzebos and Gerner, 2010). These weights/values can be assigned either in a more objective way by applying a formalised procedure (e.g., based on Bayesian logic, as in Rorije et al., 2013), by using meta-analyses (either weighted or unweighted) or by using expert judgement. Examples of tools to evaluate the quality include the Klimisch scores for experimental data (Klimisch et al., 1997) and Hill's criteria for evaluation of epidemiological data (Hill, 1965), as well as the JRC's ToxRTTool for scoring in vivo and in vitro data (Schneider et al., 2009). Under the GHS (UN, [20152023](#)), in sub-chapter 3.3.2-[2-9](#) a weight of evidence approach based on expert judgement is also recommended.

87-90. The weight given to the available evidence will be influenced by factors such as the quality of the data, consistency of results/data, nature and severity of effects, relevance of the information for the given regulatory endpoint. For each study/data, the relevance, reliability and adequacy for the purpose have to be considered. All available information that can contribute to the determination of classification for an endpoint is considered

Commented [EA6]: Update requested by the Netherlands (comment No. 45): A paragraph on the limited mechanistic information there is on eye irritation/serious eye damage. Considered relevant as the WoE assessment refers to mechanism or mode of action (paragraph 85).

Suggestion to include the following sentence in this paragraph: *There is limited mechanistic information available on eye irritation processes and therefore testing methods are generally an integration of different mechanisms leading to corneal opacity, the main effects leading to classification.*

together. In the following paragraphs a suggestion of the steps and elements of WoE is given.

4.9.1. Place/role of WoE in the IATA

~~88-91~~ 91. WoE should be carried out before any new prospective in vitro or in vivo testing is performed. A combination of physico-chemical information, (Q)SAR, read-across, grouping information and/or existing in vivo, in vitro and/or human data might be considered sufficient to conclude on serious eye damage and eye irritation effects.

4.9.2. Coverage of relevant sources of information

~~89-92~~ 92. The IATA specifies several types of existing information that can be used, with the condition that these are of sufficient quality. Structural information, physico-chemical properties, data on structurally-related chemicals obtained by read-across or grouping approaches, (Q)SAR modelling data, existing human and relevant laboratory animal data as well as in vitro data are listed. In the WoE analysis, the availability of specified types of data should be checked. The sources of those data obviously vary, ranging from clinical study reports, scientific publications, data from poison information centres, guideline tests, up to worker surveillance data of the chemical companies.

4.9.3. Assessment of data quality

~~90-93~~ 93. The quality of the data that is obtained for a WoE needs to be assessed, since the quality will contribute to the value/weight of each data element. In case the quality of a certain study is deemed to be inappropriate, it is recommendable not to consider those data in the WoE, but focus on other pieces of information which are of sufficient quality. Quality might be inappropriate e.g., due to negative outcome in the validation of the methodology, "non-adherence" to the relevant test guideline/method, lack of adequate controls, deficiencies in data reporting etc. Furthermore, quality may need to be evaluated based on expert judgement in case of e.g. incomplete or unavailable validation of a test method.

~~94-94~~ 94. The quality of the study, the method, the reporting of the results, and the conclusions that are drawn, must be evaluated carefully. Reasons why existing study data may vary in quality include the use of outdated test guidelines, the failure to characterise the test chemical properly (in terms of purity, physical characteristics, etc.) and the use of crude techniques/procedures that have since become refined, moreover, other reasons could be poor reporting of information and poor quality assurance.

~~92-95~~ 95. For many existing test chemicals, at least some of the available information could have been generated prior to the requirements of Good Laboratory Practice (GLP) and the standardisation of testing methods. While such information may still be usable, both the data and the methodology used must be evaluated in order to determine their reliability. Such an evaluation would ideally require an evidence-based evaluation i.e., a systematic and consistent evaluation following pre-defined, transparent and independently reviewed criteria before making decisions. These should always include justifications for the use of particular data sets on the basis of the criteria-based evaluation. For some test chemicals in contrast, information may be available from tests conducted according to OECD Test Guidelines (or other standards like CEN, ISO, ASTM, OSPAR methods, national standard methods), and in compliance with the principles of GLP or equivalent standards.

4.9.5. Adequacy of information

~~93-96~~ 96. Adequacy defines the usefulness of information for the purpose of hazard and risk assessment, in other words whether the available information allows clear decision-making about whether the test chemical induces (or not) serious eye damage and eye irritation and an adequate classification can be derived. The evaluation of adequacy of test results and

documentation for the intended purpose is particularly important for test chemicals where there may be (a number of) results available, but where some or all of them have not been carried out according to current standards. Where there is more than one study, the greatest weight is attached to the studies that are the most relevant and reliable. For each endpoint, robust summaries need to be prepared for the key studies. Sound scientific judgement is an important principle in considering the adequacy of information and determining the key study.

4.9.6. Non-testing data

(Q)SAR data

~~94-97.~~ It is important to distinguish between the proposed validity of the (Q)SAR model per se, and the reliability and adequacy of an individual (Q)SAR estimate (i.e., the application of the (Q)SAR model to a specific substance), and the appropriateness of the documentation (e.g., QMRF) associated with models and their predictions.

~~95-98.~~ Guidance on how to characterise (Q)SAR models according to the OECD (Q)SAR validation principles is provided in the OECD GD 69 (OECD, 2007). Other useful guidance has also been published to aid in determining how to use and report on QSAR models .

~~96-99.~~ The information in the QMRF and QPRF should be used when assessing whether a prediction is adequate for the purpose of classification and labelling and/or risk assessment. The assessment will also need to take into account the regulatory context. This means that the assessments of (Q)SAR validity (typically proposed in scientific publications) and (Q)SAR estimate reliability need to be supplemented with an assessment of the relevance of the prediction for the regulatory purposes, which includes an assessment of completeness, i.e., whether the information is sufficient to make the regulatory decision, and if not, what additional (experimental) information is needed. The decision will be taken on a case-by-case basis.

~~97-100.~~ (Q)SAR predictions may be gathered from databases (in which the predictions have already been generated and documented) or generated de novo through the available models.

Data obtained by grouping approaches

~~98-101.~~ Conclusions about the likely properties of a substance can also be based on the knowledge of the properties of one or more similar chemicals, by applying grouping methods.

~~99-102.~~ The corresponding OECD guidance document No. 194 provides information on the use of grouping of chemicals and read-across approaches (OECD, 2014b).

~~100-103.~~ ———As with (Q)SARs, grouping approaches can be used to indicate either the presence or the absence of an effect.

4.9.7. Existing human data

~~104-104.~~ ———The strength of the epidemiological evidence for specific health effects depends, among other things, on the type of analyses and on the magnitude and specificity of the response. Human data other than from epidemiological studies can be obtained from e.g., case reports, clinical studies, occupational disease registries or other occupational surveillance schemes and from poison centre information. In principle all types of toxic effects can be reported in such studies. Confidence in the findings is increased when comparable results are obtained in several independent studies on populations exposed to the same agent under different conditions. Other characteristics that support causal

associations are the presence of a dose-response relationship, a consistent correlation in time and (biological) plausibility, i.e., aspects covered by epidemiological criteria such as those described by Hill (1965), Fedak et al. (2015) and Lucas & McMichael (2005).

402.105. ——— A comprehensive guidance of both the evaluation and use of epidemiological evidence for risk assessment purposes is provided by Kryzanowski et al. (WHO, 2000).

403.106. ——— High quality human data may be considered as one of the strongest basis for classification and labelling decision making (subject to the ethical considerations relevant for the respective regulatory programme). However, when contradictory human and animal (OECD TG 405) data are available and WoE analysis including all other existing data and (Q)SAR profiling is not conclusive towards one or the other result, confirmatory in vitro testing should be performed.

404.107. ——— It is emphasised that testing with human volunteers for hazard identification is strongly discouraged for ethical reasons, but data from accidental human exposures, while not necessarily of the highest quality, can be used to support WoE conclusions.

4.9.8. Evaluation of consistency of the data

405.108. ——— The consistency of the existing data coming from various sources is crucial and should therefore be thoroughly evaluated in a WoE analysis. Consistent data which come from several studies/sources may be considered sufficient for regulatory purposes. In case the data elements are of comparable weight but give inconsistent evidence (e.g., (Q)SAR is positive and available limited human data is negative), usually WoE analysis will not be conclusive and prospective in vitro and/or in vivo testing will have to be conducted (Part 3 of the IATA). In case the weights of the individual pieces of evidence differ considerably, a WoE conclusion may be drawn according to the evidence carrying the highest weight. If high quality human (Module 1), in vitro (Module 3) and/or in vivo (Module 2) data are available, these should carry the highest weight in the WoE assessment.

4.9.10. Assessment of the coverage of relevant parameters and observations

406.109. ——— In a standard in vivo test guideline the required parameters/observations have been specified and often build the basis for decision making (e.g., classification and labelling for serious eye damage and eye irritation is mainly derived from the in vivo rabbit eye test). However, when taking together (in an integrating phase), it is not always possible to extract information equivalent to those parameters from non-testing data. Therefore, an important element of WoE is to consider to what extent the parameters and observations were addressed by each data element of the WoE.

4.9.11. Conclusions of WoE

407.110. ——— In the final analysis of the WoE, each data element will be characterised for its quality, relevance, coverage (e.g., serious eye damage and eye irritation) and associated uncertainty. The assessor would either decide to include or exclude the existing information based on these. When consistency is seen among "qualified" data elements, WoE may reach a conclusion that the relevant endpoint or information requirement has been sufficiently covered and further testing is not necessary. When on the other hand, insufficient information remains after the "non-qualified" data have been rejected/put aside and/or when the remaining information is inconsistent or contradictory, WoE would reach to a conclusion that the relevant endpoint or information requirement has not been sufficiently covered and further testing is necessary, depending

on the specific legal/regulatory framework, and inform on which test to conduct to fill the data gap.

~~108-111.~~ The WoE assessment needs to be transparently explained and documented to enable a logical flow leading to the decision/conclusion. An example for a simple approach to the documentation of the WoE is presented in Annex 1. Furthermore examples of evaluations are given for detergents and agrochemical mixtures in annex 2.

DRAFT 6 Nov 2023

5. REFERENCES

Abraham M.H., Hassanisadi M., Jalali-Heravi M., Ghafourian T., Cain W.S., Cometto-Muniz J.E. (2003) Draize rabbit eye test compatibility with eye irritation thresholds in humans: a quantitative structure-activity relationship analysis. *Toxicol. Sci.* 76, 384-91.

[Abo T, Yuki T, Xu R, Araki D, Takahashi Y, Sakaguchi H, Itagaki H. Expansion of the applicability domain for highly volatile substances on the Short Time Exposure test method and the predictive performance in assessing eye irritation potential. *J Toxicol Sci.* 2018;43\(7\):407-422. doi: 10.2131/jts.43.407. PMID: 29973473.](#)

Adriaens E., Barroso J., Eskes C., Hoffmann S., McNamee P., Alépée N., Bessou-Touya S., De Smedt A, de Wever B., Pfannenbecker U., Tailhardat M., Zuang V. (2014). Draize test for serious eye damage / eye irritation: importance of the endpoints evaluated with regard to UN GHS / EU CLP classification. *Archives of Toxicology* 88, 701-723.

[Adriaens E., Dhondt M.M., Remon J.P. \(2005\). Refinement of the Slug Mucosal Irritation test as an alternative screening test for eye irritation. *Toxicology In Vitro* 19, 79-89.](#)

Adriaens E., Bytheway H., De Wever B., Eschrich D., Guest R., Hansen E., Vanparys P., Schoeters G., Warren N., Weltens R., Whittingham A., Remon J.P. (2008). Successful prevalidation of the slug mucosal irritation test to assess the eye irritation potency of chemicals. *Toxicology In Vitro* 22, 1285-1296.

~~[Adriaens E., Dhondt M.M., Remon J.P. \(2005\). Refinement of the Slug Mucosal Irritation test as an alternative screening test for eye irritation. *Toxicology In Vitro* 19, 79-89.](#)~~

Adriaens E., Verstraelen S., Alépée N., Kandarova H., Drzewiecka A., Gruszka K., Guest R., Willoughby J.A., Van Rompay A.R. (2017a). CON4EI: Development of testing strategies for hazard identification and labelling for serious eye damage and eye irritation of chemicals. *Toxicology In Vitro* doi: 10.1016/j.tiv.2017.09.008.

Adriaens, E., Alépée, N., Kandarova, H., Drzewiecka, A., Gruszka, K., Guest, R., Willoughby, J.A. Sr., Verstraelen, S., Van Rompay, A.R. (2017b). CON4EI: Selection of Reference Chemicals for Hazard Identification and Labelling of Eye Irritating Chemicals. *Toxicol. In Vitro* 44, 44-48. <https://doi.org/10.1016/j.tiv.2017.06.001>

Adriaens, E., Willoughby, J. A., Meyer, B. R., Blakeman, L. C., Alepee, N., Drzewiecka, A., Gruszka, K., Guest, R., Kandarova, H., Verstraelen, S., & Van Rompay, A.R. (2017c). CON4EI: Short Time Exposure (STE) Test Method for Hazard Identification and Labelling of Eye Irritating Chemicals. *Toxicol. In Vitro* doi.org/10.1016/j.tiv.2017.08.002

Adriaens E, Guest R, Willoughby JA Sr, Fochtman P, Kandarova H, Verstraelen S, Van Rompay AR. (2017d). CON4EI: Slug Mucosal Irritation (SMI) test method for hazard identification and labelling of serious eye damaging and eye irritating chemicals. *Toxicol In Vitro*. doi: 10.1016/j.tiv.2017.08.020

[Adriaens E, Verstraelen S, Desprez B, Alépée N, Abo T, Bagley D, Hibatallah J, Mewes KR, Pfannenbecker U, Van Rompay AR. Overall performance of Bovine Corneal Opacity and Permeability \(BCOP\) Laser Light-Based Opacitometer \(LLBO\) test method with regard to solid and liquid chemicals testing. *Toxicol In Vitro.* 2021 Feb;70:105044. doi: 10.1016/j.tiv.2020.105044. Epub 2020 Oct 31. PMID: 33130054.](#)

A.I.S.E. (2006). Low Volume Eye Test (LVET): A.I.S.E. Submission to ECVAM, January 2006.

Alépée N., Barroso J., De Smedt A., De Wever B., Hibatallah J., Klaric M., Mewes K.R., Millet M., Pfannenbecker U., Tailhardat M., Templier M., McNamee P. (2015) Use of HPLC/UPLC Spectrophotometry for Detection of Formazan in In vitro Reconstructed Human Tissue (RhT)-Based Test Methods Employing the MTT-Reduction Assay to Expand their Applicability to Strongly Coloured Test Chemicals. *Toxicology In Vitro* 29, 741-761.

Alépée N., Leblanc V., Adriaens E., Grandidier M.H., Lelièvre D., Meloni M., Nardelli L., Roper C.S., Santirocco E., Toner F., Van Rompay A., Vinall J., Cotovio J. (2016a). Multi-laboratory validation of SkinEthic HCE test method for testing serious eye damage/eye irritation using liquid chemicals. *Toxicology In Vitro* 31, 43-53.

Alépée N., Adriaens E., Grandidier M.H., Meloni M., Nardelli L., Vinall C.J., Toner F., Roper C.S., Van Rompay A.R., Leblanc V., Cotovio J. (2016b). Multi-laboratory evaluation of SkinEthic HCE test method for testing serious eye damage/eye irritation using solid chemicals and overall performance of the test method with regard to solid and liquid chemicals testing. *Toxicology In Vitro* 34, 55-70.

[Alépée N., Adriaens E., Abo T., Bagley D., Desprez B., Hibatallah J., Mewes KR, Pfannenbecker U., Sala A., Van Rompay AR, Verstraelen S., McNamee P. \(2019a\). Development of a defined approach for eye irritation or serious eye damage for neat liquids based on Cosmetics Europe Analysis of in vitro RhCE and BCOP test methods. *Toxicology In Vitro* 59, 100-114. \[https://doi: 10.1016/j.tiv.2019.04.011\]\(https://doi.org/10.1016/j.tiv.2019.04.011\).](#)

[Alépée N., Adriaens E., Abo T., Bagley D., Desprez B., Hibatallah J., Mewes KR, Pfannenbecker U., Sala A., Van Rompay AR, Verstraelen S., McNamee P. \(2019b\). Development of a defined approach for eye irritation or serious eye damage for liquids, neat and in dilution, based on cosmetics Europe analysis of in vitro STE and BCOP test methods. *Toxicology In Vitro* 57, 154-163. doi: 10.1016/j.tiv.2019.02.019.](#)

[Alépée, N., Leblanc, V., Grandidier, MH., Teluob, S., Tagliati, V., Adriaens, E., Michaut, V. \(2020\) Development of the SkinEthic HCE Time-to-Toxicity test method for identifying liquid chemicals not requiring classification and labelling and liquids inducing serious eye damage and eye irritation. *Toxicol. In Vitro* 69, 104960.](#)

[Alépée, N., Leblanc, V., Grandidier, MH., Teluob, S., Viricel, A., Adriaens, E., Michaut, V. \(2021\). SkinEthic HCE Time-to-Toxicity on Solids: A test method for distinguishing chemicals inducing serious eye damage, eye irritation and not requiring classification and labelling. *Toxicol. in Vitro* 75, 105203.](#)

[Alépée, N., Grandidier, MH., Teluob, S., Amaral, F., Caviola, E., De Servi, B., Martin, S., Meloni, M., Nardelli, L., Padelou, C., Tagliati, V., Viricel, A., Adriaens, E., Michaut, V. \(2022\). Validation of the SkinEthic HCE Time-to-Toxicity test method for eye hazard classification of chemicals according to UN GHS. *Toxicol. in Vitro*, 80, 105319. <https://doi.org/10.1016/j.tiv.2022.105319>](#)

Bagley D.M., Bruner L.H., Silva O. de, Cottin M., O'Brien K.A.F., Uttley M., Walker A.P. (1992). An evaluation of five potential alternatives in vitro to the rabbit eye irritation test in vivo. *Toxicology In Vitro* 6, 136-149.

Bagley D.M., Cervin D., Harbell J.W. (1999). Assessment of the chorioallantoic membrane vascular assay (CAMVA) in the COLIPA in vitro eye irritation validation study. *Toxicology In Vitro* 13, 285-293.

Ball N., Cronin M.T.D., Shen J., Adenuga M.D., Blackburn K., Booth E.D., Bouhifd M., Donley E., Egnash L., Freeman J.J., Hastings C., Juberg D.R., Kleinsang A., Kleinstreuer N., Kroese E.D., Lee A.C.,

Luechtefeld T., Maertens A., Marty S., Naciff J.M., Palmer J., Pamies D., Penman M., Richarz A.-N., Russo D.P., Stuard S.B., Patlewicz G., van Ravenzwaay B., Wu S., Zhu H., Hartung T. (2016). Toward good read-across practice (GRAP) guidance. *ALTEX* 33,149-166.

Barratt M.D. (1995) The role of structure-activity relationships and expert systems in alternative strategies for the determination of skin sensitisation, skin corrosivity and eye irritation. *Altern. Lab. Anim.* 23, 111-22.

Barratt M.D. (1997) QSARs for the Eye Irritation Potential of Neutral Organic Chemicals. *Toxicology In Vitro* 11, 1-8.

Barroso J., Alépée N., Cole T., Eskes C., Freeman S.J., Liška R., McNamee P., Pfannenbecker U., Reus A.A., Rubingh C.M., Schaeffer M.W., Zuang V. (2014). The EURL ECVAM - Cosmetics Europe prospective validation study of Reconstructed human Cornea-like Epithelium (RhCE)-based test methods for identifying chemicals not requiring classification and labelling for serious eye damage/eye irritation: Validation Study Report. EUR 28125 EN; doi:10.2787/41680. Available at: <http://publications.jrc.ec.europa.eu/repository/handle/JRC100280>.

Barroso J., Pfannenbecker U., Adriaens E., Alépée N., Cluzel M., De Smedt A., Hibatallah J., Klaric M., Mewes K.R., Millet M., Templier M., McNamee P. (2017). Cosmetics Europe compilation of historical serious eye damage/eye irritation in vivo data analysed by drivers of classification to support the selection of chemicals for development and evaluation of alternative methods/strategies: the Draize eye test Reference Database (DRD). *Archives of Toxicology* 91, 521-547.

Bartok M., Gabel D., Zorn-Kruppa M., Engelke M. (2015). Development of an in vitro ocular test system for the prediction of all three GHS categories. *Toxicology In Vitro* 29, 72-80.

Berta A. (1992). Chapter 17: Enzymatic changes in tears in caustic injuries. In *Enzymology of the tears*. Boca Raton, CRC Press, p. 285-295.

Blazka M.E., Harbell J.W., Klauzner M., Raabe H., Kubilus J., Hsia F., Minerath B., Kotler M., Bagley D.M. (2000). Colgate-Palmolive's program to validate the EpiOcular™ human tissue construct model. *The Toxicologist* 54, 188.

Blazka M., Harbell J.W., Klausner M., Merrill J.C., Kubilus J., Kloss C., Bagley D.M. (2003). Evaluating the ocular irritation potential of 54 test articles using the EpiOcular human tissue construct model (OCL-200). *The Toxicologist* 72, 221.

Brantom P.G., Bruner L.H., Chamberlain M., Desilva O., Dupuis J., Earl L.K., Lovell D.P., Pape W.J.W., Uttley M., Bagley D.M., Baker F.W., Brachter M., Courtellemont P., Declercq L., Freeman S., Steiling W., Walker A.P., Carr G.J., Dami N., Thomas G., Harbell J., Jones P.A., Pfannenbecker U., Southee J.A., Tchong M., Argembeaux H., Castelli D., Clothier R., Esdaile D.J., Itigaki H., Jung K., Kasai Y., Kojima H., Kristen U., Larnicol M., Lewis R.W., Marenus K., Moreno O., Peterson A., Rasmussen E.S., Robles C., Stern M. (1997). A summary report of the COLIPA international validation study on alternatives to the Draize rabbit eye irritation test. *Toxicology In Vitro* 11, 141-179.

Cazelle E., Eskes C., Hermann M., Jones P., McNamee P., Prinsen M., Taylor H., Wijnands M.V.W. (2014). Suitability of histopathology as an additional endpoint to the isolated chicken eye test for classification of non-extreme pH detergent and cleaning products. *Toxicology In Vitro* 28, 657-666.

Cazelle E., Eskes C., Hermann M., Jones P., McNamee P., Prinsen M., Taylor H., Wijnands M.V.W. (2015). Suitability of the Isolated Chicken Eye Test for Classification of Extreme pH Detergents and Cleaning Products. *Toxicology In Vitro* 29, 609-616.

Cerven D., Moreno O. (1998). Bovine corneal opacity and permeability test validation as an alternative to the Draize eye irritation assay. In *Adv Anim Altern Saf Efficacy Test* (ed. H. Salem & S.A. Katz), pp. 261-267. Washington, DC, USA: Taylor & Francis.

Clippinger A.J., Hill E., Curren R., Bishop P. (2016). Bridging the gap between regulatory acceptance and industry use of non-animal methods. *ALTEX* 33, 453-458.

[Clippinger AJ, Raabe HA, Allen DG, Choksi NY, van der Zalm AJ, Kleinstreuer NC, Barroso J, Lowit AB. Human-relevant approaches to assess eye corrosion/irritation potential of agrochemical formulations. *Cutan Ocul Toxicol.* 2021 Jun;40\(2\):145-167. doi: 10.1080/15569527.2021.1910291. Epub 2021 Apr 20. PMID: 33830843.](#)

Cormier E.M., Hunter J.E., Billhimer W., May J., Farage M.A. (1995) Use of clinical and consumer eye irritation data to evaluate the low-volume eye test. *J. Toxicol. Cut. Ocul. Toxicol.* 14, 197-205.

Cormier E.M., Parker R.D., Henson C., Cruze L.W., Merritt A.K., Bruce R.D., Osborne R. (1996). Determination of the intra- and inter-laboratory reproducibility of the Low Volume Eye Test and its statistical relationship to the Draize tes. *Reg. Tox. Pharmac.* 23, 156-161.

Corvaro M., Gehen S., Macleod F., Metha J, (submitted). A retrospective analysis of in vivo skin and eye irritation and skin sensitisation studies with agrochemical formulations: setting the scene for development of alternative strategies.

DB-ALM (2013). Protocol no 71: Fluorescein Leakage Test. European Union Reference Laboratory for Alternatives to Animal Testing (EURL ECVAM), Ispra, Italy: Available at: <http://ecvam-dbalm.jrc.ec.europa.eu>

DB-ALM (2015). Protocol no 164: EpiOcular™ Eye Irritation Test – Summary. European Union Reference Laboratory for Alternatives to Animal Testing (EURL ECVAM), Ispra, Italy: Available at: <http://ecvam-dbalm.jrc.ec.europa.eu>

DB-ALM (2017). Protocol no 190: SkinEthic™ HCE Eye Irritation Test Liquid (EITL) – Summary. European Union Reference Laboratory for Alternatives to Animal Testing (EURL ECVAM), Ispra, Italy: Available at: <http://ecvam-dbalm.jrc.ec.europa.eu>

DB-ALM (2017). Protocol no 191: SkinEthic™ HCE Eye Irritation Test Solid (EITS) – Summary. European Union Reference Laboratory for Alternatives to Animal Testing (EURL ECVAM), Ispra, Italy: Available at: <http://ecvam-dbalm.jrc.ec.europa.eu>

Draize J.H., Woodard G., Calvery H.O. (1944). Methods for the study of irritation and toxicity of substances applied topically to the skin and mucous membranes. *Journal of Pharmacology and Experimental Therapeutics* 82, 377-390.

ECETOC (2002) ECETOC Monograph No.32, Use of Human Data in Hazard Classification for Irritation and Sensitisation. Available at: <http://www.ecetoc.org/publications>.

ECHA (2015). Skin corrosion / irritation, serious eye damage / eye irritation and respiratory tract corrosion / irritation. In: *Guidance on Information Requirements and Chemical Safety Assessment - Chapter R.7a:*

endpoint specific guidance, Version 4.1, October 2015, p. 168-235. Available at: http://echa.europa.eu/documents/10162/13632/information_requirements_r7a_en.pdf]

Commented [EA7]: https://echa.europa.eu/documents/10162/23249/09/ir_csa_r7a_r7-2_msc_rac_draft_en_201502_en.pdf/615a10ed-4e56-4787-a98b-2e2b1e71da8e

ESAC (2007). Statement on the conclusion of the ICCVAM retrospective study on organotypic in vitro assays as screening tests to identify potential ocular corrosives and severe eye irritants. Available at: <https://eurl-ecvam.jrc.ec.europa.eu/about-ecvam/scientific-advice-stakeholders-networks/ecvam-scientific-advisory-committee-esac/statements-opinions>.

ESAC (2009a). Statement on the use of existing Low Volume Eye Test (LVET) data for Weight-of-Evidence decisions on classification and labelling of cleaning products and their main ingredients. Available at: <https://eurl-ecvam.jrc.ec.europa.eu/about-ecvam/scientific-advice-stakeholders-networks/ecvam-scientific-advisory-committee-esac/statements-opinions>.

ESAC (2009b). Statement on the scientific validity of cytotoxicity/cell-function based in vitro assays for eye irritation testing. Available at: <https://eurl-ecvam.jrc.ec.europa.eu/about-ecvam/scientific-advice-stakeholders-networks/ecvam-scientific-advisory-committee-esac/statements-opinions>.

ESAC (2009c). ESAC Peer Review Panel consensus report the retrospective validation of cytotoxicity/cell function based in vitro assays. Available at: https://eurl-ecvam.jrc.ec.europa.eu/validation-regulatory-acceptance/docs-eye-irritation/1_Final%20draft%20PRP%20CBA.pdf.

ESAC (2014). ESAC Opinion on the EURL ECVAM Eye Irritation Validation Study (EIVS) on EpiOcular™ EIT and SkinEthic™ HCE and a related Cosmetics Europe study on HPLC/UPLC-spectrophotometry as an alternative endpoint detection system for MTT-formazan. ESAC Opinion No. 2014-03 of 17 November 2014; EUR 28173 EN; doi:10.2787/043697. Available at: <http://publications.jrc.ec.europa.eu/repository/handle/JRC103702>.

ESAC (2016a). ESAC Opinion on the SkinEthic™ Human Corneal Epithelium (HCE) Eye Irritation Test (EIT). ESAC Opinion No. 2016-02 of 24 June 2016; EUR 28175 EN; doi:10.2787/390390. Available at: <http://publications.jrc.ec.europa.eu/repository/handle/JRC103704>.

ESAC (2016b). ESAC Opinion on the Ocular Irritation® test method for prediction of serious eye damage/ eye irritation potential of chemicals. ESAC Opinion No. 2016-01 of 24 June 2016; EUR 28174 EN; doi:10.2787/991443. Available at: <http://publications.jrc.ec.europa.eu/repository/handle/JRC103703>.

Eskes C., Hoffmann S., Facchini D., Ulmer R., Wang A., Flego M., Vassallo M., Bufo M., van Vliet E., d'Abrosca F., Wilt N. (2014). Validation Study on the Ocular Irritation® Assay for Eye Irritation Testing. *Toxicology In Vitro* 28, 1046-1065.

EURL ECVAM (2008a). Fluorescein Leakage Assay Background Review Document as an Alternative Method for Eye Irritation Testing. Available under Validation & Regulatory Acceptance, Section Topical Toxicity - Eye Irritation at: <https://eurl-ecvam.jrc.ec.europa.eu>.

EURL ECVAM (2008b). Cytotoxicity/cell function based in vitro assays: Recommendations from the Validation Management Group. Available at: <https://eurl-ecvam.jrc.ec.europa.eu/validation-regulatory-acceptance/docs-eye-irritation/VMG%20CBA%20recommendations%20-%20final.pdf>.

Fedak K.M., Bernal A., Capshaw Z.A., Gross S. (2015). Applying the Bradford Hill criteria in the 21st century: how data integration has changed causal inference in molecular epidemiology. *Emerging Themes in Epidemiology* 12, 14. doi: 10.1186/s12982-015-0037-4.

Freeberg F.E., Griffith J.F., Bruce R.D., Bay P.H.S. (1984) Correlation of animal test methods with human experience for household products. *J. Toxicol. Cut. Ocul. Toxicol.* 1, 53-64.

Freeberg F.E., Hooker D.T., Griffith J.F. (1986a) Correlation of animal eye test data with human experience for household products: an update. *J. Toxicol. Cut. Ocul. Toxicol.* 5, 115-23.

Freeberg F.E., Nixon G.A., Reer P.J., Weaver J.E., Bruce R.D., Griffith J.F., Sanders III L.W. (1986b) Human and rabbit eye responses to chemical insult. *Fund Appl. Toxicol.* 7, 626-34.

Furukawa M., Sakakibara T., Itoh K., Kawamura K., Sasaki S., Matsuura M. (2015). Histopathological evaluation of the ocular-irritation potential of shampoos, make-up removers and cleansing foams in the bovine corneal opacity and permeability assay. *J. Toxicol. Pathol.* 28, 243-248.

Gallegos Saliner A., Patlewicz G., Worth A.P. (2006). Review of literature-based models for skin and eye irritation and corrosion. JRC report EUR 22320 EN. European Chemicals Bureau, Ispra, Italy.

Gallegos Saliner A., Patlewicz G., Worth A.P. (2008) A Review of (Q)SAR Models for Skin and Eye Irritation and Corrosion. *QSAR Comb. Sci.* 27, 49-59.

Geerts L., Adriaens E., Alépée N., Guest R., Willoughby JA. Sr, Kandarova H., Drzewiecka A., Fochtman P., Verstraelen S., Van Rompay A.R. (2017) CON4EI: Evaluation of QSAR models for hazard identification and labelling of eye irritating chemicals. *Toxicol In Vitro* doi: 10.1016/j.tiv.2017.09.004

Gerner I., Liebsch M., Spielmann H. (2005). Assessment of the Eye Irritating Properties of Chemicals by Applying Alternatives to the Draize Rabbit Eye Test: The Use of QSARs and In vitro Tests for the Classification of Eye Irritation. *Alternatives to Laboratory Animals* 33, 215-237.

Gershbein L.L., McDonald J.E. (1977). Evaluation of the corneal irritancy of test shampoos and detergents in various animal species. *Food and Cosmetics Toxicology* 15, 131-134.

Griffith J.F., Nixon G.A., Bruce R.D., Reer P.J., Bannan E.A. (1980). Dose-response studies with chemical irritants in the albino rabbit eye as basis for selecting optimum testing conditions for predicting hard to the human eye. *Toxicology and Applied Pharmacology* 55, 501-513.

Guo X., Yang X.F., Yang Y., Hans R., Cai J.H., Xue J.Y., Tan X.H., Xie X.P., Xiong X.K., Huang J.M. (2010). Prediction of ocular irritancy of 26 chemicals and 26 cosmetic products with isolated rabbit eye (IRE) test. *Biomed. Environ. Sci.* 25, 359-366.

Harbell J.W., Koontz S.W., Lewis R.W., Lovell D., Acosta D. (1997) IRAG Working Group 4. Cell cytotoxicity assays. *Food and Chemical Toxicology* 35, 79-126.

Harbell, J.W., Mun G., Curren, R.D. (2006), Application of histological evaluation to enhance the bovine opacity and permeability (BCOP) assay. *Toxicologist* 90: 326. Available at: http://www.iivs.org/pages/publication_view.php?doc_id=135.

Hayashi K., Mori T., Abo T., Koike M., Takahashi Y., Sakaguchi H., Nishiyama N. (2012a). A tiered approach combining the short time exposure (STE) test and the bovine corneal opacity and permeability (BCOP) assay for predicting eye irritation potential of chemicals. *J. Toxicol. Sci.* 37, 269-280.

Hayashi K., Mori T., Abo T., Ooshima K., Hayashi T., Komano T., Takahashi Y., Sakaguchi H., Takatsu A., Nishiyama N. (2012b). Two-stage bottom-up tiered approach combining several alternatives for identification of eye irritation potential of chemicals including insoluble or volatile substances. *Toxicology In Vitro* 26, 1199-1208.

Hill A.B. (1965). The environment and disease: association or causation? *Proc. R. Soc. Med.* 58, 295-300.

Hoffmann, S., Saliner, A.G., Paltewicz, G., Eskes, C., Zuang, V., Worth, A.P., 2008. A feasibility study developing an integrated testing strategy assessing skin irritation potential of chemicals *Toxicol. Lett* 180, 9-20.

[Huang AJ, Tseng SC. Corneal epithelial wound healing in the absence of limbal epithelium. Invest Ophthalmol Vis Sci. 1991 Jan;32\(1\):96-105. PMID: 1702774.](#)

Hulzebos E., Gerner I. (2010). Weight factors in an Integrated Testing Strategy using adjusted OECD principles for (Q)SARs and extended Klimisch codes to decide on skin irritation classification. *Regulatory Toxicology and Pharmacology* 58,131-144.

ICCVAM (2006). Test Method Evaluation Report: In vitro Ocular Toxicity Test Methods for Identifying Ocular Severe Irritants and Corrosives. NIH Publication No.: 07-4517. Available: <http://ntp.niehs.nih.gov/pubhealth/evalatm/test-method-evaluations/ocular/in-vitro/tmer/index.html>.

ICCVAM (2010a). ICCVAM Test Method Evaluation Report: Recommendations to Discontinue Use of the Low Volume Eye Test for Ocular Safety Testing. NIH Publication No. 10-7515. Available at: https://ntp.niehs.nih.gov/iccvam/docs/ocutox_docs/lvet/front.pdf.

ICCVAM (2010b). Test Method Evaluation Report: Current Validation Status of In vitro Test Methods Proposed for Identifying Eye Injury Hazard Potential of Chemicals and Products. National Institutes of Health Publication Number 10-7553A. National Toxicology Program, North Carolina, USA. Available at: <http://ntp.niehs.nih.gov/pubhealth/evalatm/test-method-evaluations/ocular/in-vitro-test-methods/index.html>.

ICCVAM (2010c). ICCVAM Test Method Evaluation Report: Current validation status of a proposed in vitro testing strategy for U.S. Environmental Protection Agency ocular hazard classification and labeling of antimicrobial cleaning products. NIH Publication No. 10-7513. Available at: <http://ntp.niehs.nih.gov/pubhealth/evalatm/test-method-evaluations/ocular/amcp/index.html>.

ICCVAM (2013). Short Time Exposure (STE) Test Method Summary Review Document, NIH. Available at: http://www.ntp.niehs.nih.gov/iccvam/docs/ocutox_docs/STE-SRD-NICEATM-508.pdf.

ILSI Health and Environmental Sciences Institute (1996). Replacing the Draize eye irritation test : scientific background and research needs. *J. Toxicol. Cut. & Ocular Toxicol.* 15, 211-234.

Ishii S., Ishii K., Nakadate M., Yamasaki K. (2013). Correlation study in skin and eye irritation between rabbits and humans based on published literatures. *Food Chem. Toxicol.* 55, 596-601.

Jester J.V., Li, H.F., Petroll, W.M., Parker, R.D., Cavanagh, H.D., Carr, G.J., Smith, B., Maurer, J.K. (1998). Area and Depth of Surfactant Induced Corneal Injury Correlates with Cell Death. *Invest. Ophthalmol. Vis. Sci.* 39, 922-936.

Kandarova H., Letasiova S., Adriaens E., Guest R., Willoughby J.A., Drzewiecka A., Gruszka K., Alépée N., Verstraelen S., Van Rompay A.R. (2017a). CON4EI: EpiOcular™ Eye Irritation Test (EpiOcular™ EIT) for hazard identification and labelling of eye irritating chemicals. *Toxicology In Vitro*. doi: 10.1016/j.tiv.2017.07.002.

Kandarova H., Letasiova S., Adriaens E., Guest R., Willoughby J.A. Sr, Drzewiecka A., Gruszka K., Alépée N., Verstraelen S., Van Rompay A.R.(2017b). CON4EI: CONSortium for in vitro Eye Irritation testing strategy - EpiOcular™ time-to-toxicity (EpiOcular ET-50) protocols for hazard identification and labelling of eye irritating chemicals. *Toxicol In Vitro*. doi: 10.1016/j.tiv.2017.08.019.

Katoh M., Hamajima F., Ogasawara T., Hata K. (2013). Establishment of a new in vitro test method for evaluation of eye irritancy using a reconstructed human corneal epithelial model, LabCyte CORNEA-MODEL. *Toxicology In Vitro* 27, 2184-2192.

Klimisch H.J., Andreae M., Tillmann U. (1997) A Systematic Approach for Evaluating the Quality of Experimental Toxicological and Ecotoxicological Data. *Regulatory Toxicology and Pharmacology* 25, 1-5.

Kojima H, Hayashi K., Sakaguchi H., Omori T., Otoizumi T., Sozu T., Kuwahara H., Hayashi T., Sakaguchi M., Toyoda A., Goto H., Watanabe S., Ahiko K., Nakamura T., Morimoto T.. (2013). Second-Phase Validation of Short Time Exposure Tests for Assessment of Eye Irritation Potency of Chemicals. *Toxicology In Vitro* 27, 1855-1869.

Kolle S.N., Kandárová H., Wareing B., van Ravenzwaay B., Landsiedel R. (2011). In-house validation of the EpiOcular(TM) eye irritation test and its combination with the bovine corneal opacity and permeability test for the assessment of ocular irritation. *Altern. Lab. Anim.* 39, 365-387.

Kolle S.N., Moreno, M.C.R., Mayer, W., van Cott, A., van Ravenzwaay, B., Landsiedel, R. (2015). The EpiOcular™ Eye Irritation Test is the Method of Choice for In vitro Eye Irritation Testing of Agrochemical Formulations: Correlation Analysis of EpiOcular™ Eye Irritation Test and BCOP Test Data to UN GHS, US EPA and Brazil ANIVSA Classifications. *ATLA* 43, 1-18.

Kolle S.N., Sauer U.G., Moreno M.C., Teubner W., Wohlleben W., Landsiedel R. (2016). Eye irritation testing of nanomaterials using the EpiOcular™ eye irritation test and the bovine corneal opacity and permeability assay. *Part. Fibre Toxicol.* 13, 18. doi: 10.1186/s12989-016-0128-6.

Lemp M.A. (1974). Cornea and sclera. *Arch Ophthalmol* 92, 158-170.

Lenoir J., Claerhout I., Kestelyn R., Komp A., Remon J.P., Adriaens E. (2011). The slug mucosal irritation (SMI) assay: development of a screening tool for the evaluation of ocular discomfort caused by shampoos. *Toxicol In Vitro* 25, 1919-1925.

Lewis R.W., McCall J.C., Botham P.A. (1993). A comparison of two cytotoxicity tests for predicting the ocular irritancy of surfactants. *Toxicology In Vitro* 7, 155-158.

Liew C.Y., Yap C.W. (2013). QSAR and predictors of eye and skin effects. *Mol. Inform.* 32, 281-290.

[Lim SE, Ha SJ, Jang WH, Jung KM, Jung MS, Yeo KW, Kim JS, Jeong TC, Kang MJ, Kim SY, Lee SH, Ko KY, Kim TS, Park KS, Bae S, Lim KM. Me-too validation study for in vitro eye irritation test with 3D-reconstructed human cornea epithelium. MCTT HCETM. *Toxicol In Vitro*. 2019 Mar;55:173-184. doi: 10.1016/j.tiv.2018.12.003. Epub 2018 Dec 17. PMID: 30572010.](#)

Lucas R.M., McMichael A.J. (2005). Association or causation: evaluating links between "environment and disease". *Bull World Health Organ.* 83, 792-795.

Luechtefeld T., Maertens A., Russo D.P., Rovida C., Zhu H., Hartung T. (2016). Analysis of Draize eye irritation testing and its prediction by mining publicly available 2008-2014 REACH data. *ALTEX.* 33, 123-134.

Luepke N.P. (1985). Hen's egg chorioallantoic membrane test for irritation potential. *Food and Chemical Toxicology* 23, 287-291.

Marzulli F.N., Ruggles D.I. (1973). Rabbit eye irritation test: collaborative study. *J. Ass. Off. Analyt. Chem.* 56, 905-914.

Maurer J.K., Parker R.D., Jester J.V. (2002). Extent of corneal injury as the mechanistic basis for ocular irritation: key findings and recommendations for the development of alternative assays. *Reg. Tox. Pharmacol.* 36,106-117.

McCulley J.P. (1987). Chemical injuries. In *Cornea, scientific foundations and clinical practice* (Smolin G, Thoft RA Ed.). Boston, Little, Brown p. 527-542.

Mewes K.R., Engelke M., Zorn-Kruppa M., Bartok M., Tandon R., Brandner J.M., Petersohn D. (in press), In vitro eye irritation testing using the open source reconstructed hemicornea - a ring trial. *ALTEX*, 2017 Jan. 14. Epub ahead of print. doi: 10.14573/altex.

Mosmann T. (1983). Rapid Colorimetric Assay for Cellular Growth and Survival: Application to Proliferation and Cytotoxicity Assays. *J. Immunol. Methods* 65, 55-63.

Muir C.K., Flower C., Abb N.J. van (1983). A novel approach to the search for in vitro alternatives to in vivo eye irritancy testing. *Toxicology Letters* 18, 1-5.

OECD (2005). Series on Testing and Assessment No.34. Guidance Document the Validation and International Acceptance of New or Updated Test Methods for Hazard Assessment. Organisation for Economic Cooperation and Development, Paris. Available at: <http://www.oecd.org/env/testguidelines>.

~~OECD (2007). Series on Testing and Assessment No. 69. Guidance Document on the Validation of (Quantitative) Structure-Activity Relationships [(Q)SAR] Models. Organisation for Economic Cooperation and Development, Paris. Available at: <http://www.oecd.org/env/testguidelines>.~~

~~OECD (2017). Series on Testing and Assessment No. 160. Guidance Document on "The Bovine Corneal Opacity and Permeability (BCOP) and Isolated Chicken Eye (ICE) Test Methods: Collection of Tissues for Histological Evaluation and Collection of Data on Non-Severe Irritants (Second Edition)". OECD, Paris. Available at: <http://www.oecd.org/env/testguidelines>.~~

~~OECD (2012a). Guideline for Testing of Chemicals No. 405. Acute Eye Irritation/Corrosion. Organisation for Economic Cooperation and Development, Paris. Available at: -~~

~~OECD (2012b). Guidelines for Testing of Chemicals No. 460. Fluorescein Leakage Test Method for Identifying Ocular Corrosives and Severe Irritants. Organisation for Economic Co-operation and Development, Paris. Available at: <http://www.oecd.org/env/testguidelines>.~~

OECD (2012e2012a). Series on Testing and Assessment No. 180: Streamlined summary document related to the Fluorescein Leakage (FL) test method for identification of ocular corrosives and severe irritants (TG 460). Organisation for Economic Co-operation and Development, Paris. Available at: <http://www.oecd.org/env/testguidelines>.

OECD (2012d2012b). Draft Test Guideline on the Cytosensor Microphysiometer Test Method: An in vitro Method for Identifying Ocular Corrosive and Severe Irritant Chemicals as well as Chemicals not Classified as Ocular Irritants. Available at: <http://www.oecd.org/env/testguidelines>.

~~OECD (2013a). Guidelines for Testing of Chemicals No. 437. Bovine Corneal Opacity and Permeability Test Method for Identifying i) Chemicals Inducing Serious Eye Damage. Organisation for Economic Cooperation and Development, Paris. Available at: <http://www.oecd.org/env/testguidelines>.~~

~~OECD (2013b). Guideline for Testing of Chemicals No. 438. Isolated Chicken Eye Test Method for Identifying i) Chemicals Inducing Serious Eye Damage and ii) Chemicals Not Requiring Classification. Organisation for Economic Cooperation and Development, Paris. Available at: .~~

OECD (2013ac2013b). Series on Testing and Assessment No. 189: Streamlined summary document supporting OECD Test Guideline 437 on the Bovine Corneal Opacity and Permeability for eye irritation/corrosion. Organisation for Economic Co-operation and Development, Paris. Available at: <http://www.oecd.org/env/testguidelines>.

~~OECD (2013d). Series on Testing and Assessment No. 188: Streamlined summary document supporting OECD Test Guideline 438 on the Isolated Chicken Eye for eye irritation / corrosion. Organisation for Economic Co-operation and Development, Paris. Available at: <http://www.oecd.org/env/testguidelines>.~~

OECD (2013be2013e). Guideline for the Testing of Chemicals No. 122. Determination of pH, Acidity and Alkalinity. OECD, Paris Available at: <http://www.oecd.org/env/testguidelines>.

~~OECD (2014a). Series on Testing and Assessment No. 203: Guidance Document on Integrated Approaches to Testing and Assessment of Skin Irritation/Corrosion. Organisation for Economic Co-operation and Development, Paris. Available at: <http://www.oecd.org/env/testguidelines>.~~

~~OECD (2014a). *Guidance Document on the Validation of (Quantitative) Structure-Activity Relationship [(Q)SAR] Models*. OECD Series on Testing and Assessment, No. 69. OECD Publishing, Paris. <https://doi.org/10.1787/9789264085442-en>.~~

OECD (2014b). Series on Testing and Assessment No. 194. Guidance on Grouping of Chemicals, second edition. Organisation for Economic Cooperation and Development, Paris. Available at: <https://doi.org/10.1787/9789264274679-en>~~<http://www.oecd.org/env/testguidelines>.~~

OECD (2014c). Series on Testing and Assessment No 211. Guidance Document for Describing Non-guideline In Vitro Test Methods. Organisation for Economic Cooperation and Development, Paris. Available at: <https://doi.org/10.1787/9789264274730-en>~~<http://www.oecd.org/env/testguidelines>.~~

~~OECD (2015a). Guideline for the Testing of Chemicals No. 491. Short Time Exposure In vitro Test Method for Identifying i) Chemicals Inducing Serious Eye Damage and ii) Chemicals Not Requiring Classification for Eye Irritation or Serious Eye Damage. Organisation for Economic Cooperation and Development, Paris. Available at: <http://www.oecd.org/env/testguidelines>.~~

~~OECD (2015b). Guideline for the Testing of Chemicals No. 492. Reconstructed human Cornea-like Epithelium (RhCE) test method for identifying chemicals not requiring classification and labelling for eye irritation or serious eye damage. Organisation for Economic Cooperation and Development, Paris. Available at: <http://www.oecd.org/env/testguidelines>.~~

OECD (2015~~b~~^c). Series on Testing and Assessment No. 216. ~~Peformance~~^{Performance} Standards for the Assessment of Proposed Similar or Modified In vitro Reconstructed human Cornea-like Epithelium (RhCE) test methods for eye hazard. Organisation for Economic Cooperation and Development, Paris. Available at: <http://www.oecd.org/env/testguidelines>.

OECD (~~2016~~^c). Series on Testing and Assessment ~~No.n.~~^{No.} 255. Guidance Document on the Reporting of Defined Approaches and Individual Information Sources to be used within Integrated Approaches to Testing and Assessment (IATA). Organisation for Economic Cooperation and Development, Paris. Available at: [https://one.oecd.org/document/env/jm/mono\(2016\)28/en/pdf](https://one.oecd.org/document/env/jm/mono(2016)28/en/pdf)~~http://www.oecd.org/env/testguidelines~~.

~~OECD (2017). Guideline for the Testing of Chemicals No. 492. Reconstructed human Cornea-like Epithelium (RhCE) test method for identifying chemicals not requiring classification and labelling for eye irritation or serious eye damage. Organisation for Economic Cooperation and Development, Paris. Available at: <http://www.oecd.org/env/testguidelines>.~~

~~OECD (2013b/2018a). Guideline for Testing of Chemicals No. 438. Isolated Chicken Eye Test Method for Identifying i) Chemicals Inducing Serious Eye Damage and ii) Chemicals Not Requiring Classification. Organisation for Economic Cooperation and Development, Paris. Available at: <http://www.oecd.org/env/testguidelines>.~~

~~OECD (2013d/2018b). Series on Testing and Assessment No. 188. Streamlined summary document supporting OECD Test Guideline 438 on the Isolated Chicken Eye for eye irritation / corrosion. Organisation for Economic Co-operation and Development, Paris. Available at: <http://www.oecd.org/env/testguidelines>.~~

~~OECD (2017/2018c). Series on Testing and Assessment No. 160. Guidance Document on "The Bovine Corneal Opacity and Permeability (BCOP) and Isolated Chicken Eye (ICE) Test Methods: Collection of Tissues for Histological Evaluation and Collection of Data on Non-Severe Irritants (Second Edition). OECD, Paris. Available at: -~~

~~OECD (2017). *Guidance Document on an Integrated Approach on Testing and Assessment (IATA) for Skin Corrosion and Irritation*, OECD Series on Testing and Assessment, No. 203, OECD Publishing, Paris, <https://doi.org/10.1787/9789264274693-en>.~~

~~OECD (2018a). Series on Testing and Assessment No. 160. Guidance Document on "The Bovine Corneal Opacity and Permeability (BCOP) and Isolated Chicken Eye (ICE) Test Methods: Collection of Eye Tissues for Histological Evaluation and Collection of Data on Non-Severe Irritants (Second Edition). OECD, Paris. Available at: [https://one.oecd.org/document/env/jm/mono\(2011\)45/rev1/en/pdf](https://one.oecd.org/document/env/jm/mono(2011)45/rev1/en/pdf).~~

~~OECD (2018b). *Peer Review Report on Validation status of the LabCyte CORNEA-MODEL24 EYE IRRITATION TEST*, OECD Series on Testing and Assessment No.282. ENV Publications, Organisation for Economic Cooperation and Development, Paris.~~

[OECD \(2019\). Series on Testing and Assessment No. 188: Streamlined summary document supporting OECD Test Guideline 438 on the Isolated Chicken Eye for eye irritation / corrosion. Organisation for Economic Co-operation and Development, Paris. Available at: http://www.oecd.org/env/testguidelines.](http://www.oecd.org/env/testguidelines)

[OECD \(2021\). Test No. 494: Vitrigel-Eye Irritancy Test Method for Identifying Chemicals Not Requiring Classification and Labelling for Eye Irritation or Serious Eye Damage. OECD Guidelines for the Testing of Chemicals, Section 4, OECD Publishing, Paris, https://doi.org/10.1787/9f20068a-en.](https://doi.org/10.1787/9f20068a-en)

[OECD \(2022a\). Test No. 492B: Reconstructed Human Cornea-like Epithelium \(RHCE\) Test Method for Eye Hazard Identification. OECD Guidelines for the Testing of Chemicals, Section 4, OECD Publishing, Paris, https://doi.org/10.1787/0d603916-en.](https://doi.org/10.1787/0d603916-en)

[OECD \(2022b\). Test No. 467: Defined Approaches for Serious Eye Damage and Eye Irritation. OECD Guidelines for the Testing of Chemicals, Section 4, OECD Publishing, Paris, https://doi.org/10.1787/28fe2841-en.](https://doi.org/10.1787/28fe2841-en)

[OECD \(2023a\). Test No. 405: Acute Eye Irritation/Corrosion. OECD Guidelines for the Testing of Chemicals, Section 4, OECD Publishing, Paris, https://doi.org/10.1787/9789264185333-en.](https://doi.org/10.1787/9789264185333-en)

[OECD \(2023b\). Test No. 437: Bovine Corneal Opacity and Permeability Test Method for Identifying i\) Chemicals Inducing Serious Eye Damage and ii\) Chemicals Not Requiring Classification for Eye Irritation or Serious Eye Damage. OECD Guidelines for the Testing of Chemicals, Section 4, OECD Publishing, Paris, https://doi.org/10.1787/9789264203846-en.](https://doi.org/10.1787/9789264203846-en)

[OECD \(2023c\). Test No. 438: Isolated Chicken Eye Test Method for Identifying i\) Chemicals Inducing Serious Eye Damage and ii\) Chemicals Not Requiring Classification for Eye Irritation or Serious Eye Damage. OECD Guidelines for the Testing of Chemicals, Section 4, OECD Publishing, Paris, https://doi.org/10.1787/9789264203860-en.](https://doi.org/10.1787/9789264203860-en)

[OECD \(2023d\). Test No. 460: Fluorescein Leakage Test Method for Identifying Ocular Corrosives and Severe Irritants. OECD Guidelines for the Testing of Chemicals, Section 4, OECD Publishing, Paris, https://doi.org/10.1787/9789264185401-en.](https://doi.org/10.1787/9789264185401-en)

[OECD \(2023e\). Test No. 491: Short Time Exposure In Vitro Test Method for Identifying i\) Chemicals Inducing Serious Eye Damage and ii\) Chemicals Not Requiring Classification for Eye Irritation or Serious Eye Damage. OECD Guidelines for the Testing of Chemicals, Section 4, OECD Publishing, Paris, https://doi.org/10.1787/9789264242432-en.](https://doi.org/10.1787/9789264242432-en)

[OECD \(2023f\). Test No. 492: Reconstructed human Cornea-like Epithelium \(RhCE\) test method for identifying chemicals not requiring classification and labelling for eye irritation or serious eye damage. OECD Guidelines for the Testing of Chemicals, Section 4, OECD Publishing, Paris, https://doi.org/10.1787/9789264242548-en.](https://doi.org/10.1787/9789264242548-en)

[OECD \(2023g\). Test No. 496: In vitro Macromolecular Test Method for Identifying Chemicals Inducing Serious Eye Damage and Chemicals Not Requiring Classification for Eye Irritation or Serious Eye Damage. OECD Guidelines for the Testing of Chemicals, Section 4, OECD Publishing, Paris, https://doi.org/10.1787/970e5cd9-en.](https://doi.org/10.1787/970e5cd9-en)

Pape, W.J.W., Pfannenbecker, U. & Hoppe, U. (1987). Validation of the red blood cell test system as in vitro assay for the rapid screening of irritation potential of surfactants. *Molecular Toxicology* 1, 525-536.

- Pape W.J.W., Hoppe U. (1990). Standardization of an in vitro red blood cell test for evaluating the acute cytotoxic potential of tensides. *Arzneimittelforschung* 40, 498-502.
- Pape W.J.W., Pfannenbecker U., Argembeaux H., Bracher M., Esdaile D.J., Hagino S., Kasai Y., Lewis R.W. (1999). COLIPA validation project on in vitro eye irritation tests for cosmetic ingredients and finished products (phase I): the red blood cell test for the estimation of acute eye irritation potentials. Present status. *Toxicology In Vitro* 13, 343-354.
- Patlewicz G., Rodford R., Walker J.D. (2003). Quantitative structure-activity relationships for predicting skin and eye irritation. *Environ. Toxicol. Chem.* 22, 1862-1869.
- Pfister R.R. (1983). Chemical injuries of the eye. *Ophtamology* 90, 1246-1253.
- Piehl M., Gilotti A., Donovan A., DeGeorge G., Cerven D. (2010). Novel cultured porcine corneal irritancy assay with reversibility endpoint. *Toxicology In Vitro* 24, 231-239.
- Piehl M., Carathers M., Soda R., Cerven D., DeGeorge G. (2011). Porcine Corneal Ocular Reversibility Assay (PorCORA) predicts ocular damage and recovery for global regulatory agency hazard categories. *Toxicology In Vitro* 25, 1912-1918.
- Prinsen M.K. (2006). The Draize eye test and in vitro alternatives, a left-handed marriage? *Toxicology In Vitro* 20, 78-81.
- Prinsen M.K., Schipper M.E.I., Wijnands M.V.W. (2011). Histopathology in the Isolated Chicken Eye Test and comparison of different stainings of the cornea. *Toxicology In Vitro* 25, 1475-1479.
- Roggeband R., York M., Pericoi M., Braun W. (2000). Eye irritation responses in rabbit and man after single applications of equal volumes of undiluted model liquid detergent products. *Food and Chemical Toxicology* 38, 727-734.
- Rorije E., Aldenberg T., Buist H., Kroese D., Schüürmann G. (2013). The OSIRIS weight of evidence approach: ITS for skin sensitization. *Regul. Toxicol. Pharmacol.* 67, 146-156.
- Saito K., Miyazawa M., Nukada Y., Ei K., Abo T., Sakaguchi H. (2015). Predictive performance of the Short Time Exposure test for identifying eye irritation potential of chemical mixtures. *Toxicology In Vitro* 29, 617-20.
- Sakaguchi H., Ota N., Omori T., Kuwahara H., Sozu T., Takagi Y., Takahashi Y., Tanigawa K., Nakanishi M., Nakamura T., Morimoto T., Wakuri S., Okamoto Y., Sakaguchi M., Hayashi T., Hanji T., Watanabe S. (2011). Validation Study of the Short Time Exposure (STE) Test to Assess the Eye Irritation Potential of Chemicals. *Toxicology In Vitro* 25, 796-809.
- Schaeffer M., Barroso J., Worth A., Zuang V. (2014). Chapter 4. Serious eye damage and eye irritation. In: *Alternative methods for regulatory toxicology – a state-of-the-art review*. EURL-ECVAM Report EUR 26797 EN. Luxembourg: Publications Office of the European Union. Available at: http://publications.jrc.ec.europa.eu/repository/bitstream/JRC91361/echa_jrc_sla_report_public_05-09-14_withcover%20ipo.pdf.
- Schneider K., Schwarz M., Burkholder I., Kopp-Schneider A., Edler L., Kinsner-Ovaskainen A., Hartung T., Hoffmann S. (2009). "ToxRTTool", a new tool to assess the reliability of toxicological data. *Toxicol. Lett.* 189, 138-144.

Scott L., Eskes C., Hoffmann S., Adriaens E., Alépée N., Bufo M., Clothier R., Facchini D., Faller C., Guest R., Harbell J., Hartung T., Kamp H., Le Varlet B., Meloni M., McNamee P., Osborne R., Pape W., Pfannenbecker U., Prinsen M., Seaman C., Spielman H., Stokes W., Trouba K., Van den Berghe C., Van Goethem F., Vassallo M., Vinardell P., Zuang V. (2010). A proposed eye irritation testing strategy to reduce and replace in vivo studies using Bottom-Up and Top-Down approaches. *Toxicology In Vitro* 24,1-9.

Settivari R.S., Amado R.A., Corvaro M., Visconti N.R., Kan L., Carney E.W., Boverhof D.R., Gehen S.C. (2016). Tiered application of the neutral red release and EpiOcular™ assays for evaluating the eye irritation potential of agrochemical formulations. *Regul. Toxicol. Pharmacol.* 81, 407-420.

Solimeo R., Zhang J., Kim M., Sedykh A., Zhu H. (2012). Predicting Chemical Ocular Toxicity Using a Combinatorial QSAR Approach. *Chem. Res. Toxicol.* 25, 2763-2769.

Spielmann H., Gerner I., Kalweit S., Moog R., Wirnsberger T., Krauser K., Kreiling R., Kreuzer H., Luepke N.P., Miltenburger H.G., Mueller N., Muermann P., Pape W., Siegemund B., Spengler J., Steiling W., Wiebel, F.J. (1991). Interlaboratory assessment of alternatives to the Draize eye irritation test in Germany. *Toxicology In Vitro* 5, 539-542.

Steiling W., Bracher M., Courtellemont P., Silva O. de (1999). The HET-CAM, a useful in vitro assay for assessing the eye irritation properties of cosmetic formulations and indigents. *Toxicology In Vitro* 13, 375-384.

Stürer A., Seidel C., Sauer O., Koch I., Zilker T., Hermanns-Clausen M., Hruba K., Hüller G., Tutdibi E., Heppner H.J., Desel H. (2010). Poisons Centres' Data for Expert Judgement within Classification, Labelling and Packaging Regulation: Solid Household Automatic Dishwashing Products do not Cause Serious Eye Damage. *Clinical Toxicology* 48, 245.

Takahashi Y., Koike M., Honda H., Ito Y., Sakaguchi H., Suzuki H., Nishiyama N. (2008). Development of the Short Time Exposure (STE) Test: an In vitro Eye Irritation Test Using SIRC Cells. *Toxicology In Vitro* 22, 760-770.

Tandon R., Bartok M., Zorn-Kruppa M., Brandner J.M., Gabel D., Engelke M. (2015). Assessment of the eye irritation potential of chemicals: A comparison study between two test methods based on human 3D hemi-cornea models. *Toxicology In Vitro* 30, 561-568.

Tsakovska I., Netzeva T., Worth A.P. (2005) Evaluation of (Q)SARs for the prediction of Eye Irritation/Corrosion Potential - physico-chemical exclusion rules. JRC Report EUR 21897 EN, 42pp. European Chemicals Bureau, Ispra, Italy. Available at: https://eurl-ecvam.jrc.ec.europa.eu/laboratories-research/predictive_toxicology/doc/Evaluation_of_Eye_Irritation_QSARs.pdf

Tsakovska I., Gallegos Saliner A., Netzeva T., Pavan M., Worth A.P. (2007) Evaluation of SARs for the prediction of eye irritation/corrosion potential - structural inclusion rules in the BfR decision support system. *SAR QSAR Environ Res* 18, 221-235.

UN (2015/2023). United Nations Globally Harmonized System of Classification and Labelling of Chemicals (GHS). ST/SG/AC.10/30/Rev.6/10, ~~Sixth~~ Tenth Revised Edition, New York and Geneva: United Nations. Available at: <https://unece.org/sites/default/files/2023-07/GHS%20Rev10e.pdf>
http://www.unece.org/trans/danger/publi/ghs/ghs_rev06/06files_e.html.

US EPA (2015). Use of an alternate testing framework for classification of eye irritation potential of EPA pesticide products. Office of Pesticide Programs, US Environmental Protection Agency, Washington DC, 20460, 3-2-2015. Available at: http://www.epa.gov/sites/production/files/2015-05/documents/eye_policy2015update.pdf.

U.S. CPSC. 2015b. Test for Eye Irritants. 16 C.F.R. §1500.42. Available at: <https://www.gpo.gov/fdsys/pkg/CFR-2016-title16-vol2/pdf/CFR-2016-title16-vol2-sec1500-42.pdf>

Van Goethem, F., Hansen, E., Sysmans, M., De Smedt, A., Vanparys, P., Van Gompel, J., (2010). Development of a new opacimeter for the bovine corneal opacity and permeability (BCOP) assay. *Toxicol. in Vitro* 24, 1854–1861.

Van Rompay A.R., Alépée N., Nardelli L., Hollanders K., Leblanc V., Drzewiecka A., Gruszka K., Guest R., Kandarova H., Willoughby J.A., Verstraelen S., Adriaens E. (2017). CON4EI: SkinEthic™ Human Corneal Epithelium Eye Irritation Test (SkinEthic™ HCE EIT) for hazard identification and labelling of eye irritating chemicals. *Toxicology In Vitro*. doi: 10.1016/j.tiv.2017.06.012.

Verma R.P., Matthews E.J. (2015a). Estimation of the chemical-induced eye injury using a weight-of-evidence (WoE) battery of 21 artificial neural network (ANN) c-QSAR models (QSAR-21): part I: irritation potential. *Regul. Toxicol. Pharmacol.* 71, 318-330.

Verma R.P., Matthews E.J. (2015b). Estimation of the chemical-induced eye injury using a Weight-of-Evidence (WoE) battery of 21 artificial neural network (ANN) c-QSAR models (QSAR-21): part II: corrosion potential. *Regul. Toxicol. Pharmacol.* 71, 331-336.

Verma R.P., Matthews E.J. (2015c). An in silico expert system for the identification of eye irritants. *SAR QSAR Environ. Res.* 26, 383-395.

Verstraelen, S., Maglennon, Hollanders, K., Boonen, F., Adriaens, E., Alépée, N., Drzewiecka, A., Gruszka, K., Kandarova, H., Willoughby J.A., Guest, R., Schofield, J., Van Rompay, A.R. (2017). CON4EI: Bovine Corneal Opacity and Permeability (BCOP) Test for Hazard Identification and Labelling of Eye Irritating Chemicals. *Toxicol. In Vitro* 44, 122-133

Weil C.S., Scala A. (1971). Study of intra- and inter- laboratory variability in the results of rabbit eye and skin irritation tests. *Toxicology and Applied Pharmacology* 19, 276-360.

WHO (2000). Working Group report on Evaluation and use of epidemiological evidence for environmental health risk assessment: WHO Guideline Document. *Environm. Health Perspect.* 108, 997-1002.

Wilhelmus K.R. (2001). The Draize eye test. *Survey of Ophthalmology* 45, 493-515.

Yamaguchi H., Kojima H., Takezawa T. (2016). Predictive performance of the Vitrigel-eye irritancy test method using 118 chemicals. *J. Appl. Toxicol.* 36, 1025-1037.

Young J.R., How M.J., Walker A.P., Worth W.M.H. (1988). Classification as corrosive or irritant to skin of preparations containing acidic or alkaline substances, without testing on animals. *Toxicology In Vitro* 2, 19-26.

Zhu H., Bouhifd M., Kleinstreuer N., Kroese E.D., Liu Z., Luechtefeld T., Pamies D., Shen J., Strauss V., Wu S., Hartung T. (2016). Supporting read-across using biological data. *ALTEX* 33,167-182.

114 | ENV/JM/MONO(2017)15/REV1

Zorn-Kruppa M., Houdek P., Wladykowski E., Engelke M., Bartok M., Mewes K.R., Moll I., Brandner J.M. (2014). Determining the Depth of Injury in Bioengineered Tissue Models of Cornea and Conjunctiva for the Prediction of All Three Ocular GHS Categories. PLoS One 9, e114181.

[Peer Review Report on Validation status of the MCTT HCE™ EYE IRRITATION TEST, unpublished.](#)

DRAFT 6 Nov 2023

Unclassified

DRAFT 6 Nov 2023

ANNEX I – EXAMPLE OF MATRIX FOR WEIGHT OF EVIDENCE ANALYSES

For those modules having available data, entries are filled in the respective cases. For the rest of the entries, NA shall be indicated in column 2. It is recommended to use short and conclusive wording. For assessment of the evidence, refer to the Part 2 of this guidance document. Note that WoE should be assessed before any new experimental data is generated.

Module	Title of document/full reference; or data not available (N)	Study Result and/or positive or negative evidence obtained	Data quality according to the Klimisch score when appropriate*	Adequacy and relevance, short statement	Coverage of relevant parameters/ observations Yes/ NO	Consistency with other information	Conclusive remark**
1. Existing human data							
2. Existing data on skin corrosion							
3. In vivo animal study							
4. In vitro data from OECD adopted test methods							
5. Other animal data from non OECD adopted test methods							
6. Other data from non-OECD adopted							

alternative test methods							
7. Physico-chemical properties							
8. Non-testing methods ((Q)SAR, grouping, bridging & additivity approaches)							
Overall conclusion	<p>1. WoE allows decision/assessment of the potential of the test chemical to induce serious eye damage and eye irritation. The substance should be classified as UN GHS No Cat., Cat. 2 (2A or 2B), Cat. 1, or</p> <p>2. WoE does not allow decision/assessment of the potential of the test chemical to induce serious eye damage and eye irritation. Recommendation or specification of the most appropriate additional testing strategy to be undertaken.</p>						

*) An electronic tool supporting the quality assessment of in vivo and vitro data through the application of consistent criteria leading to scored results has been developed by EURL ECVAM (described in Schneider et al., 2009). The ToxRTTool can be downloaded from the EURL ECVAM page: <https://eurl-ecvam.jrc.ec.europa.eu/about-ecvam/archive-publications/toxrtool>**) For example: "This data (any entry except 3 and 4) is consistent with the existing in vitro studies".

***) For example: "The existing human data suggest that the substance is an eye irritant. Due to poor reporting of this data, and low quality in terms of exposure information, the data is inconclusive, and has a low weight in the final evaluation. "

ANNEX II – EXAMPLES OF WEIGHT OF EVIDENCE EVALUATIONS

Disclaimer: the examples presented below do not imply acceptance or endorsement by any Member Country or OECD. They are intended only to provide an illustration on how individual information sources may be reported and combined in a WoE approach to derive a final classification.

Example 1: Weight of evidence analyses for classification of a Soluble Liquid (SL) Agrochemical formulation, DD-001, for effects on eyes

	Full Reference	Study Result	Data quality Klimisch score	Adequacy and relevance	Coverage of relevant parameters/ observations Yes/ NO	Consistency	Conclusive remark
1. Existing human data	Not available)						
2. In vivo animal study	Non available						
3. In vitro data from OECD adopted test methods	Arthur, 2015. In vitro Eye irritation test (Neural red release method)	The relative tissue viability (mean) in the Epi Ocular was 67%. The mixture does not require classification for effect on eyes according to the predication model	1	Key study conducted according to GLP. OECD 492 allows discrimination between materials not requiring classification from those requiring classification (Cat. 2/ Cat.1)	Yes	Consistent with existing in vitro studies.	Key data. Data supports that the mixture does not require classification as eye irritant. Proposed classification: GHS not classified
4. Other animal data from non OECD	Non available.						

Unclassified

adopted test methods							
5. Other data from non-OECD adopted alternative methods	Arthur, 2015. In vitro Eye irritation test (Neural red release method)	The EC50 in the NRR assay was 630. The mixture did not show eye irritation potentially both according to original interpretation criteria. (Reader, 1989) and proposed revised criteria (Settivari, 2016).	2	Internal screening, non-GLP compliant but performed in a GLP facility in the spirit of GLP. Supportive information, limitation due to lack of predictivity for GHS cat 2 agrochemicals formulations.	Yes	Consistent with existing in vitro studies (low cytotoxicity)	Supportive data. Data supports that the mixture does not require classification as eye irritant. Proposed classification: GHS not classified.
6. Existing data on skin corrosion	Smith, 2011. Acute skin irritation study in the White Zealand Rabbit	Not corrosive nor irritant to the skin. Mean scores (at 24, 48, 72 hours): -Erythema: 0.7, 1.0, 1.3 -Oedema: 0.0, 0.3, 0.0 Recovery by day 7	1	GLP compliant. Study confirms low skin irritation potential	Yes	Consistent with other in vitro evidence	Supportive data. Effects on skin except for skin corrosion do not allow assessment for effects on eyes. Data supports that the mixture does not require classification as severe eye irritant.
7. Physico-chemical properties	Acosta, 2001. Determination of pH, acidity and alkalinity measurement according to OECD 122	pH is 5.2 is therefore not pH-extreme	1	Supportive information because pH alone does not allow assessment of the eye irritancy	Yes		Supportive data. Data supports that the mixture does not require classification as severe eye irritant.
8. Non-testing methods ((Q)SAR, grouping, bridging & additivity approaches)	Chatfield, 2014. Additivity approach, requested in the European assessment report, Part C, confidential information	GHS or CLP classification for all ingredients (2 active substances and 4 co-formulants) is available from the corresponding MSDS. There are no GHS cat 1	Not applicable as Klimisch score is applicable to assessing the reliability of toxicological studies	Supportive information. Usable for Classification purposes in EU	Yes	Consistent with existing in vitro studies	Key data. Data supports that the mixture does not require classification as eye irritant.

		classified ingredients and 2 ingredients (surfactants) classified as GHS cat 2A, accounting for a total of 6.72% w/w of the mixture composition. No classification for eye irritation is triggered according to GHS criteria (UN, 2015).					Proposed classification: GHS not classified
Overall conclusion	<p>No human data are available.</p> <p>pH and skin effects do not lead to a direct UN GHS Cat. 1 classification.</p> <p>In vitro data on two independent cytotoxicity based assays indicate that the test item has low cytotoxic potential and classification is not required.</p> <p>Non testing data (additivity approach based on concentration thresholds), support that classification is not required.</p> <p>In conclusion, a WoE evaluation of the consistency, quality and relevance of all available data allows a decision on the eye irritation/serous eye damage potential of the Agrochemical formulation, DD-001. DD-001 should not be classified for eye hazards.</p>						

Note: This example has been developed only to illustrate how the classification of an untested mixture could be derived and justified. It does not contain any recommendation for a testing strategy. However, the BCOP has very recently been included in a testing strategy for antimicrobial cleaning products (AMCPs) under the U.S. EPA classification and labelling system (Clippinger et al., 2016).

Example 2: Weight of evidence analyses for classification of a Hand Dish Washing Liquid W07 for effects on eyes

	Full Reference	Study result	Data quality Klimisch score	Adequacy and relevance	Coverage of relevant parameters/observations Yes/NO	Consistency	Conclusive remarks
Existing human data on company-owned mixture W07	Poison Control Centre data collected over a 12 months period	9 cases of mild to moderate eye effects only were reported out of all sold products*. In the cases where follow-up information was available, all ocular effects were fully reversible within a few days. *This is an example, in reality the number of cases will need to be identified relative to the number of products sold in a specific geographical area.	Not applicable to Poison Control Centre data as Klimisch score is applicable to assessing the reliability of toxicological studies.	Supportive information. Limitation due to unknown dose and exposure duration. No criteria for C&L based on human data.	No, not in every case all relevant parameters are covered (e.g. exposure conditions, detailed tissue effects).	Consistent with existing in vitro studies and other human experience, which identify the hand dish washing liquid W07 as inducing fully reversible ocular effects.	Supportive data.
Existing human data on similar mixtures	MAGAM II Multicentre multinational prospective, study of human eye exposures reported to poisons centres, over a 24 months	28 reported cases related to hand dish washing liquids: mild to moderate but no severe eye irritation after exposure. In	Not applicable to Poison Control Centre data as Klimisch score is applicable to assessing the reliability of	Supportive information. Scoring based on Poison Control Centre severity scoring system complemented by	Although not in every case all relevant parameters are available (e.g. exposure conditions), tissue observations are conducted typically by an ophthalmologist and	Consistent with existing in vivo and in vitro studies, which identify the hand dish washing liquid W07 as inducing fully reversible ocular	Supportive data.

	period	the cases where follow-up information was available, all ocular effects were fully reversible within a few days.	toxicological studies.	MAGAM reported symptoms. No criteria for C&L based on human data. Information provided as a product category containing different products vs. an individual named product.	reported in a standardized way.	effects	
In vitro data on eye irritation corrosion	Isolated Chicken Eye Test OECD 438 with histopathology as an additional endpoint, 2015	No Prediction can be Made based on a combination of the endpoint categories of II;II;III. This combination of endpoint categories is much lower than those used to identify classification as Cat. 1. Not identified as UN GHS Cat. 1 based on criteria developed by Cazelle et al. (2014) for histopathological evaluation of non-pH-extreme detergents and cleaning products.	1	Key and supportive study conducted according to GLP	Yes	Consistent with existing in vitro studies and human experience data which does not identify the hand dish washing liquid W07 as a UN GHS Cat. 1.	Key and supportive data. OECD 438 study with histopathology as an additional endpoint.
In vitro data on eye irritation corrosion	Reconstructed human Cornea-like Epithelium (RhCE) Test Method OECD 492, 2016	Tissue viability in the EpiOcular™ EIT was 45 %, identifying that the mixture requires classification for effect on eyes	1	Key study conducted according to GLP. Study allows judgement on need or no need for classification. OECD 492 allows discrimination	Yes	Consistent with existing in vitro studies and human experience data	Key data.

				between materials not requiring classification from those requiring classification (Cat. 2/ Cat. 1).			
In vitro data on eye irritation corrosion	Bovine Corneal Opacity and Permeability Test OECD 437, 2015	No Prediction can be Made based on In Vitro Irritancy Score (IVIS) of 10.3. The IVIS is far below the threshold of 55.1 for classification as Cat. 1	1	Key study conducted according to GLP. Study allows judgement on severity of effects but not persistence of effects and it does not allow identification of Cat. 2 specifically.	Yes	Consistent with existing in vitro studies and human experience data	Key data.
In vitro data on skin irritation	In Vitro Skin Irritation: Reconstructed Human Epidermis (RHE) Test Method OECD 439, 2014	Tissue viability in EpiSkin™ test method was 75 %, identifying that the tested mixture does not require classification for skin irritation.	1	Study confirms low skin irritation potential.	Yes	Consistent with existing in vitro studies and human experience data	Supportive data. Effects on skin except for skin corrosion do not allow assessment for effects on eyes.
Physico-chemical properties	Determination of pH, acidity and alkalinity measurement according to OECD 122	pH is 6.0, W07 is therefore not pH-extreme	1	Supportive information because pH alone does not allow assessment of the eye irritancy.	Yes		Supportive data.
Overall Conclusion	<p>Human data indicates only mild to moderate and fully reversible effects.</p> <p>In vitro data indicates classification required but mixture not identified as UN GHS Cat. 1.</p> <p>pH and skin effects do not indicate corrosive effects.</p> <p>In conclusion, a WoE evaluation of the consistency, quality and relevance of all available data allows a decision on the eye irritation/serious eye damage potential of the Hand Dish Washing Liquid W07. The Hand Dish Washing Liquid W07 should be classified as UN GHS Cat. 2.</p>						

Note: This example has been developed only to illustrate how the classification of an untested mixture could be derived and justified. It does not contain any recommendation for a testing strategy. However, the BCOP has very recently been included in a testing strategy for antimicrobial cleaning products (AMCPs) under the U.S. EPA classification and labelling system (Clippinger et al., 2016).

DRAFT 6 Nov 2023

